

Lecture 8: More MCMC: Metropolis-Hastings, Gibbs and Blocking

Merlise Clyde

September 26



Metropolis-Hastings (MH)

- Metropolis requires that the proposal distribution be symmetric



Metropolis-Hastings (MH)

- Metropolis requires that the proposal distribution be symmetric
- Hastings (1970) generalizes Metropolis algorithms to allow asymmetric proposals - aka Metropolis-Hastings or MH $q(\theta^* | \theta^{(s)})$ does not need to be the same as $q(\theta^{(s)} | \theta^*)$



Metropolis-Hastings (MH)

- Metropolis requires that the proposal distribution be symmetric
- Hastings (1970) generalizes Metropolis algorithms to allow asymmetric proposals - aka Metropolis-Hastings or MH $q(\theta^* | \theta^{(s)})$ does not need to be the same as $q(\theta^{(s)} | \theta^*)$
- propose $\theta^* | \theta^{(s)} \sim q(\theta^* | \theta^{(s)})$



Metropolis-Hastings (MH)

- Metropolis requires that the proposal distribution be symmetric
- Hastings (1970) generalizes Metropolis algorithms to allow asymmetric proposals - aka Metropolis-Hastings or MH $q(\theta^* | \theta^{(s)})$ does not need to be the same as $q(\theta^{(s)} | \theta^*)$
- propose $\theta^* | \theta^{(s)} \sim q(\theta^* | \theta^{(s)})$
- Acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / q(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / q(\theta^{(s)} | \theta^*)} \right\}$$



Metropolis-Hastings (MH)

- Metropolis requires that the proposal distribution be symmetric
- Hastings (1970) generalizes Metropolis algorithms to allow asymmetric proposals - aka Metropolis-Hastings or MH $q(\theta^* | \theta^{(s)})$ does not need to be the same as $q(\theta^{(s)} | \theta^*)$
- propose $\theta^* | \theta^{(s)} \sim q(\theta^* | \theta^{(s)})$
- Acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / q(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / q(\theta^{(s)} | \theta^*)} \right\}$$

- adjustment for asymmetry in acceptance ratio is key to ensuring convergence to stationary distribution!



Special cases

- Metropolis



Special cases

- Metropolis
- Independence chain



Special cases

- Metropolis
- Independence chain
- Gibbs samplers



Special cases

- Metropolis
- Independence chain
- Gibbs samplers
- Metropolis-within-Gibbs



Special cases

- Metropolis
 - Independence chain
 - Gibbs samplers
 - Metropolis-within-Gibbs
 - combinations of the above!



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)
- Replace Gaussian by a Student-t with low degrees of freedom



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)
- Replace Gaussian by a Student-t with low degrees of freedom
- transformations of θ



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)
- Replace Gaussian by a Student-t with low degrees of freedom
- transformations of θ



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)
- Replace Gaussian by a Student-t with low degrees of freedom
- transformations of θ



Independence Chain

- suppose we have a good approximation $\tilde{\pi}(\theta | y)$ to $\pi(\theta | y)$
- Draw $\theta^* \sim \tilde{\pi}(\theta | y)$ *without* conditioning on $\theta^{(s)}$
- acceptance probability

$$\min \left\{ 1, \frac{\pi(\theta^*) \mathcal{L}(\theta^*) / \tilde{\pi}(\theta^* | \theta^{(s)})}{\pi(\theta^{(s)}) \mathcal{L}(\theta^{(s)}) / \tilde{\pi}(\theta^{(s)} | \theta^*)} \right\}$$

- what happens if the approximation is really accurate?
- probability of acceptance is ≈ 1
- Important caveat for convergence: tails of the posterior should be at least as heavy as the tails of the posterior (Tweedie 1994)
- Replace Gaussian by a Student-t with low degrees of freedom
- transformations of θ



Gibbs Sampler

special case of Blocked MH



Gibbs Sampler

special case of Blocked MH

- proposal distribution q_k for the k th block is the **full conditional distribution** for $\theta_{[k]}$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) = \frac{\pi(\theta_{[k]}, \theta_{[-k]} \mid y)}{\pi(\theta_{[-k]} \mid y)} \propto \pi(\theta_{[k]}, \theta_{[-k]} \mid y)$$



Gibbs Sampler

special case of Blocked MH

- proposal distribution q_k for the k th block is the **full conditional distribution** for $\theta_{[k]}$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) = \frac{\pi(\theta_{[k]}, \theta_{[-k]} \mid y)}{\pi(\theta_{[-k]} \mid y)} \propto \pi(\theta_{[k]}, \theta_{[-k]} \mid y)$$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) \propto \mathcal{L}(\theta_{[k]}, \theta_{[-k]})\pi(\theta_{[k]}, \theta_{[-k]})$$



Gibbs Sampler

special case of Blocked MH

- proposal distribution q_k for the k th block is the **full conditional** distribution for $\theta_{[k]}$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) = \frac{\pi(\theta_{[k]}, \theta_{[-k]} \mid y)}{\pi(\theta_{[-k]} \mid y)} \propto \pi(\theta_{[k]}, \theta_{[-k]} \mid y)$$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) \propto \mathcal{L}(\theta_{[k]}, \theta_{[-k]})\pi(\theta_{[k]}, \theta_{[-k]})$$

$$\min \left\{ 1, \frac{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^* \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})}{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^{(s-1)} \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})} \right\}$$

Gibbs Sampler

special case of Blocked MH

- proposal distribution q_k for the k th block is the **full conditional** distribution for $\theta_{[k]}$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) = \frac{\pi(\theta_{[k]}, \theta_{[-k]} \mid y)}{\pi(\theta_{[-k]} \mid y)} \propto \pi(\theta_{[k]}, \theta_{[-k]} \mid y)$$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) \propto \mathcal{L}(\theta_{[k]}, \theta_{[-k]})\pi(\theta_{[k]}, \theta_{[-k]})$$

$$\min \left\{ 1, \frac{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^* \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})}{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^{(s-1)} \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})} \right\}$$

- acceptance probability is always 1!



Gibbs Sampler

special case of Blocked MH

- proposal distribution q_k for the k th block is the **full conditional** distribution for $\theta_{[k]}$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) = \frac{\pi(\theta_{[k]}, \theta_{[-k]} \mid y)}{\pi(\theta_{[-k]} \mid y)} \propto \pi(\theta_{[k]}, \theta_{[-k]} \mid y)$$

$$\pi(\theta_{[k]} \mid \theta_{[-k]}, y) \propto \mathcal{L}(\theta_{[k]}, \theta_{[-k]})\pi(\theta_{[k]}, \theta_{[-k]})$$

$$\min \left\{ 1, \frac{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^*, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^* \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})}{\pi(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})\mathcal{L}(\theta_{[<k]}^{(s)}, \theta_{[k]}^{(s-1)}, \theta_{[>k]}^{(s-1)})/q_k(\theta_{[k]}^{(s-1)} \mid \theta_{[<k]}^{(s)}, \theta_{[>k]}^{(s-1)})} \right\}$$

- acceptance probability is always 1!
- even though joint distribution is messy, full conditionals may be (conditionally) conjugate and easy to sample from!



Univariate Normal Example

Model

$$\begin{aligned} Y_i \mid \mu, \sigma^2 &\stackrel{iid}{\sim} N(\mu, 1/\phi) \\ \mu &\sim N(\mu_0, 1/\tau_0) \\ \phi &\sim \text{Gamma}(a/2, b/2) \end{aligned}$$



Univariate Normal Example

Model

$$\begin{aligned} Y_i \mid \mu, \sigma^2 &\stackrel{iid}{\sim} N(\mu, 1/\phi) \\ \mu &\sim N(\mu_0, 1/\tau_0) \\ \phi &\sim \text{Gamma}(a/2, b/2) \end{aligned}$$

- Joint prior is a product of independent Normal-Gamma



Univariate Normal Example

Model

$$\begin{aligned} Y_i \mid \mu, \sigma^2 &\stackrel{iid}{\sim} N(\mu, 1/\phi) \\ \mu &\sim N(\mu_0, 1/\tau_0) \\ \phi &\sim \text{Gamma}(a/2, b/2) \end{aligned}$$

- Joint prior is a product of independent Normal-Gamma
- Is $\pi(\mu, \phi \mid y_1, \dots, y_n)$ also a Normal-Gamma family?



Full Conditional for the Mean

The full conditional distributions $\mu | \phi, y_1, \dots, y_n$

$$\begin{aligned}\mu | \phi, y_1, \dots, y_n &\sim N(\hat{\mu}, 1/\tau_n) \\ \hat{\mu} &= \frac{\tau_0 \mu_0 + n\phi\bar{y}}{\tau_0 + n\phi} \\ \tau_n &= \tau_0 + n\phi\end{aligned}$$



Full Conditional for the Precision

$$\begin{aligned}\phi \mid \mu, y_1, \dots, y_n &\sim \text{Gamma}(a_n/2, b_n/2) \\ a_n &= a + n \\ b_n &= b + \sum_i (y_i - \mu)^2\end{aligned}$$



Full Conditional for the Precision

$$\begin{aligned}\phi \mid \mu, y_1, \dots, y_n &\sim \text{Gamma}(a_n/2, b_n/2) \\ a_n &= a + n \\ b_n &= b + \sum_i (y_i - \mu)^2\end{aligned}$$

$$\mathbb{E}[\phi \mid \mu, y_1, \dots, y_n] = \frac{(a + n)/2}{(b + \sum_i (y_i - \mu)^2)/2}$$



Full Conditional for the Precision

$$\begin{aligned}\phi \mid \mu, y_1, \dots, y_n &\sim \text{Gamma}(a_n/2, b_n/2) \\ a_n &= a + n \\ b_n &= b + \sum_i (y_i - \mu)^2\end{aligned}$$

$$\mathbb{E}[\phi \mid \mu, y_1, \dots, y_n] = \frac{(a + n)/2}{(b + \sum_i (y_i - \mu)^2)/2}$$

What happens with a non-informative prior i.e

$a = b = \epsilon$ as $\epsilon \rightarrow 0$?



Full Conditional for the Precision

$$\begin{aligned}\phi \mid \mu, y_1, \dots, y_n &\sim \text{Gamma}(a_n/2, b_n/2) \\ a_n &= a + n \\ b_n &= b + \sum_i (y_i - \mu)^2\end{aligned}$$

$$\mathbb{E}[\phi \mid \mu, y_1, \dots, y_n] = \frac{(a + n)/2}{(b + \sum_i (y_i - \mu)^2)/2}$$

What happens with a non-informative prior i.e

$a = b = \epsilon$ as $\epsilon \rightarrow 0$?



Normal Linear Regression Example

Model

$$\begin{aligned} Y_i \mid \beta, \phi &\stackrel{iid}{\sim} \mathbf{N}(x_i^T \beta, 1/\phi) \\ Y \mid \beta, \phi &\sim \mathbf{N}(X\beta, \phi^{-1}I_n) \\ \beta &\sim \mathbf{N}(b_0, \Phi_0^{-1}) \\ \phi &\sim \mathbf{N}(v_0/2, s_0/2) \end{aligned}$$



Normal Linear Regression Example

Model

$$\begin{aligned} Y_i | \beta, \phi &\stackrel{iid}{\sim} \mathbf{N}(x_i^T \beta, 1/\phi) \\ Y | \beta, \phi &\sim \mathbf{N}(X\beta, \phi^{-1}I_n) \\ \beta &\sim \mathbf{N}(b_0, \Phi_0^{-1}) \\ \phi &\sim \mathbf{N}(v_0/2, s_0/2) \end{aligned}$$

x_i is a $p \times 1$ vector of predictors and X is $n \times p$ matrix



Normal Linear Regression Example

Model

$$Y_i | \beta, \phi \stackrel{iid}{\sim} \mathbf{N}(x_i^T \beta, 1/\phi)$$

$$Y | \beta, \phi \sim \mathbf{N}(X\beta, \phi^{-1}I_n)$$

$$\beta \sim \mathbf{N}(b_0, \Phi_0^{-1})$$

$$\phi \sim \mathbf{N}(v_0/2, s_0/2)$$

x_i is a $p \times 1$ vector of predictors and X is $n \times p$ matrix

β is a $p \times 1$ vector of coefficients



Normal Linear Regression Example

Model

$$Y_i | \beta, \phi \stackrel{iid}{\sim} \mathbf{N}(x_i^T \beta, 1/\phi)$$

$$Y | \beta, \phi \sim \mathbf{N}(X\beta, \phi^{-1}I_n)$$

$$\beta \sim \mathbf{N}(b_0, \Phi_0^{-1})$$

$$\phi \sim \mathbf{N}(v_0/2, s_0/2)$$

x_i is a $p \times 1$ vector of predictors and X is $n \times p$ matrix

β is a $p \times 1$ vector of coefficients

Φ_0 is a $p \times p$ prior precision matrix



Normal Linear Regression Example

Model

$$\begin{aligned} Y_i \mid \beta, \phi &\stackrel{iid}{\sim} \mathbf{N}(x_i^T \beta, 1/\phi) \\ Y \mid \beta, \phi &\sim \mathbf{N}(X\beta, \phi^{-1}I_n) \\ \beta &\sim \mathbf{N}(b_0, \Phi_0^{-1}) \\ \phi &\sim \mathbf{N}(v_0/2, s_0/2) \end{aligned}$$

x_i is a $p \times 1$ vector of predictors and X is $n \times p$ matrix

β is a $p \times 1$ vector of coefficients

Φ_0 is a $p \times p$ prior precision matrix

Multivariate Normal density for β

$$\pi(\beta \mid b_0, \Phi_0) = \frac{|\Phi_0|^{1/2}}{(2\pi)^{p/2}} \exp \left\{ -\frac{1}{2} (\beta - b_0)^T \Phi_0 (\beta - b_0) \right\}$$



Full Conditional for β

$$\beta \mid \phi, y_1, \dots, y_n \sim \mathbf{N}(b_n, \Phi_n^{-1})$$

$$b_n = (\Phi_0 + \phi X^T X)^{-1} (\Phi_0 b_0 + \phi X^T X \hat{\beta})$$

$$\Phi_n = \Phi_0 + \phi X^T X$$



Derivation continued



Full Conditional for ϕ

$$\phi \mid \beta, y_1, \dots, y_n \sim \text{Gamma}((v_0 + n)/2, (s_0 + \sum_i (y_i - x_i^T \beta)))$$



Choice of Prior Precision

Non-Informative $\Phi_0 \rightarrow 0$



Choice of Prior Precision

Non-Informative $\Phi_0 \rightarrow 0$

- Formal Posterior given ϕ

$$\beta \mid \phi, y_1, \dots, y_n \sim \mathbf{N}(\hat{\beta}, \phi^{-1}(X^T X)^{-1})$$



Choice of Prior Precision

Non-Informative $\Phi_0 \rightarrow 0$

- Formal Posterior given ϕ

$$\beta \mid \phi, y_1, \dots, y_n \sim N(\hat{\beta}, \phi^{-1}(X^T X)^{-1})$$

- needs $X^T X$ to be full rank for distribution to be unique



Invariance and Choice of Mean/Precision

- the model in vector form

$$Y \sim \mathbb{N}_n(X\beta, \phi^{-1}I_n)$$



Invariance and Choice of Mean/Precision

- the model in vector form

$$Y \sim \mathbb{N}_n(X\beta, \phi^{-1}I_n)$$

- What if we transform the X matrix by $\tilde{X} = XH$ where H is $p \times p$ and invertible



Invariance and Choice of Mean/Precision

- the model in vector form

$$Y \sim \mathbb{N}_n(X\beta, \phi^{-1}I_n)$$

- What if we transform the X matrix by $\tilde{X} = XH$ where H is $p \times p$ and invertible
- obtain the posterior for $\tilde{\beta}$ using Y and \tilde{X}

$$Y \sim \mathbb{N}_n(\tilde{X}\tilde{\beta}, \phi^{-1}I_n)$$

- since $\tilde{X}\tilde{\beta} = XH\tilde{\beta} = X\beta$ invariance suggests that the posterior for β and $H\tilde{\beta}$ should be the same



Invariance and Choice of Mean/Precision

- the model in vector form

$$Y \sim \mathbf{N}_n(X\beta, \phi^{-1}I_n)$$

- What if we transform the X matrix by $\tilde{X} = XH$ where H is $p \times p$ and invertible
- obtain the posterior for $\tilde{\beta}$ using Y and \tilde{X}

$$Y \sim \mathbf{N}_n(\tilde{X}\tilde{\beta}, \phi^{-1}I_n)$$

- since $\tilde{X}\tilde{\beta} = XH\tilde{\beta} = X\beta$ invariance suggests that the posterior for β and $H\tilde{\beta}$ should be the same
- or the posterior of $H^{-1}\beta$ and $\tilde{\beta}$ should be the same



Invariance and Choice of Mean/Precision

- the model in vector form

$$Y \sim \mathbf{N}_n(X\beta, \phi^{-1}I_n)$$

- What if we transform the X matrix by $\tilde{X} = XH$ where H is $p \times p$ and invertible
- obtain the posterior for $\tilde{\beta}$ using Y and \tilde{X}

$$Y \sim \mathbf{N}_n(\tilde{X}\tilde{\beta}, \phi^{-1}I_n)$$

- since $\tilde{X}\tilde{\beta} = XH\tilde{\beta} = X\beta$ invariance suggests that the posterior for β and $H\tilde{\beta}$ should be the same
- or the posterior of $H^{-1}\beta$ and $\tilde{\beta}$ should be the same
- with some linear algebra we can show that this is true if $b_0 = 0$ and Φ_0 is $kX^T X$ for some k (show!)



Zellner's g-prior

Popular choice is to take $k = \phi/g$ which is a special case of Zellner's g-prior

$$\beta | \phi, g \sim N \left(0, \frac{g}{\phi} (X^T X)^{-1} \right)$$



Zellner's g-prior

Popular choice is to take $k = \phi/g$ which is a special case of Zellner's g-prior

$$\beta | \phi, g \sim N \left(0, \frac{g}{\phi} (X^T X)^{-1} \right)$$

- Full conditional

$$\beta | \phi, g \sim N \left(\frac{g}{1+g} \hat{\beta}, \frac{1}{\phi} \frac{g}{1+g} (X^T X)^{-1} \right)$$



Zellner's g-prior

Popular choice is to take $k = \phi/g$ which is a special case of Zellner's g-prior

$$\beta | \phi, g \sim N \left(0, \frac{g}{\phi} (X^T X)^{-1} \right)$$

- Full conditional

$$\beta | \phi, g \sim N \left(\frac{g}{1+g} \hat{\beta}, \frac{1}{\phi} \frac{g}{1+g} (X^T X)^{-1} \right)$$

- one parameter g controls shrinkage



Zellner's g-prior

Popular choice is to take $k = \phi/g$ which is a special case of Zellner's g-prior

$$\beta | \phi, g \sim N \left(0, \frac{g}{\phi} (X^T X)^{-1} \right)$$

- Full conditional

$$\beta | \phi, g \sim N \left(\frac{g}{1+g} \hat{\beta}, \frac{1}{\phi} \frac{g}{1+g} (X^T X)^{-1} \right)$$

- one parameter g controls shrinkage

if $\phi \sim \text{Gamma}(v_0/2, s_0/2)$ then posterior is

$$\phi | y_1, \dots, y_n \sim \text{Gamma}(v_n/2, s_n/2)$$



Zellner's g-prior

Popular choice is to take $k = \phi/g$ which is a special case of Zellner's g-prior

$$\beta | \phi, g \sim N \left(0, \frac{g}{\phi} (X^T X)^{-1} \right)$$

- Full conditional

$$\beta | \phi, g \sim N \left(\frac{g}{1+g} \hat{\beta}, \frac{1}{\phi} \frac{g}{1+g} (X^T X)^{-1} \right)$$

- one parameter g controls shrinkage

if $\phi \sim \text{Gamma}(v_0/2, s_0/2)$ then posterior is

$$\phi | y_1, \dots, y_n \sim \text{Gamma}(v_n/2, s_n/2)$$

Conjugate so we could skip Gibbs sampling and sample directly from gamma and then conditional normal!



Ridge Regression

If $X^T X$ is nearly singular, certain elements of β or (linear combinations of β) may have huge variances under the g -prior (or flat prior) as the MLEs are highly unstable!



Ridge Regression

If $X^T X$ is nearly singular, certain elements of β or (linear combinations of β) may have huge variances under the g -prior (or flat prior) as the MLEs are highly unstable!

Ridge regression protects against the explosion of variances and ill-conditioning with the conjugate priors:

$$\beta | \phi \sim N(0, \frac{1}{\phi \lambda} I_p)$$



Ridge Regression

If $X^T X$ is nearly singular, certain elements of β or (linear combinations of β) may have huge variances under the g -prior (or flat prior) as the MLEs are highly unstable!

Ridge regression protects against the explosion of variances and ill-conditioning with the conjugate priors:

$$\beta | \phi \sim N(0, \frac{1}{\phi\lambda} I_p)$$

Posterior for β (conjugate case)

$$\beta | \phi, \lambda, y_1, \dots, y_n \sim N \left((\lambda I_p + X^T X)^{-1} X^T Y, \frac{1}{\phi} (\lambda I_p + X^T X)^{-1} \right)$$



Bayes Regression

- Posterior mean (or mode) given λ is biased, but can show that there **always** is a value of λ where the frequentist's expected squared error loss is smaller for the Ridge estimator than MLE!



Bayes Regression

- Posterior mean (or mode) given λ is biased, but can show that there **always** is a value of λ where the frequentist's expected squared error loss is smaller for the Ridge estimator than MLE!
- related to penalized maximum likelihood estimation



Bayes Regression

- Posterior mean (or mode) given λ is biased, but can show that there **always** is a value of λ where the frequentist's expected squared error loss is smaller for the Ridge estimator than MLE!
 - related to penalized maximum likelihood estimation
 - Choice of λ



Bayes Regression

- Posterior mean (or mode) given λ is biased, but can show that there **always** is a value of λ where the frequentist's expected squared error loss is smaller for the Ridge estimator than MLE!
 - related to penalized maximum likelihood estimation
 - Choice of λ
 - Bayes Regression and choice of Φ_0 in general is a very important problem and provides the foundation for many variations on shrinkage estimators, variable selection, hierarchical models, nonparameteric regression and more!



Bayes Regression

- Posterior mean (or mode) given λ is biased, but can show that there **always** is a value of λ where the frequentist's expected squared error loss is smaller for the Ridge estimator than MLE!
 - related to penalized maximum likelihood estimation
 - Choice of λ
 - Bayes Regression and choice of Φ_0 in general is a very important problem and provides the foundation for many variations on shrinkage estimators, variable selection, hierarchical models, nonparameteric regression and more!
 - Be sure that you can derive the full conditional posteriors for β and ϕ as well as the joint posterior in the conjugate case!



Comments

- Why don't we treat each individual β_j as a separate block?



Comments

- Why don't we treat each individual β_j as a separate block?
- Gibbs always accepts, but can mix slowly if parameters in different blocks are highly correlated!



Comments

- Why don't we treat each individual β_j as a separate block?
- Gibbs always accepts, but can mix slowly if parameters in different blocks are highly correlated!
- Use block sizes in Gibbs that are as big as possible to improve mixing (proven faster convergence)



Comments

- Why don't we treat each individual β_j as a separate block?
- Gibbs always accepts, but can mix slowly if parameters in different blocks are highly correlated!
- Use block sizes in Gibbs that are as big as possible to improve mixing (proven faster convergence)
- Collapse the sampler by integrating out as many parameters as possible (as long as resulting sampler has good mixing)



Comments

- Why don't we treat each individual β_j as a separate block?
- Gibbs always accepts, but can mix slowly if parameters in different blocks are highly correlated!
- Use block sizes in Gibbs that are as big as possible to improve mixing (proven faster convergence)
- Collapse the sampler by integrating out as many parameters as possible (as long as resulting sampler has good mixing)
- can use Gibbs steps and (adaptive) Metropolis Hastings steps together



Comments

- Why don't we treat each individual β_j as a separate block?
- Gibbs always accepts, but can mix slowly if parameters in different blocks are highly correlated!
- Use block sizes in Gibbs that are as big as possible to improve mixing (proven faster convergence)
- Collapse the sampler by integrating out as many parameters as possible (as long as resulting sampler has good mixing)
- can use Gibbs steps and (adaptive) Metropolis Hastings steps together
- Introduce latent variables (data augmentation) to allow Gibbs steps (Next class)

