

# Bayesian Model Choice

Hoff Chapter 9, Liang et al 2007, Hoeting et al (1999), Clyde & George (2004) Statistical Science

October 31, 2022

# Posterior Distribution

With a Normal-Gamma Prior  $\text{NG}(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$ , the posterior is  $\text{NG}(\mathbf{b}_n, \Phi_n, \nu_n, SS_n)$ : with

$$\boldsymbol{\beta} \mid \boldsymbol{\phi}, \mathbf{Y} \sim \text{N}(\mathbf{b}_n, (\boldsymbol{\phi} \Phi_n)^{-1})$$

# Posterior Distribution

With a Normal-Gamma Prior  $\text{NG}(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$ , the posterior is  $\text{NG}(\mathbf{b}_n, \Phi_n, \nu_n, SS_n)$ : with

$$\begin{aligned}\beta \mid \phi, Y &\sim \text{N}(\mathbf{b}_n, (\phi \Phi_n)^{-1}) \\ \phi \mid Y &\sim \text{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

and hyper-parameters

$$\Phi_n = X^T X + \Phi_0$$

# Posterior Distribution

With a Normal-Gamma Prior  $\text{NG}(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$ , the posterior is  $\text{NG}(\mathbf{b}_n, \Phi_n, \nu_n, SS_n)$ : with

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \text{N}(\mathbf{b}_n, (\phi \Phi_n)^{-1}) \\ \phi \mid \mathbf{Y} &\sim \text{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

and hyper-parameters

$$\begin{aligned}\Phi_n &= \mathbf{X}^T \mathbf{X} + \Phi_0 \\ \mathbf{b}_n &= \Phi_n^{-1} (\mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \Phi_0 \mathbf{b}_0)\end{aligned}$$

# Posterior Distribution

With a Normal-Gamma Prior  $\text{NG}(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$ , the posterior is  $\text{NG}(\mathbf{b}_n, \Phi_n, \nu_n, SS_n)$ : with

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \text{N}(\mathbf{b}_n, (\phi \Phi_n)^{-1}) \\ \phi \mid \mathbf{Y} &\sim \text{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

and hyper-parameters

$$\begin{aligned}\Phi_n &= \mathbf{X}^T \mathbf{X} + \Phi_0 \\ \mathbf{b}_n &= \Phi_n^{-1} (\mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \Phi_0 \mathbf{b}_0) \\ \nu_n &= n + \nu_0\end{aligned}$$

# Posterior Distribution

With a Normal-Gamma Prior  $NG(b_0, \Phi_0, \nu_0, SS_0)$ , the posterior is  $NG(b_n, \Phi_n, \nu_n, SS_n)$ : with

$$\begin{aligned}\beta \mid \phi, Y &\sim N(b_n, (\phi \Phi_n)^{-1}) \\ \phi \mid Y &\sim G\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

and hyper-parameters

$$\begin{aligned}\Phi_n &= X^T X + \Phi_0 \\ b_n &= \Phi_n^{-1}(X^T X \hat{\beta} + \Phi_0 b_0) \\ \nu_n &= n + \nu_0 \\ SS_n &= SSE + SS_0 + \hat{\beta}^T X^T X \hat{\beta} + b_0^T \Phi_0 b_0 - b_n^T \Phi_n b_n\end{aligned}$$

# Marginal Distribution from Normal–Gamma

## Theorem

Let  $\boldsymbol{\theta} \mid \phi \sim N(m, \frac{1}{\phi}\Sigma)$  and  $\phi \sim G(\nu/2, \nu\hat{\sigma}^2/2)$ . Then  $\boldsymbol{\theta}$  ( $p \times 1$ ) has a  $p$  dimensional multivariate  $t$  distribution

$$\boldsymbol{\theta} \sim t_{\nu}(m, \hat{\sigma}^2\Sigma)$$

with density

$$p(\boldsymbol{\theta}) \propto \left[ 1 + \frac{1}{\nu} \frac{(\boldsymbol{\theta} - m)^T \Sigma^{-1} (\boldsymbol{\theta} - m)}{\hat{\sigma}^2} \right]^{-\frac{p+\nu}{2}}$$

# Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$



# Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$p(\boldsymbol{\theta}) \propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi$$

## Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \end{aligned}$$

# Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m})} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m}) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m}) + \nu \hat{\sigma}^2}{2}} d\phi \end{aligned}$$

# Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left( \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

# Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left( \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \\ &\propto \left( (\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2 \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

## Derivation

Marginal density  $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left( \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \\ &\propto \left( (\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2 \right)^{-\frac{p+\nu}{2}} \\ &\propto \left( 1 + \frac{1}{\nu} \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)}{\hat{\sigma}^2} \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

## Marginal Posterior Distribution of $\beta$

$$\beta \mid \phi, Y \sim N(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

## Marginal Posterior Distribution of $\beta$

$$\beta \mid \phi, Y \sim N(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

$$\phi \mid Y \sim G\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)$$



## Marginal Posterior Distribution of $\beta$

$$\beta \mid \phi, Y \sim N(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

$$\phi \mid Y \sim G\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)$$

Let  $\hat{\sigma}^2 = SS_n/\nu_n$  (Bayesian MSE)

## Marginal Posterior Distribution of $\beta$

$$\begin{aligned}\beta \mid \phi, Y &\sim N(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1}) \\ \phi \mid Y &\sim G\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

Let  $\hat{\sigma}^2 = SS_n/\nu_n$  (Bayesian MSE)

Then the marginal posterior distribution of  $\beta$  is

$$\beta \mid Y \sim t_{\nu_n}(\mathbf{b}_n, \hat{\sigma}^2 \Phi_n^{-1})$$

## Marginal Posterior Distribution of $\beta$

$$\begin{aligned}\beta \mid \phi, Y &\sim N(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1}) \\ \phi \mid Y &\sim G\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

Let  $\hat{\sigma}^2 = SS_n/\nu_n$  (Bayesian MSE)

Then the marginal posterior distribution of  $\beta$  is

$$\beta \mid Y \sim t_{\nu_n}(\mathbf{b}_n, \hat{\sigma}^2 \Phi_n^{-1})$$

Any linear combination  $\mathbf{x}^T \beta$

$$\mathbf{x}^T \beta \mid Y \sim t_{\nu_n}(\mathbf{x}^T \mathbf{b}_n, \hat{\sigma}^2 \mathbf{x}^T \Phi_n^{-1} \mathbf{x})$$

has a univariate  $t$  distribution with  $\nu_n$  degrees of freedom

# Predictive Distribution

Suppose  $Y^* \mid \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

# Predictive Distribution

Suppose  $Y^* \mid \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* \mid Y$ ?

# Predictive Distribution

Suppose  $Y^* \mid \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* \mid Y$ ?

$Y^* = X^*\beta + \epsilon^*$  and  $\epsilon^*$  is independent of  $Y$  given  $\phi$

# Predictive Distribution

Suppose  $Y^* \mid \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* \mid Y$ ?

$Y^* = X^*\beta + \epsilon^*$  and  $\epsilon^*$  is independent of  $Y$  given  $\phi$

$$X^*\beta + \epsilon^* \mid \phi, Y \sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi)$$

# Predictive Distribution

Suppose  $Y^* \mid \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* \mid Y$ ?

$Y^* = X^*\beta + \epsilon^*$  and  $\epsilon^*$  is independent of  $Y$  given  $\phi$

$$\begin{aligned} X^*\beta + \epsilon^* \mid \phi, Y &\sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi) \\ Y^* \mid \phi, Y &\sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi) \end{aligned}$$



# Predictive Distribution

Suppose  $Y^* | \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* | Y$ ?

$Y^* = X^*\beta + \epsilon^*$  and  $\epsilon^*$  is independent of  $Y$  given  $\phi$

$$X^*\beta + \epsilon^* | \phi, Y \sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi)$$

$$Y^* | \phi, Y \sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi)$$

$$\phi | Y \sim G\left(\frac{\nu_n}{2}, \frac{\hat{\sigma}^2 \nu_n}{2}\right)$$

# Predictive Distribution

Suppose  $Y^* | \beta, \phi \sim N(X^*\beta, I/\phi)$  and is conditionally independent of  $Y$  given  $\beta$  and  $\phi$

What is the predictive distribution of  $Y^* | Y$ ?

$Y^* = X^*\beta + \epsilon^*$  and  $\epsilon^*$  is independent of  $Y$  given  $\phi$

$$X^*\beta + \epsilon^* | \phi, Y \sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi)$$

$$Y^* | \phi, Y \sim N(X^*b_n, (X^*\Phi_n^{-1}X^{*T} + I)/\phi)$$

$$\phi | Y \sim G\left(\frac{\nu_n}{2}, \frac{\hat{\sigma}^2 \nu_n}{2}\right)$$

$$Y^* | Y \sim t_{\nu_n}(X^*b_n, \hat{\sigma}^2(I + X^*\Phi_n^{-1}X^T))$$

# Alternative Derivation

Conditional Distribution:

$$f(Y^* | Y) = \frac{f(Y^*, Y)}{f(Y)}$$

# Alternative Derivation

Conditional Distribution:

$$\begin{aligned} f(Y^* | Y) &= \frac{f(Y^*, Y)}{f(Y)} \\ &= \frac{\iint f(Y^*, Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \end{aligned}$$

# Alternative Derivation

Conditional Distribution:

$$\begin{aligned} f(Y^* | Y) &= \frac{f(Y^*, Y)}{f(Y)} \\ &= \frac{\iint f(Y^*, Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \\ &= \frac{\iint f(Y^* | \beta, \phi) f(Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \end{aligned}$$

# Alternative Derivation

Conditional Distribution:

$$\begin{aligned} f(Y^* | Y) &= \frac{f(Y^*, Y)}{f(Y)} \\ &= \frac{\iint f(Y^*, Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \\ &= \frac{\iint f(Y^* | \beta, \phi) f(Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \\ &= \iint f(Y^* | \beta, \phi) p(\beta, \phi | Y) d\beta d\phi \end{aligned}$$

# Alternative Derivation

Conditional Distribution:

$$\begin{aligned} f(Y^* | Y) &= \frac{f(Y^*, Y)}{f(Y)} \\ &= \frac{\iint f(Y^*, Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \\ &= \frac{\iint f(Y^* | \beta, \phi) f(Y | \beta, \phi) p(\beta, \phi) d\beta d\phi}{f(Y)} \\ &= \iint f(Y^* | \beta, \phi) p(\beta, \phi | Y) d\beta d\phi \end{aligned}$$

Requires completing the square/quadratic!

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\boldsymbol{\theta}$  is conjugate for a sampling model  $p(y \mid \boldsymbol{\theta})$  if for every  $p(\boldsymbol{\theta}) \in \mathcal{P}$ ,  $p(\boldsymbol{\theta} \mid Y) \in \mathcal{P}$ .



# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data
- ▶ limiting “non-proper” form recovers MLEs

# Conjugate Priors

## Definition

A class of prior distributions  $\mathcal{P}$  for  $\theta$  is conjugate for a sampling model  $p(y \mid \theta)$  if for every  $p(\theta) \in \mathcal{P}$ ,  $p(\theta \mid Y) \in \mathcal{P}$ .

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data
- ▶ limiting “non-proper” form recovers MLEs

Choice of conjugate prior?

# Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization



# Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{J}(\boldsymbol{\theta})|^{1/2}$$

# Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{I}(\boldsymbol{\theta})|^{1/2}$$

where  $\mathcal{I}(\boldsymbol{\theta})$  is the Expected Fisher Information matrix

# Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{I}(\boldsymbol{\theta})|^{1/2}$$

where  $\mathcal{I}(\boldsymbol{\theta})$  is the Expected Fisher Information matrix

$$\mathcal{I}(\boldsymbol{\theta}) = -\mathbb{E}\left[\frac{\partial^2 \log(\mathcal{L}(\boldsymbol{\theta}))}{\partial \theta_i \partial \theta_j}\right]$$

# Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(I - P_X)Y\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

# Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(I - P_X)Y\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} = \begin{bmatrix} -\phi(X^T X) & -(X^T X)(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

# Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(I - P_X)Y\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\begin{aligned} \frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} &= \begin{bmatrix} -\phi(X^T X) & -(X^T X)(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix} \\ E\left[\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T}\right] &= \begin{bmatrix} -\phi(X^T X) & 0_p \\ 0_p^T & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix} \end{aligned}$$

# Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(I - P_X)Y\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} = \begin{bmatrix} -\phi(X^T X) & -(X^T X)(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (X^T X) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$E\left[\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T}\right] = \begin{bmatrix} -\phi(X^T X) & 0_p \\ 0_p^T & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$\mathcal{I}((\boldsymbol{\beta}, \phi)^T) = \begin{bmatrix} \phi(X^T X) & 0_p \\ 0_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

# Jeffreys Prior

Jeffreys Prior

$$p_J(\boldsymbol{\beta}, \phi) \propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2}$$



# Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left( \frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \end{aligned}$$

# Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left( \frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\ &\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \end{aligned}$$

# Jeffreys Prior

## Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left( \frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\ &\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \\ &\propto \phi^{p/2-1} \end{aligned}$$

# Jeffreys Prior

## Jeffreys Prior

$$\begin{aligned}p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\&= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left( \frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\&\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \\&\propto \phi^{p/2-1}\end{aligned}$$

Improper prior  $\iint p_J(\boldsymbol{\beta}, \phi) d\boldsymbol{\beta} d\phi$  not finite

# Formal Bayes Posterior

$$p(\boldsymbol{\beta}, \phi \mid Y) \propto p(Y \mid \boldsymbol{\beta}, \phi) \phi^{p/2-1}$$

# Formal Bayes Posterior

$$p(\beta, \phi \mid Y) \propto p(Y \mid \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

## Formal Bayes Posterior

$$p(\beta, \phi | Y) \propto p(Y | \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\beta | \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi | Y &\sim G(n/2, \|Y - X\hat{\beta}\|^2/2)\end{aligned}$$

## Formal Bayes Posterior

$$p(\beta, \phi | Y) \propto p(Y | \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\beta | \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi | Y &\sim G(n/2, \|Y - X\hat{\beta}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with  $b_0 = 0$ ,  $\Phi = 0$ ,  $\nu_0 = 0$  and  $SS_0 = 0$



## Formal Bayes Posterior

$$p(\beta, \phi | Y) \propto p(Y | \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\beta | \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi | Y &\sim G(n/2, \|Y - X\hat{\beta}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with  $b_0 = 0$ ,  $\Phi = 0$ ,  $\nu_0 = 0$  and  $SS_0 = 0$

## Formal Bayes Posterior

$$p(\beta, \phi | Y) \propto p(Y | \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\beta | \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi | Y &\sim G(n/2, \|Y - X\hat{\beta}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with  $b_0 = 0$ ,  $\Phi = 0$ ,  $\nu_0 = 0$  and  $SS_0 = 0$

Jeffreys did not recommend using this Posterior does not depend on dimension  $p$

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶  $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶  $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(X^T X) & 0_p \\ 0_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶  $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(X^T X) & 0_p \\ 0_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi X^T X|^{1/2} \propto 1$$

# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶  $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(X^T X) & 0_p \\ 0_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi X^T X|^{1/2} \propto 1$$

$$p_{IJ}(\phi) \propto \phi^{-1}$$



# Independent Jeffreys Prior

- ▶ Treat  $\beta$  and  $\phi$  separately (“orthogonal parameterization”)
- ▶  $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶  $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(X^T X) & 0_p \\ 0_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi X^T X|^{1/2} \propto 1$$

$$p_{IJ}(\phi) \propto \phi^{-1}$$

Independent Jeffreys Prior is

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta) p_{IJ}(\phi) = \phi^{-1}$$

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\beta \mid \phi, Y \sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1})$$

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi \mid Y &\sim G((n - p)/2, \|Y - X\hat{\beta}\|^2/2)\end{aligned}$$

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi \mid Y &\sim G((n - p)/2, \|Y - X\hat{\beta}\|^2/2) \\ \beta \mid Y &\sim t_{n-p}(\hat{\beta}, \hat{\sigma}^2 (X^T X)^{-1})\end{aligned}$$

# Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, Y &\sim N(\hat{\beta}, (X^T X)^{-1} \phi^{-1}) \\ \phi \mid Y &\sim G((n-p)/2, \|Y - X\hat{\beta}\|^2/2) \\ \beta \mid Y &\sim t_{n-p}(\hat{\beta}, \hat{\sigma}^2 (X^T X)^{-1})\end{aligned}$$

Bayesian Credible Sets  $p(\beta \in C_\alpha) = 1 - \alpha$  correspond to frequentist Confidence Regions

$$\frac{\lambda^T \beta - \lambda^T \hat{\beta}}{\sqrt{\hat{\sigma}^2 \lambda^T (X^T X)^{-1} \lambda}} \sim t_{n-p}$$

## Zellner's $g$ -prior

$$\text{Zellner's } g\text{-prior(s) } \beta \mid \phi \sim N(b_0, g(X^T X)^{-1} / \phi)$$



## Zellner's $g$ -prior

Zellner's  $g$ -prior(s)  $\beta \mid \phi \sim N(b_0, g(X^T X)^{-1} / \phi)$

$$\beta \mid Y, \phi \sim N \left( \frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} b_0, \frac{g}{1+g} (X^T X)^{-1} \phi^{-1} \right)$$

## Zellner's $g$ -prior

Zellner's  $g$ -prior(s)  $\beta \mid \phi \sim N(b_0, g(X^T X)^{-1}/\phi)$

$$\beta \mid Y, \phi \sim N\left(\frac{g}{1+g}\hat{\beta} + \frac{1}{1+g}b_0, \frac{g}{1+g}(X^T X)^{-1}\phi^{-1}\right)$$

- Invariance: Require posterior of  $X\beta$  equal the posterior of  $XH\alpha$

## Zellner's $g$ -prior

Zellner's  $g$ -prior(s)  $\beta \mid \phi \sim N(b_0, g(X^T X)^{-1}/\phi)$

$$\beta \mid Y, \phi \sim N\left(\frac{g}{1+g}\hat{\beta} + \frac{1}{1+g}b_0, \frac{g}{1+g}(X^T X)^{-1}\phi^{-1}\right)$$

- ▶ Invariance: Require posterior of  $X\beta$  equal the posterior of  $XH\alpha$  ( $a_0 = H^{-1}b_0$ ) ( $b_0 = 0$ )
- ▶ Choice of  $g$ ?

## Zellner's $g$ -prior

Zellner's  $g$ -prior(s)  $\beta \mid \phi \sim N(b_0, g(X^T X)^{-1}/\phi)$

$$\beta \mid Y, \phi \sim N \left( \frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} b_0, \frac{g}{1+g} (X^T X)^{-1} \phi^{-1} \right)$$

- ▶ Invariance: Require posterior of  $X\beta$  equal the posterior of  $XH\alpha$  ( $a_0 = H^{-1}b_0$ ) ( $b_0 = 0$ )
- ▶ Choice of  $g$ ?
- ▶  $\frac{g}{1+g}$  weight given to the data

## Zellner's $g$ -prior


Zellner's  $g$ -prior(s)  $\beta \mid \phi \sim N(b_0, g(X^T X)^{-1}/\phi)$

$$\beta \mid Y, \phi \sim N \left( \frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} b_0, \frac{g}{1+g} (X^T X)^{-1} \phi^{-1} \right)$$

- ▶ Invariance: Require posterior of  $X\beta$  equal the posterior of  $XH\alpha$  ( $a_0 = H^{-1}b_0$ ) ( $b_0 = 0$ )
- ▶ Choice of  $g$ ?
- ▶  $\frac{g}{1+g}$  weight given to the data
- ▶ Fixed  $g$  effect does not vanish as  $n \rightarrow \infty$
- ▶ Use  $g = n$  or place a prior distribution on  $g$

# Shrinkage

Posterior mean under  $g$ -prior with  $b_0 = 0$   $\frac{g}{1+g}\hat{\beta}$



# Ridge Regression

- ▶ If  $X^T X$  is nearly singular, certain elements of  $\beta$  or (linear combinations of  $\beta$ ) may have huge variances under the  $g$ -prior (or flat prior) as the MLEs are highly unstable!
- ▶ **Ridge regression** protects against the explosion of variances and ill-conditioning with the conjugate prior:

$$\beta \mid \phi \sim N(0, \frac{1}{\phi\lambda} I_p)$$

- ▶ Posterior for  $\beta$  (conjugate case)

$$\beta \mid \phi, \lambda, Y \sim N \left( (\lambda I_p + X^T X)^{-1} X^T Y, \frac{1}{\phi} (\lambda I_p + X^T X)^{-1} \right)$$

- ▶ induces shrinkage as well!

# Model Choice ?

- ▶ Redundant variables lead to unstable estimates



# Model Choice ?

- ▶ Redundant variables lead to unstable estimates
- ▶ Some variables may not be relevant ( $\beta_j = 0$ )

# Model Choice ?

- ▶ Redundant variables lead to unstable estimates
- ▶ Some variables may not be relevant ( $\beta_j = 0$ )
- ▶ Can we infer a "good" model from the data?
- ▶ Expand model hierarchically to introduce another latent variable  $\gamma$  that encodes models  $\mathcal{M}_\gamma$   $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_p)^T$  where

$$\gamma_j = 0 \Leftrightarrow \beta_j = 0$$

$$\gamma_j = 1 \Leftrightarrow \beta_j \neq 0$$

- ▶ Find Bayes factors and posterior probabilities of models  $\mathcal{M}_\gamma$
- ▶  $2^p$  models!

# Zellner's g-prior

Centered model:

$$Y = \mathbf{1}_n \alpha + X^c \beta + \epsilon$$

where  $X^c$  is the centered design matrix where all variables have had their mean subtracted

# Zellner's g-prior

Centered model:

$$Y = 1_n\alpha + X^c\beta + \epsilon$$

where  $X^c$  is the centered design matrix where all variables have had their mean subtracted

►  $p(\alpha, \phi) \propto 1/\phi$

# Zellner's g-prior

Centered model:

$$Y = 1_n\alpha + X^c\beta + \epsilon$$

where  $X^c$  is the centered design matrix where all variables have had their mean subtracted

- ▶  $p(\alpha, \phi) \propto 1/\phi$
- ▶  $\beta_\gamma \mid \alpha, \phi, \gamma \sim N(0, g\phi^{-1}(X_\gamma^c'X_\gamma^c)^{-1})$

## Zellner's g-prior

Centered model:

$$Y = 1_n \alpha + X^c \beta + \epsilon$$

where  $X^c$  is the centered design matrix where all variables have had their mean subtracted

- ▶  $p(\alpha, \phi) \propto 1/\phi$
- ▶  $\beta_\gamma \mid \alpha, \phi, \gamma \sim N(0, g\phi^{-1}(X_\gamma^c' X_\gamma^c)^{-1})$

which leads to marginal likelihood of  $\gamma$  that is proportional to

$$p(Y \mid \gamma) = C(1 + g)^{\frac{n-p-1}{2}} (1 + g(1 - R_\gamma^2))^{-\frac{(n-1)}{2}}$$

where  $R^2$  is the usual coefficient of determination for model  $\mathcal{M}_\gamma$ .

# Zellner's g-prior

Centered model:

$$Y = 1_n \alpha + X^c \beta + \epsilon$$

where  $X^c$  is the centered design matrix where all variables have had their mean subtracted

- ▶  $p(\alpha, \phi) \propto 1/\phi$
- ▶  $\beta_\gamma \mid \alpha, \phi, \gamma \sim N(0, g\phi^{-1}(X_\gamma^c' X_\gamma^c)^{-1})$

which leads to marginal likelihood of  $\gamma$  that is proportional to

$$p(Y \mid \gamma) = C(1 + g)^{\frac{n-p-1}{2}} (1 + g(1 - R_\gamma^2))^{-\frac{(n-1)}{2}}$$

where  $R^2$  is the usual coefficient of determination for model  $\mathcal{M}_\gamma$ .  
Trade-off of model complexity versus goodness of fit

Lastly, assign distribution to space of models

# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals



# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals
- ▶ Find inverse of  $I_n + gP_{X_\gamma}$  (properties of projections)

# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals
- ▶ Find inverse of  $I_n + gP_{X_\gamma}$  (properties of projections)
- ▶ Find determinant of  $\phi(I_n + gP_{X_\gamma})$

# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals
- ▶ Find inverse of  $I_n + gP_{X_\gamma}$  (properties of projections)
- ▶ Find determinant of  $\phi(I_n + gP_{X_\gamma})$
- ▶ Integrate out intercept (normal)

# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals
- ▶ Find inverse of  $I_n + gP_{X_\gamma}$  (properties of projections)
- ▶ Find determinant of  $\phi(I_n + gP_{X_\gamma})$
- ▶ Integrate out intercept (normal)
- ▶ Integrate out  $\phi$  (gamma)

# Sketch

- ▶ Integrate out  $\beta_\gamma$  using sums of normals
- ▶ Find inverse of  $I_n + gP_{X_\gamma}$  (properties of projections)
- ▶ Find determinant of  $\phi(I_n + gP_{X_\gamma})$
- ▶ Integrate out intercept (normal)
- ▶ Integrate out  $\phi$  (gamma)
- ▶ algebra to simplify quadratic forms to  $R_\gamma^2$

Or integrate  $\alpha$ ,  $\beta_\gamma$  and  $\phi$  (complete the square!)

# Priors on Model Space

$$p(\mathcal{M}_\gamma) \Leftrightarrow p(\gamma)$$

- ▶  $p(\gamma_j = 1) = .5 \Rightarrow P(\mathcal{M}_\gamma) = .5^p$  Uniform on space of models

# Priors on Model Space

$$p(\mathcal{M}_\gamma) \Leftrightarrow p(\gamma)$$

- ▶  $p(\gamma_j = 1) = .5 \Rightarrow P(\mathcal{M}_\gamma) = .5^p$  Uniform on space of models  
 $p_\gamma \sim \text{Bin}(p, .5)$
- ▶  $\gamma_j \mid \pi \stackrel{\text{iid}}{\sim} \text{Ber}(\pi)$  and  $\pi \sim \text{Beta}(a, b)$  then  $p_\gamma \sim \text{BB}_p(a, b)$

$$p(p_\gamma \mid p, a, b) = \frac{\Gamma(p+1)\Gamma(p_\gamma+a)\Gamma(p-p_\gamma+b)\Gamma(a+b)}{\Gamma(p_\gamma+1)\Gamma(p-p_\gamma+1)\Gamma(p+a+b)\Gamma(a)\Gamma(b)}$$

- ▶  $p_\gamma \sim \text{BB}_p(1, 1) \sim \text{Unif}(0, p)$

# Posterior Probabilities of Models

- Calculate analytically under enumeration

$$p(\mathcal{M}_\gamma | Y) = \frac{p(Y | \gamma)p(\gamma)}{\sum_{\gamma' \in \Gamma} p(Y | \gamma')p(\gamma')}$$

Express as a function of Bayes factors and prior odds!

- Use MCMC over  $\Gamma$  - Gibbs, Metropolis Hastings
- Do we need to run MCMC over  $\gamma$ ,  $\beta_\gamma$ ,  $\alpha$ , and  $\phi$ ?

Inference/Decisions ?