

STA 601: Bayesian Model Averaging

STA 601 Fall 2021

Merlise Clyde

October 19, 2021



Posteriors

Likelihood under model γ

$$p(\mathbf{y} \mid \mathbf{X}_\gamma, \gamma, \alpha, \beta_\gamma, \phi) \propto (\phi^{\frac{n}{2}} \exp \left\{ -\frac{\phi}{2} (\mathbf{y} - \mathbf{1}\alpha - \mathbf{X}_\gamma \beta_\gamma)^T (\mathbf{y} - \mathbf{1}\alpha - \mathbf{X}_\gamma \beta_\gamma) \right\})$$



Posteriors

Likelihood under model γ

$$p(\mathbf{y} \mid \mathbf{X}_\gamma, \gamma, \alpha, \boldsymbol{\beta}_\gamma, \phi) \propto (\phi^{\frac{n}{2}} \exp \left\{ -\frac{\phi}{2} (\mathbf{y} - \mathbf{1}\alpha - \mathbf{X}_\gamma \boldsymbol{\beta}_\gamma)^T (\mathbf{y} - \mathbf{1}\alpha - \mathbf{X}_\gamma \boldsymbol{\beta}_\gamma) \right\})$$

Independent Jeffrey's priors on common parameters and the g-prior

$$\begin{aligned} \pi(\alpha, \phi) &= \phi^{-1} \\ \pi(\boldsymbol{\beta}_\gamma | \phi) &= \mathbf{N}_p \left(\boldsymbol{\beta}_{0\gamma} = \mathbf{0}, \Sigma_{0\gamma} = \frac{g}{\phi} [\mathbf{X}_\gamma^T \mathbf{X}_\gamma]^{-1} \right) \end{aligned}$$



Posteriors

With those pieces, the conditional posteriors are straightforward

$$\alpha \mid \phi, y \sim \text{N} \left(\bar{y}, \frac{1}{n\phi} \right)$$

$$\beta_\gamma \mid \gamma, \phi, g, y \sim \text{N} \left(\frac{g}{1+g} \hat{\beta}_\gamma, \frac{g}{1+g} \frac{1}{\phi} [\mathbf{X}_\gamma^T \mathbf{X}_\gamma]^{-1} \right)$$

$$\phi \mid \gamma, y \sim \text{Gamma} \left(\frac{n-1}{2}, \frac{\text{TotalSS} - \frac{g}{1+g} \text{RegSS}}{2} \right)$$

$$p(\gamma \mid y) \propto p(y \mid \gamma) p(\gamma)$$

$$\text{TotalSS} \equiv \sum_i (y_i - \bar{y})^2 \quad \text{RegSS} \equiv \hat{\beta}_\gamma^T \mathbf{X}_\gamma^T \mathbf{X}_\gamma \hat{\beta}_\gamma$$

$$R_\gamma^2 = \frac{\text{RegSS}}{\text{TotalSS}} = 1 - \frac{\text{ErrorSS}}{\text{TotalSS}}$$



Posteriors

With those pieces, the conditional posteriors are straightforward

$$\alpha \mid \phi, y \sim \mathcal{N}\left(\bar{y}, \frac{1}{n\phi}\right)$$

$$\beta_\gamma \mid \gamma, \phi, g, y \sim \mathcal{N}\left(\frac{g}{1+g}\hat{\beta}_\gamma, \frac{g}{1+g}\frac{1}{\phi}\left[\mathbf{X}_\gamma^T \mathbf{X}_\gamma\right]^{-1}\right)$$

$$\phi \mid \gamma, y \sim \text{Gamma}\left(\frac{n-1}{2}, \frac{\text{TotalSS} - \frac{g}{1+g}\text{RegSS}}{2}\right)$$

$$p(\gamma \mid y) \propto p(y \mid \gamma)p(\gamma)$$

$$\text{TotalSS} \equiv \sum_i (y_i - \bar{y})^2 \quad \text{RegSS} \equiv \hat{\beta}_\gamma^T \mathbf{X}_\gamma^T \mathbf{X}_\gamma \hat{\beta}_\gamma$$

$$R_\gamma^2 = \frac{\text{RegSS}}{\text{TotalSS}} = 1 - \frac{\text{ErrorSS}}{\text{TotalSS}}$$

$$p(Y \mid \gamma) = C(1+g)^{\frac{n-p_\gamma-1}{2}}(1+g(1-R_\gamma^2))^{-\frac{(n-1)}{2}}$$



Find Posteriors



Continued



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.
- We can also compute posterior $\Pr[\gamma_j = 1 | Y]$, the posterior probability of including the j 'th predictor, often called marginal inclusion probability (MIP), allowing for uncertainty in the other predictors.



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.
- We can also compute posterior $\Pr[\gamma_j = 1 | Y]$, the posterior probability of including the j 'th predictor, often called marginal inclusion probability (MIP), allowing for uncertainty in the other predictors.
- Also straightforward to do model averaging once we all have posterior samples.



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.
- We can also compute posterior $\Pr[\gamma_j = 1 | Y]$, the posterior probability of including the j 'th predictor, often called marginal inclusion probability (MIP), allowing for uncertainty in the other predictors.
- Also straightforward to do model averaging once we all have posterior samples.
- The Hoff book works through one example and you can find the Gibbs sampler for doing inference there. I strongly recommend you go through it carefully!



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.
- We can also compute posterior $\Pr[\gamma_j = 1 | Y]$, the posterior probability of including the j 'th predictor, often called marginal inclusion probability (MIP), allowing for uncertainty in the other predictors.
- Also straightforward to do model averaging once we all have posterior samples.
- The Hoff book works through one example and you can find the Gibbs sampler for doing inference there. I strongly recommend you go through it carefully!
- Also paper by Liang et al (2008) JASA



Summaries

- We can run a collapsed Gibbs or MH sampler over just Γ !
- We can then compute marginal posterior probabilities $\Pr[\gamma|Y]$ for each model and select model with the highest posterior probability.
- We can also compute posterior $\Pr[\gamma_j = 1 | Y]$, the posterior probability of including the j 'th predictor, often called marginal inclusion probability (MIP), allowing for uncertainty in the other predictors.
- Also straightforward to do model averaging once we all have posterior samples.
- The Hoff book works through one example and you can find the Gibbs sampler for doing inference there. I strongly recommend you go through it carefully!
- Also paper by Liang et al (2008) JASA
- we will focus on using R packages for implementing



Examples with BAS

```
library(BAS)
data(usair, package="HH")
poll.bma = bas.lm(log(SO2) ~ temp + log(mfgfirms) +
                  log(popn) + wind +
                  precip + raindays,
                  data=usair,
                  prior="g-prior",
                  alpha=nrow(usair), #  $g = n$ 
                  n.models=2^6,
                  modelprior = uniform(),
                  method="deterministic")
```



Summaries


```
poll.bma
```

```
##  
## Call:  
## bas.lm(formula = log(SO2) ~ temp + log(mfgfirms) + log(popn) +  
##      wind + precip + rainedays, data = usair, n.models = 2^6, prior = "g-p  
##      alpha = nrow(usair), modelprior = uniform(), method = "deterministic  
##  
##  
## Marginal Posterior Inclusion Probabilities:  
##      Intercept      temp  log(mfgfirms)      log(popn)      wind  
##      1.0000      0.9755      0.7190      0.2757      0.765  
##      precip      rainedays  
##      0.5994      0.3104
```



Plots of Coefficients

```
beta = coef(poll.bma)  
par(mfrow=c(2,3)); plot(beta, subset=2:7,ask=F)
```



Summary of Coefficients

beta

```
##
## Marginal Posterior Summaries of Coefficients:
##
## Using BMA
##
## Based on the top 64 models
##
```


	post mean	post SD	post p(B != 0)
## Intercept	3.153004	0.082872	1.000000
## temp	-0.059724	0.020675	0.975504
## log(mfgfirms)	0.195716	0.177190	0.719031
## log(popn)	-0.026093	0.164277	0.275681
## wind	-0.126379	0.090777	0.765449
## precip	0.010821	0.011497	0.599380
## raindays	0.001803	0.004023	0.310357

Iterated Expectations!



Model Space Visualization

```
image(poll.bma, rotate=FALSE)
```



Bartlett's Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- What happens to Bayes Factors or posterior probabilities of γ as $g \rightarrow \infty$? (for fixed data)



Bartlett's Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- What happens to Bayes Factors or posterior probabilities of γ as $g \rightarrow \infty$? (for fixed data)
- What happens to Bayes Factor as $g \rightarrow 0$



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)
- Let $F = \frac{R_\gamma^2/p_\gamma}{(1-R_\gamma^2)/(n-1-p_\gamma)}$



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)
- Let $F = \frac{R_\gamma^2/p_\gamma}{(1-R_\gamma^2)/(n-1-p_\gamma)}$
- As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject γ_0 where F is the usual F statistic for comparing model γ to γ_0



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)
- Let $F = \frac{R_\gamma^2/p_\gamma}{(1-R_\gamma^2)/(n-1-p_\gamma)}$
- As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject γ_0 where F is the usual F statistic for comparing model γ to γ_0
- BF converges to a fixed constant $(1 + g)^{-p_\gamma/2}$ (does not go to infinity)



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)
- Let $F = \frac{R_\gamma^2/p_\gamma}{(1-R_\gamma^2)/(n-1-p_\gamma)}$
- As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject γ_0 where F is the usual F statistic for comparing model γ to γ_0
- BF converges to a fixed constant $(1 + g)^{-p_\gamma/2}$ (does not go to infinity)
- one predictor example



Information Paradox

$$\text{BF}(\gamma : \gamma_0) = (1 + g)^{(n-1-p_\gamma)/2} (1 + g(1 - R_\gamma^2))^{-(n-1)/2}$$

- Let g be a fixed constant and take n fixed imagine a sequence of data such that $R_\gamma^2 \rightarrow 1$ (increasing explained variation)
- Let $F = \frac{R_\gamma^2/p_\gamma}{(1-R_\gamma^2)/(n-1-p_\gamma)}$
- As $R_\gamma^2 \rightarrow 1$, $F \rightarrow \infty$ LR test would reject γ_0 where F is the usual F statistic for comparing model γ to γ_0
- BF converges to a fixed constant $(1 + g)^{-p_\gamma/2}$ (does not go to infinity)
- one predictor example

Information Inconsistency see Liang et al JASA 2008



Mixtures of g -priors & Information Consistency

Need $BF \rightarrow \infty$ if $R_\gamma^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges for $p_\gamma < n - 1$ (proof in Liang et al)



Mixtures of g -priors & Information Consistency

Need $BF \rightarrow \infty$ if $R_\gamma^2 \rightarrow 1 \Leftrightarrow \mathbb{E}_g[(1+g)^{-p_\gamma/2}]$ diverges for $p_\gamma < n-1$ (proof in Liang et al)

- Zellner-Siow Cauchy prior, $1/g \sim \text{Gamma}(1/2, 1/2)$)



Mixtures of g -priors & Information Consistency

Need $BF \rightarrow \infty$ if $R_\gamma^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges for $p_\gamma < n - 1$ (proof in Liang et al)

- Zellner-Siow Cauchy prior, $1/g \sim \text{Gamma}(1/2, 1/2)$
- hyper-g prior or hyper-g/n (Liang et al JASA 2008)



Mixtures of g -priors & Information Consistency

Need $BF \rightarrow \infty$ if $R_\gamma^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges for $p_\gamma < n - 1$ (proof in Liang et al)

- Zellner-Siow Cauchy prior, $1/g \sim \text{Gamma}(1/2, 1/2)$
- hyper- g prior or hyper- g/n (Liang et al JASA 2008)
- robust prior (Bayarri et al Annals of Statistics 2012)



Mixtures of g -priors & Information Consistency

Need $BF \rightarrow \infty$ if $R_\gamma^2 \rightarrow 1 \Leftrightarrow E_g[(1 + g)^{-p_\gamma/2}]$ diverges for $p_\gamma < n - 1$ (proof in Liang et al)

- Zellner-Siow Cauchy prior, $1/g \sim \text{Gamma}(1/2, 1/2)$
- hyper- g prior or hyper- g/n (Liang et al JASA 2008)
- robust prior (Bayarri et al Annals of Statistics 2012)

All have tails that behave like a Cauchy distribution (robustness)

