



Phytochemistry and Medicinal Significance of *Adhatoda vasica*: A Review

Mohammed Sadiq V P*

Kerala Agricultural University

Abstract

Adhatoda vasica (syn. *Justicia adhatoda* L.), commonly known as Vasaka or Malabar nut, is a medicinal plant of considerable therapeutic importance in traditional systems such as Ayurveda and Unani. This review presents a comprehensive summary of its phytochemical constituents and pharmacological activities. The plant exhibits diverse biological effects including anti-asthmatic, anti-inflammatory, antimicrobial, hepatoprotective, antioxidant, thrombolytic, and wound-healing properties, as demonstrated through various in vitro and in vivo studies. Major bioactive compounds such as vasicine and vasicinone contribute significantly to its pharmacological potential. This review highlights *A. vasica* as a potent source of bioactive molecules and underscores the need for translational research to facilitate its integration into modern therapeutic practice. Relevant literature on *Adhatoda vasica* was gathered from databases such as Google Scholar, PubMed, Scopus, and ScienceDirect using suitable keywords such as *Adhatoda vasica*, phytochemistry, pharmacology and vasicine.

Keywords: *Adhatoda vasica*, Ayurveda, phytochemicals, pharmacological activity, anti-asthmatic, bioactive compounds.

1. Introduction

Adhatoda vasica, commonly referred to as Vasaka or Malabar nut, is a perennial evergreen shrub revered in traditional Indian medicine for its broad spectrum of pharmacological activities. The Sanskrit term “Vasa” denotes “that which restores health.” Belonging to the family Acanthaceae, *A. vasica* is distributed across tropical and subtropical regions of South and Southeast Asia, including India, Sri Lanka, China, and Bangladesh. *Adhatoda* is widely distributed throughout India including states such as Kerala, Tamil Nadu, Andhra Pradesh, Madhya Pradesh, Uttarakhand, Punjab, Mizoram and Tripura (Sankara Rao, K. and Deepak Kumar 2025). It grows abundantly, even in marginal and wasteland areas (Shamsuddin *et al.* 2021), and also contributes to ecological restoration through phytoremediation of toxic elements such as mercury and chromium (Isha *et al.* 2025). The foliage has insect repellent activity, and the stem is used for cleaning tooth and strengthening gum (Kumar *et al.* 2016). *Adhatoda beddomie* is another important species of *Adhatoda*, predominantly distributed in Kerala. It has lower levels of alkaloids, phenolics and steroids but a higher terpenoid content than *A. vasica* (Nandhini and Ilango 2020). Vasica and Ajagandhi are two improved varieties of *A. vasica* released from Kerala agricultural university with vasicine content 2.55% and 2.46% respectively (Tomy *et al.* 2023)

In India, it is known by various vernacular names—Adusa, Arusha, or Bansa (Hindi); Adalodakam (Malayalam); Adathodai (Tamil); Adulsa (Marathi); Addasaramu (Telugu); and Adu-sogae (Kannada). Morphologically, it is a compact evergreen shrub, 2–3.5 m tall, with opposite, lanceolate leaves possessing a bitter taste. The white flowers with purple streaks occur in dense spikes, both axillary and terminal (Mehta 2016). Morphologically, it is a compact evergreen shrub, 1.5–2 m tall, with opposite decussate, petiolate, exstipulate leaf possessing a bitter taste. The flowers are either white or purple occur in dense spikes, both axillary and terminal (Sampath Kumar *et al.* 2010; Tomy *et al.* 2023).

2. Phytochemical composition

Adhatoda vasica contains a rich spectrum of phytochemicals, including alkaloids, flavonoids, phenols, steroids, carbohydrates, triterpenes, tannins, betaine, essential oils, and alkanes. Over 30 alkaloids have been identified, primarily quinazoline alkaloids such as vasicine and vasicinone, which are predominant in the leaves.

Phytochemical group	Major compounds identified	Functions	References
Alkaloids	Vasicine, vasicinone, vasicinolone, adhatodine, deoxyvasicine, hydroxyl vasicine, vasicol	Bronchodilatory, respiratory stimulant, antibacterial, antioxidant, abortifacient, wound healing and hepatoprotective functions	Soni <i>et al.</i> (2008); Kancharla <i>et al.</i> (2023); Doba and Goti (2023); Shoaib (2021)

Phytochemical group	Major compounds identified	Functions	References
Flavonoids	Apigenin, astragalin, kaempferol, quercetin, vitexin	Antioxidant and anti-inflammatory functions	Maurya and Singh (2010)
Terpenoids	α -amyrin, epitaraxerol, β -carotene	Antithyroid, antioxidant and cardioprotective functions	Maurya and Singh (2010); Shahzad <i>et al.</i> (2020); Banerji <i>et al.</i> (1999); Jha <i>et al.</i> (2012)
Steroids	β -sitosterol, α -sitosterol	Hepatoprotective, antibacterial, antioxidant and immunomodulatory functions	Roy <i>et al.</i> (2013); Jain <i>et al.</i> (1980)
Glycosides	β -glucoside galactose	Antidiabetic function	Jain <i>et al.</i> (1980)

Table 1: Phytochemical composition of *Adhatoda vasica*

and responsible for the plant's bronchodilatory and respiratory-stimulant effects. In addition, other significant alkaloids like vasicinolone, adhatodine, deoxyvasicine, hydroxyl vasicine, and vasicol contribute to the plant's therapeutic potential (Soni *et al.* 2008; Kancharla *et al.* 2023). The medicinal benefits of *Adhatoda vasica* are further supported by the presence of major flavonoids including apigenin, astragalin, kaempferol, quercetin, and vitexin, as well as terpenoids such as α -amyrin, epitaraxerol, and β -carotene (Maurya and Singh 2010; Shahzad *et al.* 2020). The essential oil extracted from the plant contains terpenoids, sesquiterpenes, and cyclic hydrocarbons. Among the major steroids, β -sitosterol and α -sitosterol are present in notable amounts.

Additionally, vitamin C and amino acids like glycine, proline, serine, and valine are found in considerable quantities (Megha *et al.* 2012). Spectrophotometric analysis has confirmed the presence of important major elements such as potassium, sodium, calcium, and magnesium, along with trace elements including zinc, copper, chromium, nickel, cobalt, cadmium, lead, manganese, and iron (Kumar *et al.* 2014). Fatty acids such as Crystalline acid, arachidic acid, linoleic acid, be-henic acid and oleic acid are also present in *Adhatoda vasica* which possess anticancerous, hepatoprotective, cardioprotective, muscle relaxant and gastroprotective properties (Singh *et al.* 2011).

3. Traditional and ethnomedicinal uses

In Ayurveda, *A. vasica* is highly valued for treating respiratory disorders including asthma, bronchitis, cough, and tuberculosis (Sampath Kumar *et al.* 2010). Its bitter-astringent taste (Tikta-Kashaya Rasa), pungent post-digestive effect (Katu Vipaka), and cooling potency

(Sheeta Virya) contribute to its therapeutic action. Various preparations such as Kvatha (decoction), Avaleha (paste/jam), Sneha (medicated oil), and Sandhana (fermented products) are traditionally employed (Gupta and Prajapati 2010).

Ancient Ayurvedic texts—Charaka Samhita, Sushruta Samhita, and Bhavaprakasha Nighantu—document its role in balancing Kapha and Pitta doshas (Kumar 2016). The roots are used to promote postpartum recovery and treat gonorrhea, while flowers are employed for their bronchodilatory and expectorant effects (Hussain and Hoq 2016; Shoaib 2021).

In the Unani system, the plant is recognized for its antispasmodic, expectorant, antipyretic, and antibiotic properties, and is prescribed for conditions such as influenza, tuberculosis, bronchitis, and gastric ulcers (Shamsi *et al.* 2019).

4. Pharmacological activities

4.1. Bronchodilator and anti-asthmatic activity

Ethanollic extracts of *A. vasica* leaves significantly increased pre-convulsion time in histamine-induced bronchospasm in guinea pigs, demonstrating bronchodilatory efficacy comparable to ketotifen (Dangi *et al.* 2015; Khandelwal *et al.* 2024). Vasicine and vasicinone act synergistically as bronchodilators, antitussives, and anti-inflammatory agents. It has expectorant activity, promoting the loosening and expulsion of mucus from the respiratory tract. The alkaloids vasicinone and vasicine have potent bronchodilator and anti-allergic activity (Dhuley 1999; Gangwar and Ghosh 2014; Jyoti *et al.* 2018).

4.2. Antibacterial and antimicrobial activity

Leaf, stem, and root extracts of *A. vasica* exhibit broad-spectrum antibacterial activity against pathogens including *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, and *Streptococcus pyogenes* (Sharma and Agarwal 2021). Compounds extracted from the stem, such as β -sitosterol, monopalmitin, vanillin, vasicinolone, and vasicinone, showed strong antibacterial activity against *Escherichia coli* and *Staphylococcus aureus* (Ibrahim *et al.* 2019). While predominantly antibacterial, certain compounds like daucosterolpalmitate and vanillic acid also exhibited antifungal activity against *Candida albicans*, *Aspergillus niger*, and *Trichoderma reesei* (Abdel-Rahman *et al.* 2017). The efficacy varies with the solvent used, with ethanolic extracts generally showing superior inhibition.

4.3. Anti-diabetic activity

Diabetes mellitus is a chronic condition marked by high blood glucose due to low insulin or insulin resistance. Methanolic extracts of *A. vasica* (50-100 mg/kg) significantly reduced blood glucose levels in alloxan-induced diabetic rats, suggesting the hypoglycemic role of flavonoids, alkaloids, phenols and saponins (Mehta *et al.* 2023; Ramachandran *et al.* 2016). Enhanced glucose uptake in rat hemidiaphragm models indicates insulin-like activity.

4.4. Anti-ulcer activity

A peptic ulcer is a painful sore in the stomach or small intestine caused by stomach acid eroding the protective mucus layer. Alcoholic, chloroform, and ether extracts of *A. vasica* (100-200 mg kg⁻¹) markedly reduced gastric acidity (Guth *et al.* 1976). The extracts significantly reduced the volume of gastric acid secretion, free and total acidity, and provided direct protective effects on the gastric mucosal lining. Furthermore, the ulcer index, a quantitative measure of gastric ulcer severity was decreased in the animals treated with the plant extracts (Vinothapooshan and Sundar 2011).

4.5. Uterotonic and abortifacient activity

Vasicine exhibits potent uterotonic activity comparable to oxytocin in human myometrial tissues (Gupta *et al.* 1979). Oral administration of ethanolic extracts (175 mg kg⁻¹) in guinea pigs induced 100% abortion when administered post-conception, indicating prostaglandin-mediated uterine stimulation (Atal 1980; Chandhoke 1982).

4.6. Antioxidant activity

Adhatoda vasica exhibits significant antioxidant activity, primarily due to its high content of phenolic and flavonoid compounds. *A. vasica* scavenge DPPH radicals effectively, with flower extracts showing the highest activity (69% inhibition at 80 g/ml) (Khan *et al.* 2019). In presence of plant extract, the Mo(IV) is found to be reduced to Mo(V) and forms ammonium molybdenum complex which can be measured using UV- visible spectrophotometer (Prathiba 2019; Rao *et al.* 2013).

4.7. Anti-inflammatory and analgesic activity

The alkaloid fraction of *A. vasica* extract exhibited strong anti-inflammatory activity in carrageenan- and formalin-induced edema models, comparable to hydrocortisone (Chakraborty and Brantner 2001; Basit *et al.* 2022). It also suppressed inflammatory cytokines such as TNF- α , IL-1 β , IL-6, and IL-8 (Amala and Sujatha 2019). Furthermore, it inhibits the histamine release from mast cells contributing to reduced allergic responses (Pawar *et al.*). Experiment conducted in formalin induced zebra fish (*Danio rerio*) has proved the antinociceptive action of *A. vasica* plant extract at concentrations of 5, 10, 15, and 20 mg/ml (Gao *et al.* 2022).

4.8. Wound-healing activity

A. vasica enhances the wound healing process by promoting the production of connective tissue components such as collagen and elastin. A one per cent methanolic extract of the aerial parts of the plant, formulated as an ointment, enhanced wound contraction and collagen synthesis in excision models in mice. Treated wounds showed higher hydroxyproline content and increased granulation tissue formation, confirming its tissue-repairing and antioxidant role (Subhashini and Arunachalam 2011; Doba and Goti 2023).

4.9. Anti-tubercular activity

Two semi-synthetic derivatives of vasicine, such as bromhexine and ambroxol, exhibit mucolytic and antimycobacterial properties by accumulating in macrophages and inhibiting bacterial fatty acid synthesis (Grange and Snell 1996; Narimanyan *et al.* 2005; Jha *et al.* 2012). Quinazoline alkaloids of *A. vasica* show promise as leads for anti-TB drug development.

4.10. Thrombolytic activity

Methanolic extracts demonstrated moderate thrombolytic activity (53.23% clot lysis), attributed to alkaloids, tannins, flavonoids, and saponins (Shahriar 2023).

4.11. Hepatoprotective activity

Adhatoda vasica exhibits hepatoprotective action by reducing oxidative stress and protecting the liver from damage induced by various toxins. In CCl₄-induced hepatotoxic rats, *A. vasica* extract (250–500 mg kg⁻¹) significantly reduced elevated serum AST, ALT, ALP, and bilirubin levels, indicating hepatoprotection through antioxidant mechanisms (Afzal *et al.* 2013; Kumar *et al.* 2015).

4.12. Insecticidal and antifeedant activity

Adhatoda has been used as an insect repellent for decades in India. Its leaves are applied to control pests of oilseeds (Singh and Tiwari 2016). The alkaloid vasicinol has an antifertility effect on some insect species by blocking their oviducts (Saxena *et al.* 1986). Studies have shown the effectiveness of crude leaf extract of *Adhatoda* against pests like cabbage aphid (*Brevicoryne brassicae*) and pink hibiscus mealy bug (*Maconellicoccus hirsutus*) (Haifa and Ali 2016). The 24-hour LC₅₀ of ethanol and water extracts of *Adhatoda* leaves against *Maconellicoccus hirsutus* are 25.70 ppm and 39.81 ppm, respectively. In the case of *Brevicoryne brassicae*, the acetonic and ethanolic extracts showed average killing rates of 57.2% and 47.1%, respectively. *Adhatoda* has also been studied for its larvicidal effects on the filariasis vector *Culex quinquefasciatus* and the dengue vector *Aedes aegypti* (Thanigaivel *et al.* 2012). The antifeedant effect of methanolic extract on various pest species such as cotton leaf worm (*Spodoptera littoralis*) and rice brown plant hopper has also been reported (Sadek 2003; Rana *et al.* 2015).

Target insect	Solvent	Bioassay parameter	Quantitative result	Reference
Cabbage aphid (<i>Brevicoryne brassicae</i>)	Ethanolic leaf extract	LC ₅₀	25.70 ppm	Haifa and Ali (2016)
Cabbage aphid (<i>Brevicoryne brassicae</i>)	Aqueous leaf extract	LC ₅₀	39.81 ppm	Haifa and Ali (2016)
Pink hibiscus mealy bug (<i>Maconellicoccus hirsutus</i>)	Ethanolic leaf extract	LC ₅₀	25.70 ppm	Kalitha <i>et al.</i> (2021)

Target insect	Solvent	Bioassay parameter	Quantitative result	Reference
Pink hibiscus mealy bug (<i>Maconellicoccus hirsutus</i>)	Aqueous leaf extract	LC50	39.81 ppm	Kalitha <i>et al.</i> (2021)
Southern house mosquito (<i>Culex quinquefasciatus</i>)	Methanolic leaf extract	LC50	56.13 ppm	Thanigaivel <i>et al.</i> (2012)

Table 2: Insecticidal activity of *Adhatoda vasica* against selected insects

4.13. Anti-cancer activity

Hexane and methanolic extracts showed cytotoxic effects against leukemia (MOLM-14, NB-4) and solid tumor cell lines (HeLa, MCF-7, HepG2) with dose-dependent inhibition of proliferation (Balachandran *et al.* 2017; Nirmala *et al.* 2019; Nikhitha *et al.* 2021). The mechanism involves apoptosis induction and p53/p21 gene modulation.

4.14. Immunomodulatory activity

Methanolic and chloroform extracts enhanced immune response in rats by increasing neutrophil adhesion, macrophage activity, and delayed type hypersensitivity (Vinothapreeshan and Sundar 2011; Sutare and Kareppa 2020).

4.15. Anti-mutagenic activity

Plant extracts reduced cadmium-induced chromosomal aberrations and oxidative stress markers in mice (Jahangir *et al.* 2006). Vasicine also showed strong inhibition of 2-aminofluorene-induced mutagenicity in *Salmonella typhimurium* strains (Kaur *et al.* 2015). Ultra-high-performance liquid chromatography (UHPLC) analysis revealed the presence of polyphenolic compounds and flavonoids, which may be responsible for the bioprotective activity of the plant extract. The extract was found to have protective antioxidant activity against free radicals, superoxides, and hydrogen peroxide radicals, thereby conferring antimutagenic potential.

5. Conclusion

Adhatoda vasica is a medicinally significant plant possessing a broad spectrum of pharmacological activities supported by its diverse phytochemical composition. Key bioactive alkaloids such as vasicine and vasicinone confer anti-asthmatic, anti-inflammatory, antimicrobial, hepatoprotective, antioxidant, and immunomodulatory properties. It has great prospects due to these pharmacological properties, suitability to waste lands and low input requirements. However, the absence of improved varieties, along with the lack of standardized cultivation practices and extraction or quality control protocols, remains a major limitation. The

biosynthetic pathways and regulatory factors of major chemical constituents remain inadequately studied. Although preclinical findings are promising, comprehensive clinical trials and standardization of dosage, formulation, and safety parameters are imperative. Continued multidisciplinary research could facilitate the incorporation of *A. vasica* derived compounds into modern pharmacotherapy, validating its traditional use as a potent natural therapeutic agent.

References

- Abdel-Rahman SA, El-Gohary NS, El-Bendary ER, El-Ashry SM, Shaaban MI (2017). "Synthesis, antimicrobial, antiquorum-sensing, antitumor and cytotoxic activities of new series of cyclopenta(hepta)[b]thiophene and fused cyclohepta[b]thiophene analogs." *European Journal of Medicinal Chemistry*, **140**, 200–211.
- Afzal U, Gulfraz M, Hussain S, Malik F, Maqsood S, Shah I, Mahmood S (2013). "Hepato-protective effects of *Justicia adhatoda* L. against carbon tetrachloride (CCl₄) induced liver injury in Swiss albino mice." *African Journal of Pharmacy and Pharmacology*, **7**(1), 8–14.
- Amala R, Sujatha S (2019). "Presence of pyrroloquinazoline alkaloid in *Adhatoda vasica* attenuates inflammatory response through the downregulation of pro-inflammatory mediators in LPS stimulated RAW 264.7 macrophages." *BioImpacts*, **11**, 15–22.
- Atal CK (1980). *Chemistry and pharmacology of vasicine, a new oxytocic and abortifacient*. Regional Research Laboratory, CSIR, Jammu-Tawi.
- Balachandran C, Arun Y, Sangeetha B, Duraipandian V, Awale S, Emi N, Ignacimuthu S, Perumal PT (2017). "In vitro and in vivo anticancer activity of 2-acetyl-benzylamine isolated from *Adhatoda vasica* L. leaves." *Biomedicine and Pharmacotherapy*, **93**, 796–806.
- Banerji J, Chatterjee A, Saha M, Dhara KP, Kanrar S, Mukherjee P, Neuman A, Prange T (1999). "Reactions and rearrangements of triterpenoids-3-epitaraxerol and its transformation products." *Indian Journal of Chemistry Section B*, **38**(12), 1322–1330. doi: [10.1002/chin.200019166](https://doi.org/10.1002/chin.200019166).
- Basit A, Shuitian T, Khan A, Khan SM, Shahzad R, Khan MZ, Khan M (2022). "Anti-inflammatory and analgesic potential of leaf extract of *Justicia adhatoda* (L.) (Acanthaceae) in carrageenan and formalin-induced models by targeting oxidative stress." *Biomedicine and Pharmacotherapy*, **153**, 113322.
- Chakraborty A, Brantner AH (2001). "Study of alkaloids from *Adhatoda vasica* Nees on their antiinflammatory activity." *Phytotherapy Research*, **15**(6), 532–534.
- Chandhoke N (1982). "Vasicine, the alkaloid of *Adhatoda vasica*." *Indian Drugs*, **24**(9), 425–426.
- Dangi A, Patel S, Yaduvanshi PS (2015). "Phytochemical screening and assessment of *Adhatoda vasica* (leaf) for antiasthmatic activity." *Panacea Journal of Pharmacy and Pharmaceutical Sciences*, **4**(3), 680–704.
- Dhuley JN (1999). "Antitussive effect of *Adhatoda vasica* extract on mechanical or chemical stimulation-induced coughing in animals." *Journal of Ethnopharmacology*, **67**(3), 361–365.
- Doba JP, Goti D (2023). "Phytopharmacological evaluation of *Adhatoda vasica*." *World Journal of Pharmaceutical Research*, **12**(9), 759–780.
- Gangwar AK, Ghosh AK (2014). "Medicinal uses and pharmacological activity of *Adhatoda vasica*." *International Journal of Herbal Medicine*, **2**(1), 88–91.

- Gao L, Cui S, Huang Z, Cui H (2022). “Adhatoda vasica is effective in relieving pain through modulation of inflammation.” *Pharmacognosy Magazine*, **18**(79), 593–599. doi:10.4103/pm.pm_114_22.
- Grange JM, Snell NJ (1996). “Activity of bromhexine and ambroxol, semi-synthetic derivatives of vasicine from the Indian shrub *Adhatoda vasica*, against *Mycobacterium tuberculosis* in vitro.” *Journal of Ethnopharmacology*, **50**(1), 49–53.
- Gupta A, Prajapati PK (2010). “A clinical review of different formulations of Vasa (*Adhatoda vasica*) on Tamaka Shwasa (asthma).” *Ayu*, **31**(4), 520–524.
- Gupta OP, Wakhloo BL, Sharma ML, Atal CK (1979). “Vasicine—A potent uterine stimulant—Studies on human myometrium.” *Journal of Obstetrics and Gynaecology of India*, **29**(5), 935–938.
- Guth PH, Paulsen G, Lynn D, Aures D (1976). “Mechanism of prevention of aspirin-induced gastric lesions by bile duct ligation in the rat.” *Gastroenterology*, **71**(5), 750–753.
- Haifa NM, Ali SM (2016). “Insecticidal effect of crude plant extract of *Adhatoda vasica* against *Brevicoryne brassicae*.” *World Journal of Experimental Biosciences*, **4**(1), 49–52.
- Hussain MT, Hoq MO (2016). “Therapeutic use of *Adhatoda vasica*.” *Asian Journal of Medical and Biological Research*, **2**(2), 156–163.
- Ibrahim D, Ba FMA, Elgaml A, Gedara SR, Gohar AA (2019). “Antimicrobial, Antiquorum-sensing and Ex-vivo antispasmodic activity of *Adhatoda vasica*.” *Journal of Pharmaceutical Research International*, **31**(6), 1–16.
- Isha, Kumar P, Singh AN (2025). “An overview of *Justicia adhatoda*: A medicinal plant but native invader in India.” *Conservation*, **5**(1), 2.
- Jahangir T, Khan TH, Prasad L, Sultana S (2006). “Reversal of cadmium chloride-induced oxidative stress and genotoxicity by *Adhatoda vasica* extract in Swiss albino mice.” *Biological Trace Element Research*, **111**(1–3), 217–228.
- Jain M, Koul S, Atal CK (1980). “Novel nor-harmal alkaloid from *Adhatoda vasica*.” *Phytochemistry*, **19**(9), 1880–1882. doi:10.1016/S0031-9422(00)83845-5.
- Jha DK, Panda L, Lavanya P, Ramaiah S, Anbarasu A (2012). “Detection and confirmation of alkaloids in leaves of *Justicia adhatoda* and bioinformatics approach to elicit its anti-tuberculosis activity.” *Applied Biochemistry and Biotechnology*, **168**(5), 980–990.
- Jyoti R, Shikha P, Bihari KV, Kumar SU (2018). “Role of *Adhatoda vasica* in management of respiratory disorders.” *Environment Conservation Journal*, **19**(3), 73–75.
- Kalitha PPM, Sundaralingam MS, Sukumaran A, Mano C (2021). “Insecticidal effect of *Adhatoda vasica* (leaf) and *Trigonella foenum-graecum* (seed) extracts against mealy bugs (*Maconellicoccus hirsutus*) on *Hibiscus rosa-sinensis* plant.” *Asian Journal of Biological and Life Sciences*, **10**(2), 500–506. doi:10.5530/ajbls.2021.10.66.

- Kancharla B, Singh R, Gajendra TA, Ramakrishna K, Singh SK, Kumar A, Hemalatha S (2023). "Vasicinone, a pyrroloquinazoline alkaloid from *Adhatoda vasica* Nees enhances memory and cognition by inhibiting cholinesterases in Alzheimer's disease." *Phytomedicine Plus*, **3**(2), 100439.
- Kaur A, Kaur D, Arora S (2015). "Evaluation of antioxidant and anti-mutagenic potential of *Justicia adhatoda* leaves extract." *African Journal of Biotechnology*, **14**(21), 1807–1819.
- Khan AM, Bhadauria S, Yadav R (2019). "Phytochemical screening and antioxidant activity of extract of different parts of *Adhatoda vasica*." *Research Journal of Pharmacology and Technology*, **12**(12), 5699–5705.
- Khandelwal P, Wadhwani BD, Rao RS, Mali D, Vyas P, Kumar T, Nair R (2024). "Exploring the pharmacological and chemical aspects of pyrrolo-quinazoline derivatives in *Adhatoda vasica*." *Heliyon*, **10**(4), e25727.
- Kumar M, Dandapat S, Kumar A, Sinha MP (2014). "Pharmacological screening of leaf extract of *Adhatoda vasica* for therapeutic efficacy." *Global Journal of Pharmacology*, **8**(4), 494–500.
- Kumar M, Dandapat S, Sinha MP (2015). "Hepatoprotective activity of *Adhatoda vasica* and *Vitex negundo* leaf extracts against carbon tetrachloride induced hepatotoxicity in rats." *Advances in Biological Research*, **9**(4), 242–246.
- Kumar N (2016). "Pharmaceutical attributes of vasa (*Adhatoda vasica* Linn.)—A review." *World Journal of Pharmaceutical Research*, **5**(4), 437–455.
- Kumar V, Kumar R, Singh V (2016). "*Adhatoda vasica*: A traditional use cum health benefit." *Ramnathakrishn*, **11**(1), 82–83.
- Maurya S, Singh D (2010). "Quantitative analysis of flavonoids in *Adhatoda vasica* Nees extracts." *Der Pharma Chemica*, **2**(5), 242–246.
- Megha J, Palathingal T, Bhagat SS (2012). "Effect of environmental conditions on the Vitamin C and essential oil content of *Adhatoda vasica* growing in the various regions of Kerala and Maharashtra." *International Journal of Scientific Research*, **3**(8), 31–33.
- Mehta J (2016). "Phenology of *Adhatoda vasica* a multifarious useful medicinal plant." *International Journal of Applied Research*, **2**(7), 791–794.
- Mehta RK, Thapa R, Chaudhary MK (2023). "Evaluation of anti-diabetic activity of *Justicia adhatoda* (Linn.) leaves in diabetic Wistar rats." *Journal of Universal College of Medical Sciences*, **11**(1), 50–54.
- Nandhini S, Ilango K (2020). "Comparative study on pharmacognostical, phytochemical investigations and quantification of vasicine content in the extracts of *Adhatoda vasica* Nees and *Adhatoda beddomei* C. B. Clarke." *Pharmacognosy Journal*, **12**(4), 884–896. doi:10.5530/pj.2020.12.126.
- Narimanyan M, Badalyan M, Panosyan V, Gabrielyan E, Panossian A, Wikman G (2005). "Randomized trial of a fixed combination (KanJang) of herbal extracts containing *Adhatoda vasica*, *Echinacea purpurea* and *Eleutherococcus senticosus* in patients with upper respiratory tract infections." *Phytomedicine*, **12**(8), 539–547.

- Nikhitha JN, Swathy KS, Chandran RP (2021). "In vitro anticancer activity of ethanol extract of *Adhatoda vasica* Nees on human ovarian cancer cell lines." *Journal of Genetic Engineering and Biotechnology*, **19**(1), 116.
- Nirmala S, Sabapathi PN, Sudhakar M, Bathula N, Sravanthi Y (2019). "Investigation of In vitro Anti-cancer property of *Adhatoda vasica* in Hela, HepG2, MCF-7, MDAMB-231 Cell Lines." *Research Journal of Pharmacognosy and Phytochemistry*, **11**(4), 212–216.
- Prathiba HD (2019). "Free radical scavenging potential of leaf of *Adhatoda vasica* and *Lantana camara* by DPPH assay." *International Journal of Pharmaceutical and Biological Sciences*, **9**(Special Issue 1), 34–39.
- Ramachandran V, Rajendran K, Ramalingam G (2016). "In-vitro study of *Adhatoda vasica* Nees on glucose uptake by isolated rat hemi-diaphragm." *Journal of Pharmaceutical and Medical Chemistry*, **2**(2), 141–143.
- Rana MS, Bachchu MAA, Hossain MA, Ara R, Rahman H (2015). "Effect of some indigenous plant extracts against brown plant hopper, *Nilaparvata lugens* (Stal.) (Hemiptera: Delphacidae)." *Bangladesh Journal of Entomology*, **25**(1), 15–31.
- Rao KVB, Munjal M, Patnayak A, Karthik L, Kumar G (2013). "Phytochemical composition, antioxidant, antimicrobial and cytotoxic potential of methanolic extracts of *Adhatoda vasica* (Acanthaceae)." *Research Journal of Pharmacology and Technology*, **6**(9), 1004–1009.
- Roy DC, Shaik MM, Faruquee H (2013). "A brief review on phytochemistry and pharmacological properties of *Adhatoda vasica*." *Journal of Tropical Medicinal Plants*, **14**, 115–124.
- Sadek M (2003). "Antifeedant and toxic activity of *Adhatoda vasica* leaf extract against *Spodoptera littoralis* (Lep., Noctuidae)." *Journal of Applied Entomology*, **127**(7), 396–404.
- Sampath Kumar KP, Bhowmik D, Chiranjib, Tiwari P, Rawat L (2010). "Indian traditional herbs *Adhatoda vasica* and its medicinal application." *Journal of Chemical and Pharmaceutical Research*, **2**(1), 240–245.
- Sankara Rao, K and Deepak Kumar (2025). "India Flora Online: *Justicia adhatoda*." <http://indiaflora-ces.iisc.ac.in/plants.php?name=Justiciaadhatoda>. Accessed on 13 December 2025.
- Saxena BP, Tikku K, Atal CK, Koul O (1986). "Insect antifertility and antifeedant allelochemicals in *Adhatoda vasica*." *Insect Science and Its Application*, **7**(4), 489–493.
- Shahriar M (2023). "Phytochemical screenings and thrombolytic activity of the leaf extracts of *Adhatoda vasica*." *The Experiment*, **7**(4), 438–461.
- Shahzad Q, Sammi S, Mehmood A, Naveed K (2020). "Phytochemical analysis and antimicrobial activity of *Adhatoda vasica* leaves." *Pure and Applied Biology*, **9**(2), 1654–1661.
- Shamsi Y, Khan R, Nikhat S (2019). "Clinically significant improvement in a case of bronchial asthma with Unani medicine: A case report." *Traditional and Integrative Medicine*, **4**(3), 130–136.

- Shamsuddin T, Alam MS, Junaid M, Akter R, Hosen SMZ, Ferdousy S, Mouri NJ (2021). “Adhatoda vasica (Nees.): A Review on its Botany, Traditional uses, Phytochemistry, Pharmacological Activities and Toxicity.” *Mini Reviews in Medicinal Chemistry*, **21**(14), 1925–1964.
- Sharma A, Agarwal M (2021). “Antimicrobial activity of Adhatoda vasica Nees.” *World Journal of Pharmaceutical Research*, **10**(11), 1328–1342.
- Shoaib A (2021). “A systematic ethnobotanical review of Adhatoda vasica (L.) Nees.” *Cellular and Molecular Biology*, **67**(4), 248–263.
- Singh PT, Singh MO, Singh HB (2011). “Adhatoda vasica Nees: Phytochemical and pharmacological profile.” *The Natural Products Journal*, **1**(1), 29–37. doi:10.2174/2210315511101010029.
- Singh R, Tiwari A (2016). “Adhatoda vasica: A miracle and boon for asthmatic people – A review.” *Research Journal of Pharmacognosy and Phytochemistry*, **8**(4), 242–244.
- Soni S, Anandjiwala S, Patel G, Rajani M (2008). “Validation of different methods of preparation of Adhatoda vasica leaf juice by quantification of total alkaloids and vasicine.” *Indian Journal of Pharmaceutical Sciences*, **70**(1), 36–42.
- Subhashini S, Arunachalam KD (2011). “Investigations on the phytochemical activities and wound healing properties of Adhatoda vasica leave in Swiss albino mice.” *African Journal of Plant Science*, **5**(2), 133–145.
- Sutare MS, Kareppa BM (2020). “Pharmacological activities of potent medicinal plant Adhatoda zeylanica Medik.: A review.” *International Journal of Life Science Review*, **3**(2), 1–5.
- Thanigaivel A, Chandrasekaran R, Revathi K, Nisha S, Sathish-Narayanan S, Kirubakaran SA, Senthil-Nathan S (2012). “Larvicidal efficacy of Adhatoda vasica (L.) Nees against the bancroftian filariasis vector Culex quinquefasciatus Say and dengue vector Aedes aegypti L. in in vitro condition.” *Parasitology Research*, **110**(5), 1993–1999.
- Tomy TP, Ansary PY, Oommen SM, Shincymol VV (2023). “Taxonomical identification of different plant sources of Vaasa (Adhatoda spp.)” *Kerala Journal of Ayurveda*, **24**(3), 1–5. doi:10.55718/kja.212.
- Vinothapooshan G, Sundar G (2011). “Immunomodulatory activity of various extracts of Adhatoda vasica Linn. in experimental rats.” *African Journal of Pharmacy and Pharmacology*, **5**(3), 308–316.
- Vinothapreeshan G, Sundar K (2011). “Anti-ulcer activity of Adhatoda vasica leaves against gastric ulcer in rats.” *Journal of Global Pharma Technology*, **3**(2), 7–13.

! Publication & Reviewer Details**Publication Information**

- **Submitted:** 15 November 2025
 - **Accepted:** 20 December 2025
 - **Published (Online):** 23 December 2025
-

Reviewer Information

- **Reviewer 1:**
Anonymous
- **Reviewer 2:**
Anonymous

i Disclaimer/Publisher's Note

The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). The publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#), which permits non-commercial use, sharing, and reproduction in any medium, provided the original work is properly cited and no modifications or adaptations are made.

Affiliation:

Mohammed Sadiq V P*
B.Sc. (Hons) Agriculture
College of Agriculture, Vellayani
Thiruvananthapuram, Kerala India
E-mail: sadiqbinsidhee@gmail.com