# PROBABILITY THEORY LECTURE 2

Per Sidén

Division of Statistics

Dept. of Computer and Information Science
Linköping University

## **OVERVIEW LECTURE 2**

- Conditional distributions
- ► Conditional expectation, conditional variance
- ► Distributions with random parameters and the Bayesian approach
- Regression and Prediction

## CONDITIONAL DISTRIBUTIONS

▶ For events [if P(B) > 0]

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

▶ A and B are **independent** if and only if P(A|B) = P(A).

#### CONDITIONAL DISTRIBUTIONS

▶ For events [if P(B) > 0]

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- ▶ A and B are **independent** if and only if P(A|B) = P(A).
- ► For discrete random variables

$$p_{Y|X=x}(y) = p(Y = y|X = x) = \frac{p_{X,Y}(x,y)}{p_X(x)}$$
$$p_{Y|X=x}(y) = \frac{p_{X,Y}(x,y)}{\sum_{y} p_{X,Y}(x,y)}.$$

#### CONDITIONAL DISTRIBUTIONS

▶ For events [if P(B) > 0]

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- ▶ A and B are **independent** if and only if P(A|B) = P(A).
- ► For discrete random variables

$$p_{Y|X=x}(y) = p(Y = y|X = x) = \frac{p_{X,Y}(x,y)}{p_X(x)}$$
$$p_{Y|X=x}(y) = \frac{p_{X,Y}(x,y)}{\sum_{Y} p_{X,Y}(x,y)}.$$

► For continuous random variables

$$f_{Y|X=x}(y) = \frac{f_{X,Y}(x,y)}{f_X(x)}$$

$$f_{Y|X=x}(y) = \frac{f_{X,Y}(x,y)}{\int_{-\infty}^{\infty} f_{X,Y}(x,z)dz}$$

## CONDITIONAL EXPECTATION

▶ Conditional expectation of Y given X = x is

$$E(Y|X=x) = \begin{cases} \sum_{y} y \cdot p_{Y|X=x}(y) & \text{if Y is discrete} \\ \int_{-\infty}^{\infty} y \cdot f_{Y|X=x}(y) dy & \text{if Y is continuous} \end{cases}$$

#### CONDITIONAL EXPECTATION

▶ Conditional expectation of Y given X = x is

$$E(Y|X=x) = \begin{cases} \sum_{y} y \cdot p_{Y|X=x}(y) & \text{if } Y \text{is discrete} \\ \int_{-\infty}^{\infty} y \cdot f_{Y|X=x}(y) dy & \text{if } Y \text{is continuous} \end{cases}$$

▶ If h(x) = E(Y|X = x), note that h(X) = E(Y|X) is a random variable that only depends on X.

#### CONDITIONAL EXPECTATION

▶ Conditional expectation of Y given X = x is

$$E(Y|X=x) = \begin{cases} \sum_{y} y \cdot p_{Y|X=x}(y) & \text{if } Y \text{is discrete} \\ \int_{-\infty}^{\infty} y \cdot f_{Y|X=x}(y) dy & \text{if } Y \text{is continuous} \end{cases}$$

- ▶ If h(x) = E(Y|X = x), note that h(X) = E(Y|X) is a random variable that only depends on X.
- ► Ex. 2.1 page 33.  $X \sim U(0,1)$ ,  $Y|X = x \sim U(0,x)$ .

▶ Theorem 2.1. Law of iterated expectation.

$$E[E(Y|X)] = E(Y)$$

► Theorem 2.1. Law of iterated expectation.

$$E\left[E(Y|X)\right] = E(Y)$$

Note that the *inner expectation* (E(Y|X)) is with respect to  $f_{Y|X}(y)$ , while the *outer expectation* is with respect to  $f_X(x)$ .

► Theorem 2.1. Law of iterated expectation.

$$E\left[E(Y|X)\right] = E(Y)$$

- Note that the *inner expectation* (E(Y|X)) is with respect to  $f_{Y|X}(y)$ , while the *outer expectation* is with respect to  $f_X(x)$ .
- ► The law of iterated expectation is an "expectation version" of the law of total probability.

► Theorem 2.1. Law of iterated expectation.

$$E[E(Y|X)] = E(Y)$$

- Note that the *inner expectation* (E(Y|X)) is with respect to  $f_{Y|X}(y)$ , while the *outer expectation* is with respect to  $f_X(x)$ .
- ► The law of iterated expectation is an "expectation version" of the law of total probability.
- ightharpoonup E(Y|X) = E(Y) if X and Y are independent.

▶ Conditional variance of Y given X = x is

$$Var(Y|X = x) = E[(Y - E(Y|X = x))^{2}|X = x]$$

▶ Conditional variance of Y given X = x is

$$Var(Y|X = x) = E[(Y - E(Y|X = x))^{2}|X = x]$$

Note that v(X) = Var(Y|X) is a random variable that only depends on X.

▶ Conditional variance of Y given X = x is

$$Var(Y|X = x) = E[(Y - E(Y|X = x))^{2}|X = x]$$

- Note that v(X) = Var(Y|X) is a random variable that only depends on X.
- ► Corollary 2.3.1

$$Var(Y) = E[Var(Y|X)] + Var[E(Y|X)]$$

▶ Conditional variance of Y given X = x is

$$Var(Y|X = x) = E[(Y - E(Y|X = x))^{2}|X = x]$$

- Note that v(X) = Var(Y|X) is a random variable that only depends on X.
- ► Corollary 2.3.1

$$Var(Y) = E[Var(Y|X)] + Var[E(Y|X)]$$

Note the naive version Var(Y) = E[Var(Y|X)] misses the uncertainty in Y that comes from not knowing X in E(Y|X).

▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.

- ▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.
- ► Example 1:
  - ▶  $X|N = n \sim Bin(n, p)$  and  $N \sim Po(\lambda)$ .
  - If the number of potential bidders in an auction is N=n and each of them bids with probability p, then  $X \sim Bin(n,p)$  bids will be placed.
  - ▶ The number of potential bidders is uncertain,  $N \sim Po(\lambda)$ .

- ▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.
- ► Example 1:
  - ▶  $X|N = n \sim Bin(n, p)$  and  $N \sim Po(\lambda)$ .
  - ▶ If the number of potential bidders in an auction is N = n and each of them bids with probability p, then  $X \sim Bin(n, p)$  bids will be placed.
  - ▶ The number of potential bidders is uncertain,  $N \sim Po(\lambda)$ .
  - ▶ The marginal distribution for X is  $Po(\lambda \cdot p)$

- ▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.
- ► Example 1:
  - ▶  $X|N = n \sim Bin(n, p)$  and  $N \sim Po(\lambda)$ .
  - ▶ If the number of potential bidders in an auction is N = n and each of them bids with probability p, then  $X \sim Bin(n, p)$  bids will be placed.
  - ▶ The number of potential bidders is uncertain,  $N \sim Po(\lambda)$ .
  - ▶ The marginal distribution for X is  $Po(\lambda \cdot p)$

- ▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.
- ► Example 1:
  - ▶  $X|N = n \sim Bin(n, p)$  and  $N \sim Po(\lambda)$ .
  - If the number of potential bidders in an auction is N=n and each of them bids with probability p, then  $X \sim Bin(n,p)$  bids will be placed.
  - ▶ The number of potential bidders is uncertain,  $N \sim Po(\lambda)$ .
  - ▶ The marginal distribution for X is  $Po(\lambda \cdot p)$
- ► Example 2:
  - $X|(\sigma^2=1/\lambda)\sim N(0,1/\lambda)$  and  $\lambda\sim\Gamma\left(\frac{n}{2},\frac{2}{n}\right)$ , then  $X\sim t(n)$ .
  - ► X is daily stock market returns.  $X|\lambda \sim N(0, 1/\lambda)$ , where  $1/\lambda$  is the daily variance.

- ▶  $X|\theta \sim f_X(x;\theta)$  and  $\theta$  is a random variable.
- ► Example 1:
  - ▶  $X|N = n \sim Bin(n, p)$  and  $N \sim Po(\lambda)$ .
  - If the number of potential bidders in an auction is N=n and each of them bids with probability p, then  $X \sim Bin(n, p)$  bids will be placed.
  - ▶ The number of potential bidders is uncertain,  $N \sim Po(\lambda)$ .
  - ▶ The marginal distribution for X is  $Po(\lambda \cdot p)$
- ► Example 2:
  - $ilde{X}|(\sigma^2=1/\lambda)\sim N(0,1/\lambda) \text{ and } \lambda\sim \Gamma\left(\frac{n}{2},\frac{2}{n}\right), \text{ then } X\sim t(n).$
  - ► X is daily stock market returns.  $X|\lambda \sim N(0,1/\lambda)$ , where  $1/\lambda$  is the daily variance.
  - ► The daily variance varies from day to day according to  $\lambda \sim \Gamma\left(\frac{n}{2}, \frac{2}{n}\right)$ . Turbulent day: realization of  $\lambda$  is very small.

 $\triangleright$   $X_n$ =number of heads after n tosses.

$$X_n|P=p\sim Bin(n,p)$$

 $\triangleright$   $X_n$ =number of heads after n tosses.

$$X_n|P=p\sim Bin(n,p)$$

▶ Prior distribution:  $P \sim U(0, 1)$ .

 $\triangleright$   $X_n$ =number of heads after n tosses.

$$X_n|P=p\sim Bin(n,p)$$

- ▶ Prior distribution:  $P \sim U(0, 1)$ .
- ▶ Posterior distribution:  $P|(X_n = k) \sim Beta(k+1, n+1-k)$ .

 $\triangleright$   $X_n$ =number of heads after n tosses.

$$X_n|P=p\sim Bin(n,p)$$

- ▶ Prior distribution:  $P \sim U(0, 1)$ .
- ▶ Posterior distribution:  $P|(X_n = k) \sim Beta(k+1, n+1-k)$ .
- ► Marginal of X<sub>n</sub>

$$X_n \sim U(\{0, 1, 2, ..., n\})$$

▶ Conditional of  $X_{n+1}$  given  $X_n$  and p

$$P(X_{n+1} = k+1|X_n = k, P = p) = p$$

▶ Conditional of  $X_{n+1}$  given  $X_n$  and p

$$P(X_{n+1} = k+1|X_n = k, P = p) = p$$

▶ Conditional of  $X_{n+1}$  given  $X_n$ 

$$P(X_{n+1} = k+1|X_n = k) = \frac{k+1}{n+2}$$

ightharpoonup Conditional of  $X_{n+1}$  given  $X_n$  and p

$$P(X_{n+1} = k+1|X_n = k, P = p) = p$$

 $\triangleright$  Conditional of  $X_{n+1}$  given  $X_n$ 

$$P(X_{n+1} = k+1|X_n = k) = \frac{k+1}{n+2}$$

► Coin flips are no longer independent when *p* is uncertain and we learn about *p* from data.

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

► The regression function

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

▶ Predictor:  $\hat{Y} = d(X)$ .

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

- ▶ Predictor:  $\hat{Y} = d(X)$ .
- ▶ Linear predictor  $d(\mathbf{X}) = a_0 + a_1 X_1 + ... a_n X_n$ .

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

- ▶ Predictor:  $\hat{Y} = d(X)$ .
- ▶ Linear predictor  $d(\mathbf{X}) = a_0 + a_1 X_1 + ... a_n X_n$ .
- ► Expected quadratic prediction error:  $E[Y d(X)]^2$

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

- ▶ Predictor:  $\hat{Y} = d(X)$ .
- ▶ Linear predictor  $d(\mathbf{X}) = a_0 + a_1 X_1 + ... a_n X_n$ .
- ▶ Expected quadratic prediction error:  $E[Y d(X)]^2$
- ▶ The **best predictor** of Y [minimizes expected quadratic prediction error] is the regression function E(Y|X=x).

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

- ▶ Predictor:  $\hat{Y} = d(X)$ .
- ▶ Linear predictor  $d(\mathbf{X}) = a_0 + a_1 X_1 + ... a_n X_n$ .
- ▶ Expected quadratic prediction error:  $E[Y d(X)]^2$
- ▶ The **best predictor** of Y [minimizes expected quadratic prediction error] is the regression function E(Y|X=x).
- ► Best linear predictor least squares:

$$\hat{Y} = \mu_y + \rho \frac{\sigma_y}{\sigma_x} (X - \mu_x)$$

► The regression function

$$h(\mathbf{x}) = h(x_1, ..., x_n) = E(Y|X_1 = x_1, ..., X_n = x_n) = E(Y|\mathbf{X} = \mathbf{x})$$

- ▶ Predictor:  $\hat{Y} = d(X)$ .
- ▶ Linear predictor  $d(\mathbf{X}) = a_0 + a_1 X_1 + ... a_n X_n$ .
- ▶ Expected quadratic prediction error:  $E[Y d(X)]^2$
- ▶ The **best predictor** of Y [minimizes expected quadratic prediction error] is the regression function E(Y|X=x).
- ► Best linear predictor least squares:

$$\hat{Y} = \mu_y + \rho \frac{\sigma_y}{\sigma_x} (X - \mu_x)$$

▶ When (X, Y) is jointly normal, E(Y|X = x) is linear. For other distributions, this is not true in general.