LINKÖPINGS UNIVERSITET Institutionen för datavetenskap Avdelningen för statistik Per Sidén

 $\begin{array}{c} 2016\text{--}10\text{--}18 \\ \text{Probability Theory, 6 hp} \\ 732\text{A}63 \end{array}$

Exam in Probability Theory, 6 credits

Exam time: 8-12

Allowed: Pocket calculator.

Table with common formulas and moment generating functions (distributed with the exam).

Table of integrals (distributed with the exam).

Table with distributions from Appendix B in the course book (distributed with the exam).

Examinator: Mattias Villani.

Assisting teacher: Per Sidén, phone 0704-977175

Grades: Maximum is 20 points.

A: 19 points
B: 17 points
C: 14 points
D: 12 points
E: 10 points
F: <10 points

- Write clear and concise answers to the questions.
- Make sure to specify the definition region for all density functions.
 - 1. Assume $X \sim N(0,1)$ and $Y \sim N(0,4)$ and that X and Y are independent.
 - (a) Compute E[(X + 1)(Y 1)].

1p.

(b) Compute E[Y|X + Y = 6].

1.5p.

(c) Derive the distribution of $U = \frac{X}{Y}$. Does it belong to a known distribution?

2.5p.

2. Let $Y|\theta \sim Bin(n,\theta)$, where n is a known positive integer. Let the density of θ be

$$f_{\theta}(\theta) = a \cdot \theta^3$$

for $\theta \in [0,1]$ and $f_{\theta}(\theta) = 0$ otherwise.

(a) Determine the constant a, so that $f_{\theta}(\theta)$ is a proper density.

1p.

(b) Calculate the variance of Y.

2p.

(c) Compute the density of Y.

2p.

- 3. Consider a fair die with probability 1/6 of rolling a six. Consider a game where the die is rolled until a six comes up and let the random variable X denote the number of die rolls required until this event happens.
 - (a) Determine the probability function of X and state if it belongs to a known distribution.
 - (b) Assume that the same game is played n times and let X_{max} be the maximum value of X across all games. Derive the distribution function of X_{max} .
 - (c) How many times does one have to play the game for having at least 50% chance of obtaining a value of X_{max} that is greater than or equal to 30? 2p.
- 4. Let X and Y be random variables such that

$$Y|X = x \sim N(0, x)$$

with $X \sim Po(\lambda)$.

(a) Find the characteristic function of Y.

(b) Show that

$$\frac{Y}{\sqrt{\lambda}} \stackrel{d}{\to} N(0,1)$$

as $\lambda \to \infty$.

(c) Formulate and prove the Central Limit Theorm.

GOOD LUCK!

Per