

PROBABILITY THEORY

LECTURE 5

Per Siden

**Division of Statistics
Dept. of Computer and Information Science
Linköping University**

OVERVIEW LECTURE 5

- ▶ Linear algebra recap
- ▶ Multivariate normal distribution

LINEAR ALGEBRA RECAP

- **Eigen-decomposition** of an $n \times n$ symmetric matrix **A**

$$\mathbf{C}'\mathbf{A}\mathbf{C} = \mathbf{D}$$

where $\mathbf{D} = \text{Diag}(\lambda_1, \dots, \lambda_n)$ and \mathbf{C} is an orthogonal matrix.

LINEAR ALGEBRA RECAP

- ▶ **Eigen-decomposition** of an $n \times n$ symmetric matrix **A**

$$\mathbf{C}'\mathbf{A}\mathbf{C} = \mathbf{D}$$

where $\mathbf{D} = \text{Diag}(\lambda_1, \dots, \lambda_n)$ and \mathbf{C} is an orthogonal matrix.

- ▶ **Orthogonal matrix:**

- ▶ $\mathbf{C}'\mathbf{C} = \mathbf{I}$
- ▶ $\mathbf{C}^{-1} = \mathbf{C}'$
- ▶ $\det \mathbf{C} = \pm 1$

LINEAR ALGEBRA RECAP

- ▶ **Eigen-decomposition** of an $n \times n$ symmetric matrix **A**

$$\mathbf{C}'\mathbf{A}\mathbf{C} = \mathbf{D}$$

where $\mathbf{D} = \text{Diag}(\lambda_1, \dots, \lambda_n)$ and \mathbf{C} is an orthogonal matrix.

- ▶ **Orthogonal matrix:**

- ▶ $\mathbf{C}'\mathbf{C} = \mathbf{I}$
- ▶ $\mathbf{C}^{-1} = \mathbf{C}'$
- ▶ $\det \mathbf{C} = \pm 1$

- ▶ The columns of $\mathbf{C} = (\mathbf{c}_1, \dots, \mathbf{c}_n)$ are the eigenvectors, and λ_i is the i th largest eigenvalue.
- ▶ $\det \mathbf{A} = \lambda_1 \cdot \lambda_2 \cdots \lambda_n$.

QUADRATIC FORMS AND POSITIVE-DEFINITENESS

- ▶ Quadratic form

$$Q(\mathbf{x}) = \mathbf{x}'\mathbf{A}\mathbf{x}$$

- ▶ $Q(\mathbf{x})$ is **positive-definite** if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- ▶ $Q(\mathbf{x})$ is **positive-semidefinite** if $Q(\mathbf{x}) \geq 0$ for all $\mathbf{x} \neq \mathbf{0}$.

QUADRATIC FORMS AND POSITIVE-DEFINITENESS

► Quadratic form

$$Q(\mathbf{x}) = \mathbf{x}'\mathbf{A}\mathbf{x}$$

- $Q(\mathbf{x})$ is **positive-definite** if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- $Q(\mathbf{x})$ is **positive-semidefinite** if $Q(\mathbf{x}) \geq 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- $Q(\mathbf{x})$ is **positive-definite** iff all eigenvalues of \mathbf{A} are positive.
- $Q(\mathbf{x})$ is **positive-semidefinite** iff all eigenvalues of \mathbf{A} are non-negative.

MATRIX SQUARE ROOT

- ▶ If $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ is diagonal, then $\tilde{D} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$ is the square root of D :

$$\tilde{D}\tilde{D} = D$$

and we can write $D^{1/2} = \tilde{D}$.

MATRIX SQUARE ROOT

- ▶ If $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ is diagonal, then $\tilde{D} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$ is the square root of D :

$$\tilde{D}\tilde{D} = D$$

and we can write $D^{1/2} = \tilde{D}$.

- ▶ The **square root of a positive definite matrix** A

$$A = CDC'$$

can be defined as

$$A^{1/2} = C\tilde{D}C'$$

where $\tilde{D} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$.

MATRIX SQUARE ROOT

- ▶ If $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ is diagonal, then $\tilde{D} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$ is the square root of D :

$$\tilde{D}\tilde{D} = D$$

and we can write $D^{1/2} = \tilde{D}$.

- ▶ The **square root of a positive definite matrix** A

$$A = CDC'$$

can be defined as

$$A^{1/2} = C\tilde{D}C'$$

where $\tilde{D} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_n})$.

- ▶ Check:

$$A^{1/2}A^{1/2} = C\tilde{D}C'C\tilde{D}C' = C\tilde{D}\tilde{D}C' = CDC' = A$$

- ▶ We also have

$$(A^{-1})^{1/2} = (A^{1/2})^{-1}$$

which is denoted by $A^{-1/2}$.

COVARIANCE MATRIX

- Mean vector

$$\mu = E\mathbf{X} = \begin{pmatrix} EX_1 \\ \vdots \\ EX_n \end{pmatrix}$$

COVARIANCE MATRIX

- ▶ Mean vector

$$\mu = E\mathbf{X} = \begin{pmatrix} EX_1 \\ \vdots \\ EX_n \end{pmatrix}$$

- ▶ Covariance matrix

$$\Lambda = \text{Cov}(\mathbf{X}) = E(\mathbf{X} - \mu)(\mathbf{X} - \mu)'$$

COVARIANCE MATRIX

- ▶ Mean vector

$$\mu = E\mathbf{X} = \begin{pmatrix} EX_1 \\ \vdots \\ EX_n \end{pmatrix}$$

- ▶ Covariance matrix

$$\Lambda = \text{Cov}(\mathbf{X}) = E(\mathbf{X} - \mu)(\mathbf{X} - \mu)'$$

TH Every covariance matrix is positive semidefinite.

- ▶ $\det \Lambda \geq 0$.

LINEAR TRANSFORMATIONS

- Recall that if $Y = aX + b$, where $E(X) = \mu$ and $Var(X) = \sigma^2$ then

$$E(Y) = a\mu + b$$

$$Var(Y) = a^2\sigma^2$$

LINEAR TRANSFORMATIONS

- Recall that if $Y = aX + b$, where $E(X) = \mu$ and $Var(X) = \sigma^2$ then

$$E(Y) = a\mu + b$$

$$Var(Y) = a^2\sigma^2$$

TH Multivariate linear transformation

Let $\mathbf{Y} = \mathbf{B}\mathbf{X} + \mathbf{b}$, where \mathbf{X} is $n \times 1$ and \mathbf{B} is $m \times n$.

Assume $E\mathbf{X} = \mu$ and $Cov(\mathbf{X}) = \Lambda$. Then,

$$E(\mathbf{Y}) = \mathbf{B}\mu + \mathbf{b}$$

$$Cov(\mathbf{Y}) = \mathbf{B}\Lambda\mathbf{B}'$$

LINEAR TRANSFORMATIONS

- Recall that if $Y = aX + b$, where $E(X) = \mu$ and $Var(X) = \sigma^2$ then

$$E(Y) = a\mu + b$$

$$Var(Y) = a^2\sigma^2$$

TH Multivariate linear transformation

Let $\mathbf{Y} = \mathbf{B}\mathbf{X} + \mathbf{b}$, where \mathbf{X} is $n \times 1$ and \mathbf{B} is $m \times n$.

Assume $E\mathbf{X} = \mu$ and $Cov(\mathbf{X}) = \Lambda$. Then,

$$E(\mathbf{Y}) = \mathbf{B}\mu + \mathbf{b}$$

$$Cov(\mathbf{Y}) = \mathbf{B}\Lambda\mathbf{B}'$$

TH Let $\mathbf{X} = (X_1, \dots, X_n)'$ where $X_1, \dots, X_n \stackrel{iid}{\sim} N(0, 1)$. Then

$$\mathbf{Y} = \mu + \Lambda^{1/2}\mathbf{X} \sim N(\mu, \Lambda)$$

MULTIVARIATE NORMAL DISTRIBUTION

- ▶ Multivariate normal $\mathbf{X} \sim N(\mu, \Lambda)$, where \mathbf{X} is a $n \times 1$ random vector.

MULTIVARIATE NORMAL DISTRIBUTION

- ▶ Multivariate normal $\mathbf{X} \sim N(\mu, \Lambda)$, where \mathbf{X} is a $n \times 1$ random vector.
- ▶ Three equivalent definitions:
 - ▶ \mathbf{X} is (multivariate) normal iff $\mathbf{a}'\mathbf{X}$ is (univariate) normal for all \mathbf{a} .

MULTIVARIATE NORMAL DISTRIBUTION

- ▶ Multivariate normal $\mathbf{X} \sim N(\mu, \Lambda)$, where \mathbf{X} is a $n \times 1$ random vector.
- ▶ Three equivalent definitions:
 - ▶ \mathbf{X} is (multivariate) normal iff $\mathbf{a}'\mathbf{X}$ is (univariate) normal for all \mathbf{a} .
 - ▶ \mathbf{X} is multivariate normal iff its characteristic function is

$$\varphi_{\mathbf{X}}(\mathbf{t}) = Ee^{i\mathbf{t}'\mathbf{X}} = \exp\left(i\mathbf{t}'\mu - \frac{1}{2}\mathbf{t}'\Lambda\mathbf{t}\right)$$

MULTIVARIATE NORMAL DISTRIBUTION

- ▶ Multivariate normal $\mathbf{X} \sim N(\mu, \Lambda)$, where \mathbf{X} is a $n \times 1$ random vector.
- ▶ Three equivalent definitions:
 - ▶ \mathbf{X} is (multivariate) normal iff $\mathbf{a}'\mathbf{X}$ is (univariate) normal for all \mathbf{a} .
 - ▶ \mathbf{X} is multivariate normal iff its characteristic function is

$$\varphi_{\mathbf{X}}(\mathbf{t}) = Ee^{i\mathbf{t}'\mathbf{X}} = \exp\left(i\mathbf{t}'\mu - \frac{1}{2}\mathbf{t}'\Lambda\mathbf{t}\right)$$

- ▶ \mathbf{X} is multivariate normal iff its density function is of the form

$$f_{\mathbf{X}}(\mathbf{x}) = \left(\frac{1}{2\pi}\right)^{n/2} \frac{1}{\sqrt{\det \Lambda}} \exp\left\{-\frac{1}{2}(\mathbf{x} - \mu)' \Lambda^{-1}(\mathbf{x} - \mu)\right\}$$

MULTIVARIATE NORMAL DISTRIBUTION

- ▶ Multivariate normal $\mathbf{X} \sim N(\mu, \Lambda)$, where \mathbf{X} is a $n \times 1$ random vector.
- ▶ Three equivalent definitions:
 - ▶ \mathbf{X} is (multivariate) normal iff $\mathbf{a}'\mathbf{X}$ is (univariate) normal for all \mathbf{a} .
 - ▶ \mathbf{X} is multivariate normal iff its characteristic function is

$$\varphi_{\mathbf{X}}(\mathbf{t}) = Ee^{i\mathbf{t}'\mathbf{X}} = \exp\left(i\mathbf{t}'\mu - \frac{1}{2}\mathbf{t}'\Lambda\mathbf{t}\right)$$

- ▶ \mathbf{X} is multivariate normal iff its density function is of the form

$$f_{\mathbf{X}}(\mathbf{x}) = \left(\frac{1}{2\pi}\right)^{n/2} \frac{1}{\sqrt{\det \Lambda}} \exp\left\{-\frac{1}{2}(\mathbf{x} - \mu)' \Lambda^{-1}(\mathbf{x} - \mu)\right\}$$

- ▶ **Bivariate normal** ($n = 2$)

$$\Lambda = \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix}$$

where $-1 \leq \rho \leq 1$ is the correlation coefficient.

PROPERTIES OF THE NORMAL DISTRIBUTION

► Let $\mathbf{X} \sim N(\mu, \Lambda)$.

TH Linear combinations: $\mathbf{Y} = \mathbf{B}\mathbf{X} + \mathbf{b}$, where \mathbf{X} is $n \times 1$ and \mathbf{B} is $m \times n$. Then

$$\mathbf{Y} \sim N(\mathbf{B}\mu + \mathbf{b}, \mathbf{B}\Lambda\mathbf{B}')$$

PROPERTIES OF THE NORMAL DISTRIBUTION

► Let $\mathbf{X} \sim N(\mu, \Lambda)$.

TH Linear combinations: $\mathbf{Y} = \mathbf{B}\mathbf{X} + \mathbf{b}$, where \mathbf{X} is $n \times 1$ and \mathbf{B} is $m \times n$. Then

$$\mathbf{Y} \sim N(\mathbf{B}\mu + \mathbf{b}, \mathbf{B}\Lambda\mathbf{B}')$$

COR The components of \mathbf{X} are all normal ($\mathbf{B} = (0, \dots, 1, 0, \dots, 0)$)

$$Y_i \sim N(\mu_i, \Lambda_{ii})$$

PROPERTIES OF THE NORMAL DISTRIBUTION

► Let $\mathbf{X} \sim N(\mu, \Lambda)$.

TH Linear combinations: $\mathbf{Y} = \mathbf{B}\mathbf{X} + \mathbf{b}$, where \mathbf{X} is $n \times 1$ and \mathbf{B} is $m \times n$. Then

$$\mathbf{Y} \sim N(\mathbf{B}\mu + \mathbf{b}, \mathbf{B}\Lambda\mathbf{B}')$$

COR The components of \mathbf{X} are all normal ($\mathbf{B} = (0, \dots, 1, 0, \dots, 0)$)

$$Y_i \sim N(\mu_i, \Lambda_{ii})$$

COR Let $\mathbf{X} = \begin{pmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \end{pmatrix}$ where \mathbf{X}_1 is $n_1 \times 1$ and \mathbf{X}_2 is $n_2 \times 1$ ($n_1 + n_2 = n$).
Then

$$\mathbf{X}_1 \sim N(\mu_1, \Lambda_1)$$

where μ_1 are the n_1 first elements of μ and Λ_1 is the $n_1 \times n_1$ submatrix of Λ .


MARGINAL NORMAL MAY NOT BE JOINTLY NORMAL

- ▶ We know that $\mathbf{X} \sim N(\mu, \Lambda)$ implies that all marginals are normal.


MARGINAL NORMAL MAY NOT BE JOINTLY NORMAL

- ▶ We know that $\mathbf{X} \sim N(\mu, \Lambda)$ implies that all marginals are normal.
- ▶ The converse does not hold. Normal marginals does not imply that the joint distribution is normal.

Marginals are normal, joint is normal



Marginals are normal, joint is not normal



CONDITIONAL DISTRIBUTIONS FROM $N(\mu, \Lambda)$

- ▶ Let $\begin{pmatrix} X \\ Y \end{pmatrix} \sim N_2(\mu, \Lambda)$, where

$$\mu = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix} \quad \text{and} \quad \Lambda = \begin{pmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix}$$

- ▶ Then

$$Y|X = x \sim N \left[\mu_y + \rho \frac{\sigma_y}{\sigma_x} (x - \mu_x), \sigma_y^2 (1 - \rho^2) \right]$$

CONDITIONAL DISTRIBUTIONS FROM $N(\mu, \Lambda)$

- ▶ Let $\begin{pmatrix} X \\ Y \end{pmatrix} \sim N_2(\mu, \Lambda)$, where

$$\mu = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix} \quad \text{and} \quad \Lambda = \begin{pmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix}$$

- ▶ Then

$$Y|X = x \sim N \left[\mu_y + \rho \frac{\sigma_y}{\sigma_x} (x - \mu_x), \sigma_y^2 (1 - \rho^2) \right]$$

- ▶ The **regression function** $E(Y|X)$ is **linear** and $\text{Var}(Y|X)$ = residual variance.

CONDITIONAL DISTRIBUTIONS FROM $N(\mu, \Lambda)$

- Let $\begin{pmatrix} X \\ Y \end{pmatrix} \sim N_2(\mu, \Lambda)$, where

$$\mu = \begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix} \quad \text{and} \quad \Lambda = \begin{pmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{pmatrix}$$

- Then

$$Y|X = x \sim N \left[\mu_y + \rho \frac{\sigma_y}{\sigma_x} (x - \mu_x), \sigma_y^2 (1 - \rho^2) \right]$$

- The **regression function** $E(Y|X)$ is **linear** and $\text{Var}(Y|X)$ = residual variance.

TH Let $\mathbf{X} = \begin{pmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \end{pmatrix}$ and partition μ and Λ accordingly as

$$\mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix} \quad \text{and} \quad \Lambda = \begin{pmatrix} \Lambda_{11} & \Lambda_{12} \\ \Lambda_{21} & \Lambda_{22} \end{pmatrix}. \quad \text{Then}$$

$$\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2 \sim N \left[\mu_1 + \Lambda_{12} \Lambda_{22}^{-1} (\mathbf{x}_2 - \mu_2), \Lambda_{11} - \Lambda_{12} \Lambda_{22}^{-1} \Lambda_{21} \right]$$

INDEPENDENCE AND NORMALITY

- ▶ Correlation measures **linear** association (dependence).

INDEPENDENCE AND NORMALITY

- ▶ Correlation measures **linear** association (dependence).
- ▶ In general: Uncorrelated \nrightarrow Independence.

INDEPENDENCE AND NORMALITY

- ▶ Correlation measures **linear** association (dependence).
- ▶ In general: Uncorrelated \nrightarrow Independence.
- ▶ In the normal distribution: Uncorrelated \leftrightarrow Independence.

INDEPENDENCE AND NORMALITY

- ▶ Correlation measures **linear** association (dependence).
- ▶ In general: Uncorrelated \nrightarrow Independence.
- ▶ In the normal distribution: Uncorrelated \leftrightarrow Independence.
- ▶ Remember that: X and Y are jointly normal \rightarrow the regression function is linear \rightarrow the linear predictor is optimal.

INDEPENDENCE AND NORMALITY

- ▶ Correlation measures **linear** association (dependence).
- ▶ In general: Uncorrelated \nrightarrow Independence.
- ▶ In the normal distribution: Uncorrelated \leftrightarrow Independence.
- ▶ Remember that: X and Y are jointly normal \rightarrow the regression function is linear \rightarrow the linear predictor is optimal.
- ▶ $X_1, \dots, X_n \stackrel{iid}{\sim} N(\mu, \sigma^2)$, then $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ and $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$ are independent.

PRINCIPAL COMPONENTS

- ▶ Let $\mathbf{C}\mathbf{A}\mathbf{C}' = \mathbf{D} = \text{diag}(\lambda_1, \dots, \lambda_n)$.

PRINCIPAL COMPONENTS

► Let $\mathbf{C}\mathbf{\Lambda}\mathbf{C}' = \mathbf{D} = \text{diag}(\lambda_1, \dots, \lambda_n)$.

TH Let $\mathbf{X} \sim N(\boldsymbol{\mu}, \boldsymbol{\Lambda})$ and set $\mathbf{Y} = \mathbf{C}'\mathbf{X}$, then

$$\mathbf{Y} \sim N(\mathbf{C}'\boldsymbol{\mu}, \mathbf{D})$$

so that the components of \mathbf{Y} are independent and $\text{Var}(Y_i) = \lambda_i$.

