

MATH2111 Tutorial 1

QUAN Xueyang (xquan@connect.ust.hk)

T1A: Tue (18:00-18:50) 2465

T1B: Thu (18:00-18:50) 6591

1 Linear Systems

1. A **linear equation** with variables x_1, x_2, \dots, x_n is of the form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b$$

where a_i 's are coefficients, and b is a constant.

Example 1.1. Determine if the following equations are linear equations.

- (a) $x_1 = x_2$
 - (b) $x_1^2 = x_2^2$
 - (c) $\sqrt{3}x_1 - x_2 = 1$
 - (d) $3\sqrt{x_1} - x_2 = 1$
 - (e) $x_1x_2 + x_1 = e^5$
2. A **linear system** is a collection of linear equations.
 3. A **solution** to a linear system is an assignment of values to variables that make all equations in the system simultaneously true.
 4. **Consistent / Inconsistent**
Any linear system has
 - (a) A unique solution, or
 - (b) Infinitely many solutions, or
 - (c) No solutions
 5. A **matrix** is a rectangular array of numbers. A matrix with m rows and n columns is said to be " $m \times n$ " or " m by n ".
 6. **Coefficient Matrix and Augmented Matrix**
For a system of linear equations,

- (a) the **coefficient matrix** is the matrix consist of all the coefficients of the linear equations;
- (b) the **augmented matrix** is the matrix consist of all the coefficients and the constants of the linear equations.

Example 1.2. Write down the coefficient matrix and the augmented matrix of the following linear system.

$$\begin{cases} x_1 + x_5 = 1 \\ 3x_2 - x_4 = 4 \\ x_1 + 2x_3 = -3 \end{cases}$$

Example 1.3. Write down the linear system with the following augmented matrix.

$$\left[\begin{array}{cccc|c} 2 & 1 & 2 & 3 & 5 \\ 0 & -2 & 0 & 4 & 1 \end{array} \right]$$

2 Row Reduction and Echelon Forms

1. Here are the 3 **elementary row operations (EROs)** on any matrices which are helpful in solving linear equations:

- Row Replacement: $cr_j + r_i$
- Row Interchange: $r_i \leftrightarrow r_j$
- Row Scaling: $cr_i, c \neq 0$

Two matrices are **row equivalent** (denoted by \sim) if one can be transformed to the other by applying a finite sequence of these row operations.

2. Linear systems with row equivalent augmented matrices have the same solutions (are **equivalent**).
3. Any matrix is in **row echelon form (REF)** if
 - (a) The rows with all zero entries must be at the bottom.
 - (b) The leading entry (the first non-zero entry on each row) must move to the right by at least one column when going down a row.

- (c) All entries in a column below a leading entry must be zeros.
4. A **reduced row echelon form (RREF)** of any matrix is a REF with the extra properties:
- (d) On a non-zero row, the leading entry must be 1.
 - (e) On the columns containing the leading entry 1, the 1 is the only non-zero entry.
5. **Algorithm to get REF from a random matrix:**
- (1) Stop when all entries are zeros, or no visible entry.
 - (2) Locate the left-most non-zero column, select a non-zero entry, use row interchange to move it to the top row.
 - (3) Use row replacement to make all entries below 0.
 - (4) Neglect the top row, and repeat Step 1-3 for the submatrix below that row.
6. **Algorithm from a REF to a RREF:**
- (5) Use row scaling to scale all the leading entries to 1.
 - (6) Working from the rightmost leading entries to left, use row replacement to make all entries above each of them 0.
7. Each matrix A is row equivalent to exactly one matrix in reduced echelon form.
8. A pivot position in a matrix M is a location in M that corresponds to a leading 1 in the reduced echelon form of M . A pivot column is a column of M that contains a pivot position.
9. **Existence and Uniqueness Theorem:**
- (1) A linear system is consistent if and only if an echelon form of its augmented matrix has no row of the form $[0 \cdots 0 | \blacksquare]$ with $\blacksquare \neq 0$.
 - (2) If a linear system is consistent, then:
 - it has a unique solution if there are no free variables;
 - it has infinitely many solutions if there are free variables

3 Exercises

1. Write down the linear system with this augmented matrix: $\left[\begin{array}{cc|c} 1 & 2 & 3 \end{array} \right]$

2. Determine the value(s) of h such that the matrix

$$\left[\begin{array}{cc|c} 1 & h & -5 \\ 2 & -8 & 6 \end{array} \right]$$

is the augmented matrix of a consistent linear system.

3. Determine which matrices are in reduced echelon form and which others are only in echelon form.

(a) $\left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right]$ (b) $\left[\begin{array}{cccc} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$ (c) $\left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$ (d) $\left[\begin{array}{ccccc} 1 & 1 & 0 & 1 & 1 \\ 0 & 2 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 0 & 4 \end{array} \right]$

4. Describe the possible echelon forms of a nonzero 2×2 matrix. Use the symbols ■, * and 0, where the leading entries (■) may have any nonzero value; the starred entries (*) may have any value (including zero).

5. Transform the following matrices to Reduced Row Echelon Form.

$$\left[\begin{array}{cccccc} 1 & 2 & 1 & -1 & 2.5 & 0.5 \\ 3 & 6 & 2 & 2 & 0 & 4 \\ 4 & 8 & 2 & 6 & -5 & 7 \end{array} \right]$$

6. Consider the following linear systems.

$$\begin{cases} x_1 + 2x_2 + x_3 + 3x_4 = 0 \\ 2x_1 + 5x_2 + 2x_3 + 8x_4 = 1 \\ 3x_1 + 5x_2 + 4x_3 + 9x_4 = -5 \end{cases}$$

(a) Write down the augmented matrix of the linear system.

(b) Get the reduced echelon form of the augmented matrix using EROs.

(c) Solve the linear system.