

MATH 2111: Tutorial 4 Linear Independence and Introduction to Linear Transformations

T1A&T1B QUAN Xueyang
T1C&T2A SHEN Yinan
T2B&T2C ZHANG Fa

Department of Mathematics, HKUST

- Linear independence/dependence
- Linear independence of columns of matrix
- Characterization of linearly dependent sets
- Conditions for linear dependence
- Transformation

Example 1(a)

Express the general solutions of the following non-homogeneous systems in terms of the given particular solutions.

$$\begin{cases} x_1 + x_2 + 3x_3 + 4x_4 + 3x_5 = 5 \\ 2x_1 + 2x_2 + 2x_4 + 4x_5 = 4 \\ -x_1 - x_2 + x_3 - x_5 = -1 \end{cases}$$

$\begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$ is a solution of the above linear system.

1(a). For $A\vec{x}=\vec{b}$, let \vec{p} be a particular solution,

\vec{x}_h is any solution of $A\vec{x}=\vec{0}$

Then solution set of $Ax=b$ is of form: $\vec{w} = \vec{p} + \vec{x}_h$

① Solve $A\vec{x}=\vec{0}$:

$$\left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 3 & 0 \\ 2 & 2 & 0 & 2 & 4 & 0 \\ -1 & -1 & 1 & 0 & -1 & 0 \end{array} \right] \xrightarrow[R_3+R_1 \rightarrow R_3]{R_2-2R_1 \rightarrow R_2} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 3 & 0 \\ 0 & 0 & -6 & -6 & -2 & 0 \\ 0 & 0 & 4 & 4 & 2 & 0 \end{array} \right]$$

$$\xrightarrow[-\frac{1}{6}R_2 \rightarrow R_2]{\frac{1}{4}R_3 \rightarrow R_3} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 3 & 0 \\ 0 & 0 & 1 & 1 & \frac{1}{3} & 0 \\ 0 & 0 & 1 & 1 & \frac{1}{2} & 0 \end{array} \right]$$

$$\xrightarrow{R_3-R_2 \rightarrow R_3} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 3 & 0 \\ 0 & 0 & 1 & 1 & \frac{1}{3} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{6} & 0 \end{array} \right]$$

$$\xrightarrow{6R_3 \rightarrow R_3} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 3 & 0 \\ 0 & 0 & 1 & 1 & \frac{1}{3} & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

$$\xrightarrow[R_1-3R_3 \rightarrow R_1]{R_2-\frac{1}{3}R_3 \rightarrow R_2} \left[\begin{array}{ccccc|c} 1 & 1 & 3 & 4 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

$$\xrightarrow{R_1-3R_2 \rightarrow R_1} \left[\begin{array}{ccccc|c} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right] \leftarrow \text{RREF}$$

Thus $\begin{cases} x_1 + x_2 + x_4 = 0 \\ x_3 + x_4 = 0 \\ x_5 = 0 \end{cases}$

x_2, x_4 are free.

$$\therefore \vec{x}_h = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -x_2 & -x_4 \\ x_2 & \\ -x_4 & x_4 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} x_2 + \begin{bmatrix} -1 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix} x_4 \quad x_2, x_4 \in \mathbb{R}$$

$$\textcircled{2} \text{ Check } \vec{p} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} \text{ is a solution to } A\vec{x} = \vec{b}$$

Thus, solution set for $A\vec{x} = \vec{b}$ is

$$\vec{w} = \vec{p} + \vec{x}_h = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} x_2 + \begin{bmatrix} -1 \\ 0 \\ -1 \\ 1 \\ 0 \end{bmatrix} x_4 \quad x_2, x_4 \in \mathbb{R}$$

Example 1(b)

Denote the coefficient matrix as \mathbf{A} . Use as many columns of \mathbf{A} as possible to construct a matrix \mathbf{B} with the property that the equation $\mathbf{B}\mathbf{x} = 0$ has only the trivial solution. (Solve $\mathbf{B}\mathbf{x} = 0$ to verify your work.)

1(b). According to the RREF of A,
the first three columns are pivot columns

If use the corresponding columns of A to form B,
we have:

$$B = \begin{bmatrix} 1 & 3 & 3 \\ 2 & 0 & 4 \\ -1 & 1 & -1 \end{bmatrix},$$

If do row reduction on $[B|\vec{0}]$, we get $\left[\begin{array}{ccc|c} 1 & 3 & 3 & 0 \\ 2 & 0 & 4 & 0 \\ -1 & 1 & -1 & 0 \end{array} \right]$.

Since $B\vec{x} = \vec{0}$ has no free variable, it has only trivial solution.

If use any additional columns of A to form B, then $B\vec{x} = \vec{0}$
will have free variable, so $B\vec{x} = \vec{0}$ has nontrivial solution.

Example 2

Find conditions on p and q such that the set of vectors

$$\left\{ \begin{array}{c} \left[\begin{array}{c} 1 \\ 2 \\ 3 \\ 0 \end{array} \right], \left[\begin{array}{c} 2 \\ 5 \\ 5 \\ 1 \end{array} \right], \left[\begin{array}{c} 1 \\ 2 \\ 4 \\ p \end{array} \right], \left[\begin{array}{c} 3 \\ 8 \\ 9 \\ q \end{array} \right] \end{array} \right\}$$

$\begin{array}{cccc} \parallel & \parallel & \parallel & \parallel \\ \rightarrow & \rightarrow & \rightarrow & \rightarrow \\ a_1 & a_2 & a_3 & a_4 \end{array}$

is linearly independent.

2. Denote

$$A = \begin{bmatrix} 1 & 2 & 1 & 3 \\ 2 & 5 & 2 & 8 \\ 3 & 5 & 4 & 9 \\ 0 & 1 & p & q \end{bmatrix},$$

$$x_1 \vec{a}_1 + x_2 \vec{a}_2 + x_3 \vec{a}_3 + x_4 \vec{a}_4 = \vec{0}$$

$$\Leftrightarrow$$

$$A\vec{x} = \vec{0}$$

By thm, we need to guarantee $A\vec{x} = \vec{0}$ has only trivial solution.

$$\left[\begin{array}{cccc|c} 1 & 2 & 1 & 3 & 0 \\ 2 & 5 & 2 & 8 & 0 \\ 3 & 5 & 4 & 9 & 0 \\ 0 & 1 & p & q & 0 \end{array} \right]$$

$$\begin{array}{l} R_2 - 2R_1 \rightarrow R_2 \\ R_3 - 3R_1 \rightarrow R_3 \end{array} \rightarrow \left[\begin{array}{cccc|c} 1 & 2 & 1 & 3 & 0 \\ 0 & 1 & 0 & 2 & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 1 & p & q & 0 \end{array} \right]$$

$$\begin{array}{l} R_3 + R_2 \rightarrow R_3 \\ R_4 - R_2 \rightarrow R_4 \end{array} \rightarrow \left[\begin{array}{cccc|c} \underline{1} & 2 & 1 & 3 & 0 \\ 0 & \underline{1} & 0 & 2 & 0 \\ 0 & 0 & \underline{1} & 2 & 0 \\ 0 & 0 & p & q-2 & 0 \end{array} \right]$$

← REF

$$\xrightarrow{R_4 - pR_3 \rightarrow R_4} \left[\begin{array}{cccc|c} \underline{1} & 2 & 1 & 3 & 0 \\ 0 & \underline{1} & 0 & 2 & 0 \\ 0 & 0 & \underline{1} & 2 & 0 \\ 0 & 0 & 0 & q-2-2p & 0 \end{array} \right]$$

The matrix already has pivots in columns 1, 2, 3, to have a pivot in column 4, we must have $q-2-2p \neq 0$.

Example 3

Consider matrix \mathbf{A} ,

$$\mathbf{A} = \begin{bmatrix} | & | & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \mathbf{a}_4 \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} 1 & 2 & 1 & 3 \\ 2 & 5 & 2 & 8 \\ 3 & 5 & 4 & 9 \end{bmatrix},$$

Find a vector which is in $\text{Span}\{\mathbf{a}_1, \mathbf{a}_2\}$ and also in $\text{Span}\{\mathbf{a}_3, \mathbf{a}_4\}$, or explain why such a vector cannot exist.

(Given $\begin{bmatrix} 3 \\ -2 \\ -2 \\ 1 \end{bmatrix}$ is a solution to $\mathbf{Ax} = \mathbf{0}$.)

$$3. \quad \vec{v} \in \text{Span}\{\vec{a}_1, \vec{a}_2\} \text{ if } \vec{v} = c_1\vec{a}_1 + c_2\vec{a}_2 \text{ for some } c_1, c_2.$$

$$\vec{v} \in \text{Span}\{\vec{a}_3, \vec{a}_4\} \text{ if } \vec{v} = c_3\vec{a}_3 + c_4\vec{a}_4 \text{ for some } c_3, c_4.$$

$$\therefore c_1\vec{a}_1 + c_2\vec{a}_2 = \vec{v} = c_3\vec{a}_3 + c_4\vec{a}_4$$

$$\therefore c_1\vec{a}_1 + c_2\vec{a}_2 - c_3\vec{a}_3 - c_4\vec{a}_4 = \vec{0}$$

$$\therefore A \begin{bmatrix} c_1 \\ c_2 \\ -c_3 \\ -c_4 \end{bmatrix} = \vec{0}$$

$$\text{Since } \begin{bmatrix} 3 \\ -2 \\ -2 \\ 1 \end{bmatrix} \text{ is a solution to } A\vec{x} = \vec{0},$$

$$\therefore \text{take } c_1 = 3, c_2 = -2, c_3 = 2, c_4 = -1$$

$$\therefore \vec{v} = 3 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} - 2 \begin{bmatrix} 2 \\ 5 \\ 5 \end{bmatrix} = \begin{bmatrix} -1 \\ -4 \\ -1 \end{bmatrix}, \text{ which is also } 2 \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix} - 1 \begin{bmatrix} 3 \\ 8 \\ 9 \end{bmatrix}$$

Example 4

State whether each of the following statement is true or false. (If it is true, give a brief justification; if it is false, give a counterexample.)

(a) If $\mathbf{A} \begin{bmatrix} 4 \\ 0 \\ 2 \\ -3 \end{bmatrix} = \mathbf{0}$, then $\mathbf{A}\mathbf{e}_4$ is a linear combination of the first three columns of \mathbf{A} .

(b) Let \mathbf{A} be a 4×3 matrix with columns \mathbf{a}_1 , \mathbf{a}_2 , \mathbf{a}_3 , and suppose \mathbf{b} is a vector in \mathbb{R}^4 such that $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{b}\}$ is linearly dependent. Then $\mathbf{A}\mathbf{x} = \mathbf{b}$ has a solution.

4(a). True.

Let \vec{a}_i be the i -th column of A .

$$A \begin{bmatrix} 4 \\ 0 \\ 2 \\ -3 \end{bmatrix} = \vec{0} \quad \text{means} \quad 4\vec{a}_1 + 2\vec{a}_3 - 3\vec{a}_4 = \vec{0},$$

$$\therefore A\vec{e}_4 = \vec{a}_4 = \frac{4}{3}\vec{a}_1 + \frac{2}{3}\vec{a}_3.$$

$$\vec{e}_i := \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \leftarrow i\text{th row.}$$

4(b). False.

$$\text{Take } \vec{a}_1 = \vec{e}_1, \quad \vec{a}_2 = \vec{e}_2, \quad \vec{a}_3 = \vec{e}_1 + \vec{e}_2, \quad \text{and } \vec{b} = \vec{e}_3$$

Easy to check these 4 vectors are linearly dependent
($\vec{a}_3 = \vec{a}_1 + \vec{a}_2$).

However, \vec{b} can't be expressed as a linear combination of \vec{a}_1, \vec{a}_2 and \vec{a}_3 .

Remark: linear dependence only implies some vector can be expressed as a linear combination of rest vectors. ✓ NOT necessarily all

Example 5

Consider

$$F \left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \right) = \begin{bmatrix} x_1 + x_3 \\ 0 \\ 0 \\ 3x_1 - x_2 \end{bmatrix}$$

(a) What is the domain of F ?

(b) Find the image of $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ under F .

5(a). domain: \mathbb{R}^3

$$(b). F\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1+0 \\ 0 \\ 0 \\ 3 \times 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 3 \end{bmatrix}$$