

Equivalence	Name
$p \wedge T \equiv p$ $p \vee F \equiv p$	Identity laws
$p \vee T \equiv T$ $p \wedge F \equiv F$	Domination laws
$p \vee p \equiv p$ $p \wedge p \equiv p$	Idempotent laws
$\neg(\neg p) \equiv p$	Double negation law
$p \vee q \equiv q \vee p$ $p \wedge q \equiv q \wedge p$	Commutative laws

Equivalence	Name
$(p \vee q) \vee r \equiv p \vee (q \vee r)$ $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$	Associative laws
$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$ $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$	Distributive laws
$\neg(p \wedge q) \equiv \neg p \vee \neg q$ $\neg(p \vee q) \equiv \neg p \wedge \neg q$	De Morgan's laws
$p \vee (p \wedge q) \equiv p$ $p \wedge (p \vee q) \equiv p$	Absorption laws
$p \vee \neg p \equiv T$ $p \wedge \neg p \equiv F$	Negation laws

Equivalence	Name
$p \rightarrow q \equiv \neg p \vee q$ $p \rightarrow q \equiv \neg q \rightarrow \neg p$ $(p \rightarrow q) \wedge (p \rightarrow r) \equiv p \rightarrow (q \wedge r)$ $(p \rightarrow q) \vee (p \rightarrow r) \equiv p \rightarrow (q \vee r)$ $(p \rightarrow r) \wedge (q \rightarrow r) \equiv (p \vee q) \rightarrow r$ $(p \rightarrow r) \vee (q \rightarrow r) \equiv (p \wedge q) \rightarrow r$	Involving conditional statements
$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$ $p \leftrightarrow q \equiv \neg p \leftrightarrow \neg q$ $p \leftrightarrow q \equiv (p \wedge q) \vee (\neg p \wedge \neg q)$	Involving biconditional statements

Quantifications of Two Variables		
Statement	When is it true?	When is it false?
$\forall x \forall y P(x, y)$	$P(x, y)$ is true for every pair x, y .	There is a pair x, y for which $P(x, y)$ is false.
$\forall y \forall x P(x, y)$		
$\forall x \exists y P(x, y)$	For every x there is a y for which $P(x, y)$ is true.	There is an x such that $P(x, y)$ is false for every y .
$\exists x \forall y P(x, y)$	There is an x such that $P(x, y)$ is true for every y .	For every x there is a y for which $P(x, y)$ is false.
$\exists x \exists y P(x, y)$	There is a pair x, y for which $P(x, y)$ is true.	$P(x, y)$ is false for every pair x, y .
$\forall y \exists x P(x, y)$		

Rules of Inference for Propositional Logic		
Rule of inference	Tautology	Name
$\frac{p}{p \rightarrow q}$ $\therefore q$	$[p \wedge (p \rightarrow q)] \rightarrow q$	Modus ponens
$\frac{\neg q}{p \rightarrow q}$ $\therefore \neg p$	$[\neg q \wedge (p \rightarrow q)] \rightarrow \neg p$	Modus tollens
$\frac{p \rightarrow q}{q \rightarrow r}$ $\therefore p \rightarrow r$	$[(p \rightarrow q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$	Hypothetical syllogism
$\frac{p \vee q}{\neg p}$ $\therefore q$	$[(p \vee q) \wedge \neg p] \rightarrow q$	Disjunctive syllogism

Rules of Inference for Propositional Logic		
Rule of inference	Tautology	Name
$\frac{p}{p \vee q}$ $\therefore p \vee q$	$p \rightarrow (p \vee q)$	Addition
$\frac{p \wedge q}{p}$ $\therefore p$	$(p \wedge q) \rightarrow p$	Simplification
$\frac{p}{q}$ $\therefore p \wedge q$	$[(p) \wedge (q)] \rightarrow (p \wedge q)$	Conjunction
$\frac{p \vee q}{\neg p \vee r}$ $\therefore q \vee r$	$[(p \vee q) \wedge (\neg p \vee r)] \rightarrow (q \vee r)$	Resolution

Rules of Inference for Predicate Logic	
Rule of inference	Name
$\frac{\forall x P(x)}{\therefore P(c)}$	Universal instantiation
$\frac{P(c) \text{ for an arbitrary } c}{\therefore \forall x P(x)}$	Universal generalization
$\frac{\exists x P(x)}{\therefore P(c) \text{ for some element } c}$	Existential instantiation
$\frac{P(c) \text{ for some element } c}{\therefore \exists x P(x)}$	Existential generalization