HKDSE MATH EP

M2

HONG KONG EDUCATIONAL PUBLISHING COMPANY

HONG KONG DIPLOMA OF SECONDARY EDUCATION EXAMINATION

MATHEMATICS Extended Part

Module 2 (Algebra and Calculus) MOCK EXAM 7 Question-Answer Book

Time allowed: 2½ hours

This paper must be answered in English

INSTRUCTIONS

- 1. After the announcement of the start of the examination, you should first write your Candidate Number in the space provided on Page 1 and stick barcode labels in the spaces provided on Pages 1, 3, 5, 7, 9 and 11.
- 2. This paper consists of TWO sections, A and B.
- 3. Attempt **ALL** questions in this paper. Write your answers in the spaces provided in this Question-Answer Book. Do not write in the margins. Answers written in the margins will not be marked.
- 4. Graph paper and supplementary answer sheets will be supplied on request. Write your Candidate Number, mark the question number box and stick a barcode label on each sheet, and fasten them with string **INSIDE** this book.
- 5. Unless otherwise specified, all working must be clearly shown.
- 6. Unless otherwise specified, numerical answers must be exact.
- 7. In this paper, vectors may be represented by bold-type letters such as \mathbf{u} , but candidates are expected to use appropriate symbols such as \mathbf{u} in their working.
- 8. The diagrams in this paper are not necessarily drawn to scale.
- 9. No extra time will be given to candidates for sticking on the barcode labels or filling in the question number boxes after the 'Time is up' announcement.
- © 香港教育圖書公司 保留版權 Hong Kong Educational Publishing Company All Rights Reserved 2016

Please	stick t	the I	barc	ode	la l	bel	he	re.
Candidate Number			Ţ					

$$\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B)$$

$$2\cos A\cos B = \cos (A+B) + \cos (A-B)$$

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B)$$

 $\sin A + \sin B = 2\sin\frac{A+B}{2}\cos\frac{A-B}{2}$

$$\sin A - \sin B = 2\cos\frac{A+B}{2}\sin\frac{A-B}{2}$$

$$\cos A + \cos B = 2\cos\frac{A+B}{2}\cos\frac{A-B}{2}$$

$$\cos A - \cos B = -2\sin\frac{A+B}{2}\sin\frac{A-B}{2}$$

SECTION A (50 marks)

1. It is given that $(1 + kx)^n = 1 - 24x + 252x^2 + \text{terms involving higher powers of } x$, where n is a positive integer and k is a constant.

- (a) Find the values of k and n.
- (b) Find the coefficient of x^3 .

(6 marks)

Answers written in the margins will not be marked.

Please	stick	the	barco	de I	abel	here.
					عادات	
						ti bili.

		κ=	=1				(5 marl
				_			
				<u> </u>			
 			· · · · ·				
							
							•
					,		
	N-100-						
÷						-	

EP(M2) MOCK 7-3

Answers written in the margins will not be marked.

- 147 -

Go on to the next page

EP(M2) MOCK 7-4

Please stick the barcode label here.

- 4. It is given that $A B = \frac{\pi}{4}$.
 - (a) Show that $\tan A = \frac{1 + \tan B}{1 \tan B}$.
 - (b) Using the result of (a), find the value of $\tan \frac{3\pi}{8}$.

(5 marks)

	•
-	

Answers written in the margins will not be marked.

EP(M2) MOCK 7-5

Answers written in the margins will not be marked.

- 149 -

Go on to the next page

- **2012** 5. ((a) Using integration by parts, find $\int (\ln x)^2 dx$.
 - (b) In Figure 1, the shaded region is bounded by the curve $y = e^x$ for $0 \le x \le 2$ and the x-axis. Find the volume of the solid of revolution when the shaded region is revolved about the y-axis.




Figure 1

(6 marks)

Answers written in the margins will not be marked.

Please stick the barcode label here.

marked.
þe
not
will
margins
the
ij.
written in
Answers w

6.	The slope at any point (x, y) of a curve is given by	$\frac{dy}{dx} =$	$\frac{1}{x} + 1$. It	is given	that	the li	ne
	2x - y - 3 = 0 touches the curve at the point P.		-				

- (a) Find the coordinates of P.
- (b) Find the equation of the curve.
- (c) Find the equation of normal to the curve at the point P.

(6 marks)

Answers written in the margins will not be marked.

 	
 	 <u>_</u>

(a) Find the extreme value of f	•		
(b) Using the result of (a), show	w that $e^x \ge x^e$ for $x > 0$.		(6 marks)
		· · · · · · · · · · · · · · · · · · ·	
<u> </u>			
			
	· · · · · · · · · · · · · · · · · · ·		

Please stick the barcode label here.

2012 8.

Answers written in the margins will not be marked.




Figure 2

Figure 2 shows a tetrahedron \overrightarrow{OABC} . Let $\overrightarrow{OA} = 5\mathbf{i} + \mathbf{j} - 2\mathbf{k}$, $\overrightarrow{OB} = 3\mathbf{i} + \mathbf{j}$ and $\overrightarrow{OC} = 4\mathbf{i} - \mathbf{j} + 3\mathbf{k}$.

- (a) Find the area of $\triangle OAB$.
- (b) Find the distance between point C and the plane OAB.

(5 marks)

 	*	
•		

Answers written in the margins will not be marked.

EP(M2) MOCK 7-9

- 153 -

- 9. Let $A = \begin{pmatrix} 7 & 3 & 1 \\ k & 1 & -4 \\ -5 & -4 & 3 \end{pmatrix}$, where k is a positive constant.
 - (a) Show that the inverse of A is $\frac{1}{13k+26} \begin{pmatrix} 13 & 13 & 13 \\ 3k-20 & -26 & -28-k \\ 4k-5 & -13 & 3k-7 \end{pmatrix}.$
 - (b) Using the result of (a), solve the system of linear equations $\begin{cases} x + y + z = 5 \\ x 2y 3z = -3 \\ 3x y + 2z = 0 \end{cases}$

(7 marks)

(

Answers written in the margins will not be marked.

	 			-
· ·				
			<u> </u>	
	 			

·	
	·
	•
•	

Please stick the barcode label here.

(

SECTION B (50 marks)

10.




Figure 3

In Figure 3, OACB is a parallelogram, where O is the origin. D is the point of intersection of the diagonals AB and OC. CO is produced to E such that CD: DE = 1: 2. Suppose AB is a diameter of the circle passing through A, C and B, EB is the tangent to the circle at B. It is given that $\overrightarrow{OB} = m\mathbf{i} + n\mathbf{j}$ and $\overrightarrow{OC} = 6\mathbf{i} + 8\mathbf{j}$, where m > 0 and n < 0.

(a) Find OE.
(b) Find the values of m and n.
(c) Is O the incentre of ΔABE? Explain your answer.
(4 marks)

	·
· .	
	<u> </u>
	·
•	
	·

EP(M2) MOCK 7-13

- 157 - ·

Go on to the next page

11. (a) Show that
$$\begin{vmatrix} p & q & 1 \\ p^2 & q^2 & 1 \\ p^3 & q^3 & 1 \end{vmatrix} = pq(1-p)(1-q)(q-p)$$
.

(3 marks)

(b) Consider the system of linear equations in x, y, z:

(E):
$$\begin{cases} hx + 3y + z = a \\ h^2x + 9y + z = b \end{cases}$$
, where a, b, c and h are real numbers.
$$h^3x + 27y + z = c$$

- (i) Find the values of h such that (E) does not have unique solution.
- (ii) Assume h = 3 and (E) is consistent. Show that c 4b + 3a = 0 when (E) has infinitely many solutions.

(5 marks)

(c) Solve the system of equations $\begin{cases} 3x + 3y + z = 2 \\ 9x + 9y + z = 1 \\ 27x + 27y + z = -2 \\ 3x + 6y - 2z = 0 \end{cases}$

(3 marks)

Answers written in the margins will not be marked.

<u> </u>	
	

EP(M2) MOCK 7-15

. – 159 –

Go on to the next page

· .	


(a) (i) Express y in terms of x.

(ii) Find the minimum length of the road.

(6 marks)

Answers written in the margins will not be marked.

(b)



The road is built with the minimum length. In Figure 5, Sam is driving on the road from X to B at a constant speed of 50 km/h. Let $\angle YXC = \alpha$ and $\angle YCX = \beta$, where Y is the position of Sam's car.

(i) By finding $\sin \alpha$ and $\cos \alpha$, show that $XY = \frac{22\sqrt{221}}{50\cot \beta + 55}$ km.

(ii) Find the rate of change of β when CY is the shortest distance between city C and the road.

(8 marks)

· · ·			
 	-		
· · · · · · · · · · · · · · · · · · ·			
 ·			
		<u>.</u>	

EP(M2) MOCK 7-19

Answers written in the margins will not be marked.

– 163 –

Go on to the next page

		1
	<u> </u>	
	·	
Ι΄		

- 164 -

© Hong Kong Educational Publishing Company

		 ·
		-
	 	
	 1	

EP(M2) MOCK 7-21

– 165 –

Go on to the next page

13.	(a)	Let T be a positive number such	that $f(x)$	=f(T-x)	c) for a	all rea	l values	of x . By	using the
		substitution $u = T - x$, show that	$\int_0^T x f(x)$	$dx = \frac{T}{2} \int_0^T$	f(x) d	dx.			
									(4 marks)
					- π		2	- 1	1

- (b) (i) Using a suitable substitution, or otherwise, show that $\int_0^{\pi} \frac{\sin x \cos^2 x}{1 + \cos^2 x} dx = 2 \int_{-1}^{1} \frac{1}{1 + x^2} dx.$
 - (ii) Hence, or otherwise, evaluate $\int_0^{\pi} \frac{\sin x \cos^2 x}{1 + \cos^2 x} dx$.

(6 marks)

(c) Using the results of (a) and (b), evaluate $\int_0^{\pi} \frac{x \sin x \cos^2 x}{1 + \cos^2 x} dx$.

(3 marks)

Answers written in the margins will not be marked.

•			
	.	 	

	,
8	
Allowers without in the margins will not be marked.	
į	
	· · · · · · · · · · · · · · · · · · ·