

HKDSE
Mathematics
Extended Part
Module 2

Past Papers (by topic)
2012 – 2019
Solution

Content

Mathematical Induction	3
Binomial Theorem	8
Compound Angle Formulae.....	11
Differentiation (First principles).....	16
Applications of Differentiation	19
Technique of Integration	35
Applications of Integration	38
Matrices	66
Systems of Linear Equations.....	82
Vectors	92

Mathematical Induction

1.

3. For $n = 1$,

$$\text{L.H.S.} = 1 \times 2 = 2 \quad \text{and R.H.S.} = 1^2(1+1) = 2$$

\therefore L.H.S. = R.H.S. and the statement is true for $n = 1$.

Assume $1 \times 2 + 2 \times 5 + 3 \times 8 + \cdots + k(3k-1) = k^2(k+1)$, where k is a positive integer.

$$1 \times 2 + 2 \times 5 + 3 \times 8 + \cdots + k(3k-1) + (k+1)[3(k+1)-1]$$

$$= k^2(k+1) + (k+1)(3k+2) \quad \text{by the assumption}$$

$$= (k+1)(k^2 + 3k + 2)$$

$$= (k+1)^2(k+1+1)$$

Hence the statement is true for $n = k+1$.

By the principle of mathematical induction, the statement is true for all positive integers n .

1

1

1

1

Follow through

(5)

[HKDSE 2012 M2 #3]

2.

3. For $n = 1$,

$$\text{L.H.S. } 1 + \frac{1}{1 \times 4} = \frac{5}{4} \quad \text{and R.H.S.} = \frac{4(1)+1}{3(1)+1} = \frac{5}{4}$$

\therefore L.H.S. = R.H.S. and the statement is true for $n = 1$.

Assume $1 + \frac{1}{1 \times 4} + \frac{1}{4 \times 7} + \frac{1}{7 \times 10} + \cdots + \frac{1}{(3k-2)(3k+1)} = \frac{4k+1}{3k+1}$, where k is a positive integer.

1

1

$$1 + \frac{1}{1 \times 4} + \frac{1}{4 \times 7} + \frac{1}{7 \times 10} + \cdots + \frac{1}{(3k-2)(3k+1)} + \frac{1}{[3(k+1)-2][3(k+1)+1]}$$

1

$$= \frac{4k+1}{3k+1} + \frac{1}{(3k+1)(3k+4)} \quad \text{by the assumption}$$

$$= \frac{(12k^2 + 19k + 4) + 1}{(3k+1)(3k+4)}$$

$$= \frac{(3k+1)(4k+5)}{(3k+1)(3k+4)}$$

$$= \frac{4(k+1)+1}{3(k+1)+1}$$

Hence the statement is true for $n = k+1$.

By the principle of mathematical induction, the statement is true for all positive integers n .

1

1

Follow through

(5)

[HKDSE 2013 M2 #3]

3.

8. (a) Note that $\sin \frac{x}{2} \cos \frac{(1+1)x}{2} = \sin \frac{x}{2} \cos x$.

So, the statement is true for $n=1$.

Assume that $\sin \frac{x}{2} \sum_{k=1}^m \cos kx = \sin \frac{mx}{2} \cos \frac{(m+1)x}{2}$ for some positive integer m .

$$\sin \frac{x}{2} \sum_{k=1}^{m+1} \cos kx$$

$$= \sin \frac{x}{2} \sum_{k=1}^m \cos kx + \sin \frac{x}{2} \cos(m+1)x$$

$$= \sin \frac{mx}{2} \cos \frac{(m+1)x}{2} + \sin \frac{x}{2} \cos(m+1)x \quad (\text{by induction assumption})$$

$$= \frac{1}{2} \left(\sin \frac{(2m+1)x}{2} - \sin \frac{x}{2} \right) + \frac{1}{2} \left(\sin \frac{(2m+3)x}{2} - \sin \frac{(2m+1)x}{2} \right)$$

$$= \frac{1}{2} \left(\sin \frac{(2m+3)x}{2} - \sin \frac{x}{2} \right)$$

$$= \cos \frac{(2m+4)x}{4} \sin \frac{(2m+2)x}{4}$$

$$= \sin \frac{(m+1)x}{2} \cos \frac{(m+2)x}{2}$$

So, the statement is true for $n=m+1$ if it is true for $n=m$.

By mathematical induction, we have $\sin \frac{x}{2} \sum_{k=1}^n \cos kx = \sin \frac{nx}{2} \cos \frac{(n+1)x}{2}$

for all positive integers n .

1

1M

1M

1M

1M

1

(b) Putting $x = \frac{\pi}{7}$ and $n = 567$ in (a), we have

$$\sum_{k=1}^{567} \cos \frac{k\pi}{7}$$

$$= \frac{\sin \frac{(567)\pi}{7} \cos \frac{(568)\pi}{7}}{\sin \frac{\pi}{7}}$$

$$= \frac{\sin \frac{\pi}{2} \cos \left(\frac{\pi}{2} + \frac{\pi}{14} \right)}{\sin \frac{\pi}{14}}$$

$$= \frac{-\sin \frac{\pi}{14}}{\sin \frac{\pi}{14}}$$

$$= -1$$

1M

1A

----- (8)

[HKDSE 2015 M2 #8]

4.

5. (a) Note that $(-1)^1(1^2) = -1 = \frac{(-1)^1(1)(2)}{2}$.

So, the statement is true for $n=1$.

Assume that $\sum_{k=1}^m (-1)^k k^2 = \frac{(-1)^m m(m+1)}{2}$ for some positive integer m .

$$\sum_{k=1}^{m+1} (-1)^k k^2$$

$$= \sum_{k=1}^m (-1)^k k^2 + (-1)^{m+1}(m+1)^2$$

$$= \frac{(-1)^m m(m+1)}{2} + (-1)^{m+1}(m+1)^2 \quad (\text{by induction assumption})$$

$$= \frac{(-1)^m m(m+1) + (-1)^{m+1}(m+1)(2m+2)}{2}$$

$$= \frac{(-1)^m (m+1)(m-2m-2)}{2}$$

$$= \frac{(-1)^{m+1}(m+1)(m+2)}{2}$$

So, the statement is true for $n=m+1$ if it is true for $n=m$.

By mathematical induction, the statement is true for all positive integers n .

1

1M

for using induction assumption

1

(b) Putting $n=333$ in (a), we have $\sum_{k=1}^{333} (-1)^k k^2 = \frac{(-1)^{333}(333)(334)}{2}$.

1M

So, we have $-1+4+\sum_{k=3}^{333} (-1)^k k^2 = \frac{(-1)^{333}(333)(334)}{2}$.

1A

Thus, we have $\sum_{k=3}^{333} (-1)^{k+1} k^2 = 55\,614$.

[HKDSE 2016 M2 #5]

5.

6. (a) Note that $(1)(1+4) = 5 = \frac{(1)(2)(15)}{6}$.

So, the statement is true for $n=1$.

Assume that $\sum_{k=1}^n k(k+4) = \frac{m(m+1)(2m+13)}{6}$ for some positive integer m .

$$\begin{aligned} & \sum_{k=1}^{n+1} k(k+4) \\ &= \sum_{k=1}^n k(k+4) + (m+1)(m+5) \\ &= \frac{m(m+1)(2m+13)}{6} + (m+1)(m+5) \quad (\text{by induction assumption}) \\ &= \frac{(m+1)(2m^2+13m+6m+30)}{6} \\ &= \frac{(m+1)(2m^2+19m+30)}{6} \\ &= \frac{(m+1)(m+2)(2m+15)}{6} \end{aligned}$$

So, the statement is true for $n=m+1$ if it is true for $n=m$.

By mathematical induction, the statement is true for all positive integers n .

1

1M

for using induction assumption

(b) Putting $n=555$ in (a), we have

$$\sum_{k=1}^{555} k(k+4) = \frac{(555)(556)(1123)}{6} = 57\ 755\ 890.$$

Putting $n=332$ in (a), we have

$$\sum_{k=1}^{332} k(k+4) = \frac{(332)(333)(677)}{6} = 12\ 474\ 402.$$

1M

either one

$$\begin{aligned} & \sum_{k=333}^{555} \left(\binom{k}{112} \binom{k+4}{223} \right) \\ &= \left(\frac{1}{112} \binom{1}{223} \right) \sum_{k=333}^{555} k(k+4) \\ &= \frac{1}{(112)(223)} \left(\sum_{k=1}^{555} k(k+4) - \sum_{k=1}^{332} k(k+4) \right) \\ &= \frac{1}{24\ 976} (57\ 755\ 890 - 12\ 474\ 402) \\ &= 1813 \end{aligned}$$

1M

either one

1A

[HKDSE 2018 M2 #6]

6.

5. (a) Note that $\frac{1}{(1)(2)} + \frac{1}{(2)(3)} = \frac{2}{3} = \frac{1+1}{(1)(2+1)}$.

So, the statement is true for $n=1$.

Assume that $\sum_{k=m}^{2m} \frac{1}{k(k+1)} = \frac{m+1}{m(2m+1)}$, where m is a positive integer.

$$\begin{aligned} & \sum_{k=m+1}^{2m+2} \frac{1}{k(k+1)} \\ &= \sum_{k=m}^{2m} \frac{1}{k(k+1)} - \frac{1}{m(m+1)} + \frac{1}{(2m+1)(2m+2)} + \frac{1}{(2m+2)(2m+3)} \\ &= \frac{m+1}{m(2m+1)} - \frac{1}{m(m+1)} + \frac{1}{(2m+1)(2m+2)} + \frac{1}{(2m+2)(2m+3)} \\ &= \frac{(m+1)^2 - (2m+1)}{m(m+1)(2m+1)} + \frac{(2m+3)+(2m+1)}{(2m+1)(2m+2)(2m+3)} \\ &= \frac{m}{(m+1)(2m+1)} + \frac{2}{(2m+1)(2m+3)} \\ &= \frac{(2m+1)(m+2)}{(m+1)(2m+1)(2m+3)} \\ &= \frac{m+2}{(m+1)(2m+3)} \end{aligned}$$

So, the statement is true for $n = m+1$ if it is true for $n = m$.

By mathematical induction, the statement is true for all positive integers n .

1

1M

can be absorbed

1M

for using induction assumption

1

(b) Putting $n = 50$ in (a), we have $\sum_{k=50}^{100} \frac{1}{k(k+1)} = \frac{51}{(50)(101)} = \frac{51}{5050}$.

1M

either one

Putting $n = 100$ in (a), we have $\sum_{k=100}^{200} \frac{1}{k(k+1)} = \frac{101}{(100)(201)} = \frac{101}{20100}$.

So, we have $\sum_{k=50}^{200} \frac{1}{k(k+1)} = \frac{51}{5050} + \frac{101}{20100} - \frac{1}{(100)(101)}$.

Thus, we have $\sum_{k=50}^{200} \frac{1}{k(k+1)} = \frac{151}{10050}$.

1A

----- (7)

[HKDSE 2019 M2 #5]

Binomial Theorem

1.

$$2. \quad (1+ax)^n = 1 + nax + \frac{n(n-1)}{2}(ax)^2 + \dots$$

$$\text{Hence } na = 6 \text{ and } \frac{n(n-1)}{2}a^2 = 16.$$

$$\text{Solving, } \frac{n(n-1)}{2} \left(\frac{6}{n}\right)^2 = 16$$

$$18(n-1) = 16n$$

$$n = 9$$

$$\text{Therefore, } a = \frac{2}{3}.$$

	1M+1A 1M 1A 1A (5)
--	--------------------------------

[HKDSE 2012 M2 #2]

2.

$$2. \quad (1+ax)^n = 1 + C_1^n ax + C_2^n (ax)^2 + \dots$$

$$\begin{cases} na = -20 & \text{--- (1)} \\ \frac{n(n-1)}{2}a^2 = 180 & \text{--- (2)} \end{cases}$$

$$(2) \div (1)^2 :$$

$$\frac{n-1}{2n} = \frac{180}{400}$$

$$n = 10$$

$$\therefore a = -2$$

	1M 1M 1A 1A (4)
--	-----------------------------

[HKDSE 2013 M2 #2]

3.

$$1. \quad (a) \quad (1-4x)^2(1+x)^n$$

$$= (1-8x+16x^2) \left[1 + nx + \frac{n(n-1)}{2}x^2 + \dots \right] \quad (*)$$

$$\text{Coefficient of } x = n-8$$

$$\therefore n-8=1$$

$$\text{i.e., } n=9$$

$$(b) \quad \therefore (1-4x)^2(1+x)^9 = (1-8x+16x^2)(1+9x+36x^2+\dots)$$

$$\text{Coefficient of } x^2 = 36 - 8 \cdot 9 + 16$$

Alternative Solution

$$\text{Coefficient of } x^2 = \frac{n(n-1)}{2} - 8n + 16 \quad \text{by (*)}$$

$$= -20$$

	1M 1A 1M For binomial expansion of $(1+x)^n$ up to the (x^2) term 1M 1A (4)
--	---

[HKDSE 2014 M2 #1]

4.

1. $\begin{aligned} & (5+x)^4 \\ &= 5^4 + 4(5^3)x + 6(5^2)x^2 + 4(5)x^3 + x^4 \\ &= 625 + 500x + 150x^2 + 20x^3 + x^4 \end{aligned}$ $\begin{aligned} & \left(1 - \frac{2}{x}\right)^3 \\ &= 1 - 3\left(\frac{2}{x}\right) + 3\left(\frac{2}{x}\right)^2 - \left(\frac{2}{x}\right)^3 \\ &= 1 - \frac{6}{x} + \frac{12}{x^2} - \frac{8}{x^3} \end{aligned}$ The constant term $= (625)(1) + (500)(-6) + (150)(12) + (20)(-8)$ $= -735$	1M 1A 1M 1M 1A ----- (5)	withhold 1M if the step is skipped
---	---	------------------------------------

[HKDSE 2016 M2 #1]

5.

2. Note that $(1+ax)^8 = 1 + C_1^8 ax + C_2^8 (ax)^2 + \dots + (ax)^8$ and $(b+x)^9 = b^9 + C_1^9 b^8 x + C_2^9 b^7 x^2 + \dots + C_7^9 b^2 x^7 + C_8^9 b x^8 + x^9$. Also note that that $\lambda_2 : \mu_7 = 7 : 4$ and $\lambda_1 + \mu_8 + 6 = 0$. Therefore, we have $\frac{C_2^8 a^2}{C_7^9 b^2} = \frac{7}{4}$ and $8a + 9b + 6 = 0$. So, we have $4a^2 = 9b^2$ and $8a + 9b + 6 = 0$. Hence, we have $4a^2 - 9\left(\frac{-8a - 6}{9}\right)^2 = 0$. Simplifying, we have $7a^2 + 24a + 9 = 0$. Thus, we have $a = -3$ or $a = \frac{-3}{7}$.	1M 1M 1M 1M 1M 1A ----- (5)	for either one for $pa^2 + qa + r = 0$ for both correct
---	---	---

[HKDSE 2017 M2 #2]

Compound Angle Formulae


1.

10. (a) Let F be a point on AB such that $OF \perp AB$. Let OA be x .

$$\therefore AF = \frac{1}{2} \quad \text{and} \quad \angle AOF = 2\theta \quad (\text{properties of isos. } \Delta)$$

$$\text{In } \triangle OAF, \sin 2\theta = \frac{\frac{1}{2}}{x}$$

1M



OR

$$x^2 = x^2 + 1^2 - 2x \cos(90^\circ - 2\theta)$$

Alternative Solution

$$\text{In } \triangle OAB, \frac{1}{\sin 4\theta} = \frac{x}{\sin(90^\circ - 2\theta)}$$

1M

$$x = \frac{1}{2 \sin 2\theta} \quad \text{----- (1)}$$

1M

$$\text{In } \triangle OAY, \frac{y}{\sin \theta} = \frac{x}{\sin(90^\circ + \theta)} \quad \text{----- (2)}$$

$$\text{Substitute (1) into (2): } \frac{y}{\sin \theta} = \frac{1}{2 \sin 2\theta} \cdot \frac{1}{\cos \theta}$$

1M

Alternative Solution

$$\text{In } \triangle OAY, \frac{y}{\sin \theta} = \frac{OY}{\sin \angle OAY}$$

1M

$$\text{In } \triangle OBY, \frac{1-y}{\sin 3\theta} = \frac{OY}{\sin \angle OBY}$$

1M

$\because \angle OAY = \angle OBY$ (base \angle s, isos. Δ s)

$$\therefore \frac{y}{\sin \theta} = \frac{1-y}{3 \sin \theta - 4 \sin^3 \theta}$$

1M

$$3y - 4y \sin^2 \theta = 1 - y$$

$$4y(1 - \sin^2 \theta) = 1$$

1M

$$y = \frac{1}{4} \sec^2 \theta$$

1

- (b) $\because 0^\circ < 4\theta < 180^\circ$

$$\therefore 0^\circ < \theta < 45^\circ$$

$$\frac{1}{4} \sec^2 0^\circ < y < \frac{1}{4} \sec^2 45^\circ \quad [\text{since } \sec^2 \theta \text{ is an increasing function for } 0^\circ < \theta < 45^\circ]$$

$$\text{i.e. } \frac{1}{4} < y < \frac{1}{2}$$

1A

(6)

[HKDSE 2012 M2 #10]

} Accept using “≤” sign

2.

$$\begin{aligned}
 7. \quad (a) \quad R.H.S. &= \frac{\sin 2x}{1 + \cos 2x} \\
 &= \frac{2\sin x \cos x}{1 + 2\cos^2 x - 1} \\
 &= \frac{\sin x}{\cos x} \\
 &= \tan x \\
 &= L.H.S.
 \end{aligned}$$

1M For either formula

1

$$\begin{aligned}
 (b) \quad R.H.S. &= \frac{\sin 8y \cos 4y \cos 2y}{(1 + \cos 8y)(1 + \cos 4y)(1 + \cos 2y)} \\
 &= \tan 4y \cdot \frac{\cos 4y \cos 2y}{(1 + \cos 4y)(1 + \cos 2y)} \quad \text{by (a)} \\
 &= \frac{\sin 4y \cos 2y}{(1 + \cos 4y)(1 + \cos 2y)} \\
 &= \tan 2y \cdot \frac{\cos 2y}{1 + \cos 2y} \quad \text{by (a)} \\
 &= \frac{\sin 2y}{1 + \cos 2y} \\
 &= \tan y \quad \text{by (a)} \\
 &= L.H.S.
 \end{aligned}$$

1M

1M

1

(5)

[HKDSE 2013 M2 #7]

3.

$$\begin{aligned}
 7. \quad (a) \quad &\sin^2 x \cos^2 x \\
 &= \frac{(2\sin x \cos x)^2}{4} \\
 &= \frac{\sin^2 2x}{4} \\
 &= \frac{1}{4} \left(\frac{1 - \cos 4x}{2} \right) \\
 &= \frac{1 - \cos 4x}{8}
 \end{aligned}$$

1M

1

$$\begin{aligned}
 (b) \quad (i) \quad f(x) &= \sin^4 x + \cos^4 x \\
 &= (\sin^2 x + \cos^2 x)^2 - 2\sin^2 x \cos^2 x \\
 &= 1^2 - 2 \left(\frac{1 - \cos 4x}{8} \right) \quad (\text{by (a)}) \\
 &= \frac{1}{4} \cos 4x + \frac{3}{4}
 \end{aligned}$$

1M

for using (a)

1A

$$\begin{aligned}
 (ii) \quad 8f(x) = 7 \\
 8 \left(\frac{1}{4} \cos 4x + \frac{3}{4} \right) = 7
 \end{aligned}$$

1M

for using the result of (b)(i)

$$2\cos 4x + 6 = 7$$

$$\cos 4x = \frac{1}{2}$$

$$4x = \frac{\pi}{3} \quad \text{or} \quad 4x = \frac{5\pi}{3}$$

$$x = \frac{\pi}{12} \quad \text{or} \quad x = \frac{5\pi}{12}$$

1A

for both

[HKDSE 2015 M2 #7]

4.

<p>6. (a) Note that $4(-1)^3 + 2(-1)^2 - 3(-1) - 1 = -4 + 2 + 3 - 1 = 0$. Thus, $x+1$ is a factor of $4x^3 + 2x^2 - 3x - 1$.</p> <p>(b)</p> $\begin{aligned} & \cos 3\theta \\ &= \cos(\theta + 2\theta) \\ &= \cos \theta \cos 2\theta - \sin \theta \sin 2\theta \\ &= \cos \theta (\cos^2 \theta - \sin^2 \theta) - \sin \theta (2 \sin \theta \cos \theta) \\ &= \cos \theta (\cos^2 \theta - (1 - \cos^2 \theta)) - 2 \cos \theta (1 - \cos^2 \theta) \\ &= 4 \cos^3 \theta - 3 \cos \theta \end{aligned}$ <p>(c) Note that $\theta = \frac{3\pi}{5}$ satisfies $\cos 3\theta = \cos(3\pi - 2\theta)$. Therefore, $\theta = \frac{3\pi}{5}$ satisfies $\cos 3\theta = -\cos 2\theta$. By (b), $\theta = \frac{3\pi}{5}$ satisfies $4 \cos^3 \theta - 3 \cos \theta = -(2 \cos^2 \theta - 1)$. Hence, $\cos \frac{3\pi}{5}$ is a root of the equation $4x^3 + 2x^2 - 3x - 1 = 0$. By (a), we have $4x^3 + 2x^2 - 3x - 1 = (x+1)(4x^2 - 2x - 1)$. So, we have $4x^3 + 2x^2 - 3x - 1 = 4(x+1) \left(x - \frac{1+\sqrt{5}}{4} \right) \left(x - \frac{1-\sqrt{5}}{4} \right)$. Since $\frac{\pi}{2} < \frac{3\pi}{5} < \pi$, we have $-1 < \cos \frac{3\pi}{5} < 0$. Thus, we have $\cos \frac{3\pi}{5} = \frac{1-\sqrt{5}}{4}$.</p>	1	1M	1A	1M	1M	for using (b)
					1	-----(6)

[HKDSE 2016 M2 #6]

5.

7. (a)

$$\begin{aligned}
 & \sin 3x \\
 &= \sin(x + 2x) \\
 &= \sin x \cos 2x + \cos x \sin 2x \\
 &= \sin x(\cos^2 x - \sin^2 x) + 2 \sin x \cos^2 x \\
 &= \sin x(1 - 2 \sin^2 x) + 2 \sin x(1 - \sin^2 x) \\
 &= 3 \sin x - 4 \sin^3 x
 \end{aligned}$$

1M

1

(b) (i)

$$\begin{aligned}
 & \frac{\sin 3\left(x - \frac{\pi}{4}\right)}{\sin\left(x - \frac{\pi}{4}\right)} \\
 &= \frac{\sin\left(3x - \frac{3\pi}{4}\right)}{\sin\left(x - \frac{\pi}{4}\right)} \\
 &= \frac{\sin 3x \cos \frac{3\pi}{4} - \cos 3x \sin \frac{3\pi}{4}}{\sin x \cos \frac{\pi}{4} - \cos x \sin \frac{\pi}{4}} \\
 &= \frac{-\frac{1}{\sqrt{2}}(\sin 3x + \cos 3x)}{\frac{1}{\sqrt{2}}(\sin x - \cos x)} \\
 &= \frac{\cos 3x + \sin 3x}{\cos x - \sin x}
 \end{aligned}$$

1M

1

(ii)

$$\frac{\cos 3x + \sin 3x}{\cos x - \sin x} = 2$$

$$\frac{\sin 3\left(x - \frac{\pi}{4}\right)}{\sin\left(x - \frac{\pi}{4}\right)} = 2 \quad (\text{by (b)(i)})$$

1M

for using (b)(i)

Note that $\sin\left(x - \frac{\pi}{4}\right) \neq 0$.

$$3 - 4 \sin^2\left(x - \frac{\pi}{4}\right) = 2 \quad (\text{by (a)})$$

1M

for using (a)

$$1 - 4 \sin^2\left(x - \frac{\pi}{4}\right) = 0$$

$$\left(1 - 2 \sin\left(x - \frac{\pi}{4}\right)\right) \left(1 + 2 \sin\left(x - \frac{\pi}{4}\right)\right) = 0$$

1M

Since $\frac{\pi}{4} < x < \frac{\pi}{2}$, we have $\sin\left(x - \frac{\pi}{4}\right) = \frac{1}{2}$.

Therefore, we have $x - \frac{\pi}{4} = \frac{\pi}{6}$.

Thus, we have $x = \frac{5\pi}{12}$.

1A

----- (8)

[HKDSE 2017 M2 #7]

6.

3. (a) $\cot A = 3 \cot B$

$$\frac{\cos A}{\sin A} = \frac{3 \cos B}{\sin B}$$

$$3 \sin A \cos B = \cos A \sin B$$

1M

$$\sin(A+B) - 2 \sin(B-A)$$

$$= (\sin A \cos B + \cos A \sin B) - 2(\sin B \cos A - \cos B \sin A)$$

$$= 3 \sin A \cos B - \cos A \sin B$$

$$= 0$$

Thus, we have $\sin(A+B) = 2 \sin(B-A)$.

1

(b) $\cot\left(x + \frac{4\pi}{9}\right) = 3 \cot\left(x + \frac{5\pi}{18}\right)$

By letting $A = x + \frac{4\pi}{9}$ and $B = x + \frac{5\pi}{18}$, we have $\cot A = 3 \cot B$.

1M

By (a), we have $\sin(A+B) = 2 \sin(B-A)$.

With the help of $\sin\left(\frac{-\pi}{6}\right) = \frac{-1}{2}$, we have $\sin\left(2x + \frac{13\pi}{18}\right) = -1$.

1M

Noting that $0 \leq x \leq \frac{\pi}{2}$, we have $x = \frac{7\pi}{18}$.

1A

Since $\cot\left(\frac{7\pi}{18} + \frac{4\pi}{9}\right) = -\sqrt{3} = 3 \cot\left(\frac{7\pi}{18} + \frac{5\pi}{18}\right)$, the required solution

of the equation $\cot\left(x + \frac{4\pi}{9}\right) = 3 \cot\left(x + \frac{5\pi}{18}\right)$ is $x = \frac{7\pi}{18}$.

-----(5)

[HKDSE 2018 M2 #3]

Differentiation (First principles)

1.

$$f(x) = e^{2x}$$

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{e^{2(0+h)} - e^{2(0)}}{h} \\ &= \lim_{h \rightarrow 0} \frac{e^{2h} - 1}{2h} \cdot 2 \\ &= 2 \end{aligned}$$

1M

1M

1A

(3)

Accept $\lim_{h \rightarrow 0} \frac{e^{2h} - 1}{h}$

[HKDSE 2012 M2 #1]

2.

$$1. \quad \frac{d}{dx}(\sin 2x) = \lim_{h \rightarrow 0} \frac{\sin 2(x+h) - \sin 2x}{h}$$

$$= \lim_{h \rightarrow 0} \left(\frac{2}{h} \cos \frac{2x+2h+2x}{2} \sin \frac{2x+2h-2x}{2} \right)$$

$$= \lim_{h \rightarrow 0} \left[2 \cos(2x+h) \frac{\sin h}{h} \right]$$

$$= 2 \lim_{h \rightarrow 0} \cos(2x+h) \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h}$$

$$= 2 \cos 2x$$

1M

1M

1M

1A

(4)

[HKDSE 2013 M2 #1]

3.

$$2. \quad (a) \quad y = x^3 - 3x$$

$$\begin{aligned} \frac{dy}{dx} &= \lim_{h \rightarrow 0} \frac{[(x+h)^3 - 3(x+h)] - (x^3 - 3x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - 3x - 3h - x^3 + 3x}{h} \\ &= \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2 - 3) \\ &= 3x^2 - 3 \end{aligned}$$

1M

1M

1A

1M

1A

(5)

OR $\frac{h[(x+h)^2 + (x+h)x + x^2] - 3h}{h}$

Accept $3x^2 - 3 < 0$

Accept $-1 < x < 1$

[HKDSE 2014 M2 #2]

4.

$$\begin{aligned}
 1. \quad & \frac{d}{dx}(x^5 + 4) \\
 &= \lim_{h \rightarrow 0} \frac{((x+h)^5 + 4) - (x^5 + 4)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{x^5 + 5hx^4 + 10h^2x^3 + 10h^3x^2 + 5h^4x + h^5 + 4 - x^5 - 4}{h} \\
 &= \lim_{h \rightarrow 0} (5x^4 + 10hx^3 + 10h^2x^2 + 5h^3x + h^4) \\
 &= 5x^4
 \end{aligned}$$

1M

1M for binomial expansion

1M withhold 1M if the step is skipped

1A

-----(4)

[HKDSE 2015 M2 #1]

5.

$$\begin{aligned}
 2. \quad & \frac{1}{\sqrt{x}} - \frac{1}{\sqrt{x+h}} \\
 &= \frac{\sqrt{x+h} - \sqrt{x}}{\sqrt{x(x+h)}} \\
 &= \left(\frac{\sqrt{x+h} - \sqrt{x}}{\sqrt{x(x+h)}} \right) \left(\frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \right) \\
 &= \frac{x+h-x}{(x+h)\sqrt{x}+x\sqrt{x+h}} \\
 &= \frac{h}{(x+h)\sqrt{x}+x\sqrt{x+h}}
 \end{aligned}$$

1M

1

$$\begin{aligned}
 & \frac{d}{dx} \sqrt{\frac{3}{x}} \\
 &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{\sqrt{3}}{\sqrt{x+h}} - \frac{\sqrt{3}}{\sqrt{x}} \right) \\
 &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{-h\sqrt{3}}{(x+h)\sqrt{x}+x\sqrt{x+h}} \right) \\
 &= \lim_{h \rightarrow 0} \frac{-\sqrt{3}}{(x+h)\sqrt{x}+x\sqrt{x+h}} \\
 &= \frac{-\sqrt{3}}{2x\sqrt{x}}
 \end{aligned}$$

1M

1

withhold 1M if the step is omitted

1A

-----(5)

[HKDSE 2016 M2 #2]

6.

$$\begin{aligned}
 1. \quad & \frac{d}{d\theta} \sec 6\theta \\
 &= \lim_{h \rightarrow 0} \frac{\sec 6(\theta+h) - \sec 6\theta}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos 6\theta - \cos 6(\theta+h)}{h \cos 6(\theta+h) \cos 6\theta} \\
 &= \lim_{h \rightarrow 0} \frac{2\sin(6\theta+3h)\sin 3h}{h \cos 6(\theta+h) \cos 6\theta} \\
 &= 6 \left(\lim_{h \rightarrow 0} \frac{\sin 3h}{3h} \right) \left(\lim_{h \rightarrow 0} \frac{\sin(6\theta+3h)}{\cos 6(\theta+h) \cos 6\theta} \right) \\
 &= 6 \left(1 \right) \left(\frac{\sin 6\theta}{\cos^2 6\theta} \right) \\
 &= 6 \sec 6\theta \tan 6\theta
 \end{aligned}$$

1M

1M

1M

1M withhold 1M if the step is skipped

1A

-----(5)

[HKDSE 2017 M2 #1]

7.

$$\begin{aligned} 1. \quad & f(1+h) \\ &= ((1+h)^2 - 1)e^{1+h} \\ &= (2h + h^2)e^{1+h} \end{aligned}$$

1A

$$\begin{aligned} & f'(1) \\ &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(2h + h^2)e^{1+h} - 0}{h} \\ &= \lim_{h \rightarrow 0} (2 + h)e^{1+h} \\ &= 2e \end{aligned}$$

1M

1M

1A

withhold 1M if this step is skipped

----- (4)

[HKDSE 2018 M2 #1]

8.

$$\begin{aligned} 1. \quad & f(1+h) - f(1) \\ &= \frac{10(1+h)}{7+3(1+h)^2} - \frac{10}{7+3} \\ &= \frac{10+10h}{10+6h+3h^2} - 1 \\ &= \frac{10+10h-10-6h-3h^2}{10+6h+3h^2} \\ &= \frac{4h-3h^2}{10+6h+3h^2} \end{aligned}$$

1

$$\begin{aligned} & f'(1) \\ &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{4h-3h^2}{10+6h+3h^2} \right) \\ &= \lim_{h \rightarrow 0} \frac{4-3h}{10+6h+3h^2} \\ &= \frac{4-3(0)}{10+6(0)+3(0)^2} \\ &= \frac{2}{5} \end{aligned}$$

1M

1M

withhold 1M if the step is skipped

1A 0.4

----- (4)

[HKDSE 2019 M2 #1]

Applications of Differentiation

1.

5. $y = \frac{x^2 + x + 1}{x+1}$
 $= x + \frac{1}{x+1}$
 $\frac{dy}{dx} = 1 - \frac{1}{(x+1)^2}$
 $\frac{dy}{dx} = 0 \text{ when } x = -2 \text{ or } 0$

x	$x < -2$	-2	$-2 < x < -1$	-1	$-1 < x < 0$	0	$x > 0$
$\frac{dy}{dx}$	> 0	0	< 0	undefined	< 0	0	> 0

1M
1A
OR $\frac{x^2 + 2x}{(x+1)^2}$

1M

Alternative Solution

$$\frac{d^2y}{dx^2} = \frac{2}{(x+1)^3}$$

When $x = 0$, $\frac{d^2y}{dx^2} = 2 > 0$; when $x = -2$, $\frac{d^2y}{dx^2} = -2 < 0$.

} 1M

 Hence $(0, 1)$ is a minimum point.

 The vertical asymptote is $x = -1$.

$$\because \lim_{x \rightarrow \pm\infty} \frac{1}{x+1} = 0$$

 ∴ the oblique asymptote is $y = x$

 1A
1A

Accept $\lim_{x \rightarrow \infty} \frac{1}{x+1} = 0$

 1A
(6)

[HKDSE 2012 M2 #5]

2.

6. (a) Let the radius of the water surface be
- a
- cm.


By considering similar triangles, $\frac{a-3}{4-3} = \frac{h}{10}$.

i.e. $a = \frac{h+30}{10}$

$$\therefore V = \frac{\pi}{3} h \left[3^2 + 3 \left(\frac{h+30}{10} \right) + \left(\frac{h+30}{10} \right)^2 \right]$$

$$= \frac{\pi}{300} h [900 + 30(h+30) + (h^2 + 60h + 900)]$$

$$= \frac{\pi}{300} (h^3 + 90h^2 + 2700h)$$

 1M
1M
1

 1M+1A
1A
(6)

(b) $\frac{dV}{dt} = \frac{\pi}{300} (3h^2 + 180h + 2700) \frac{dh}{dt}$

$$\therefore 7\pi = \frac{\pi}{300} [3(5)^2 + 180(5) + 2700] \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{4}{7}$$

i.e. the rate of increase of depth of water is $\frac{4}{7}$ cm s⁻¹.

[HKDSE 2012 M2 #6]

3.

5. (a) $f(x) = \frac{3-3x^2}{3+x^2}$

$\therefore f(0) = 1, f(1) = 0$ and $f(-1) = 0$

\therefore maximum point is $(0, 1)$,

and points of inflection are $(1, 0)$ and $(-1, 0)$.


(b) Since $x^2 + 3 > 0$, there is no vertical asymptote.

$$f(x) = -3 + \frac{12}{x^2 + 3}$$

When $x \rightarrow \pm\infty$, $y \rightarrow -3$.

Hence $y = -3$ is a horizontal asymptote.

(c)



1A

1A

For both

1M

OR $f(x) = \frac{\frac{3}{x^2} - 3}{\frac{3}{x^2} + 1}$

1A

For shape of $y = f(x)$

For all correct

(6)

[HKDSE 2013 M2 #5]

4.

$$12. \text{ (a) (i)} \quad T = \frac{PO}{7} + \frac{QB}{1.4}$$

$$= \frac{x}{7} + \frac{5\sqrt{30^2 + (40-x)^2}}{7}$$

$$= \frac{x + 5\sqrt{x^2 - 80x + 2500}}{7}$$

(ii) When T is minimum, $\frac{dT}{dx} = 0$.

$$\frac{1}{7} \left[1 + \frac{5(2x-80)}{2\sqrt{x^2 - 80x + 2500}} \right] = 0$$

$$5(x-40) = -\sqrt{x^2 - 80x + 2500}$$


$$25x^2 - 2000x + 40000 = x^2 - 80x + 2500$$

$$24x^2 - 160x + 3125 = 0$$

$$\therefore x = 40 - \frac{5\sqrt{6}}{2} \text{ or } 40 + \frac{5\sqrt{6}}{2} \text{ (rejected by checking)}$$

1M

$$\text{OR } \frac{x}{7} + \frac{\sqrt{x^2 - 80x + 2500}}{1.4}$$



1M

1

x	$0 < x < 40 - \frac{5\sqrt{6}}{2}$	$x = 40 - \frac{5\sqrt{6}}{2}$	$x > 40 - \frac{5\sqrt{6}}{2}$
$\frac{dT}{dx}$	-	0	+

So, when T is minimum, $x = 40 - \frac{5\sqrt{6}}{2}$.

$$QB = \sqrt{30^2 + \left[40 - \left(40 - \frac{5\sqrt{6}}{2} \right) \right]^2}$$

$$= \frac{25\sqrt{6}}{2} \text{ m}$$

1A

1

(6)

$$(b) \text{ (i)} \quad \sin \beta = \frac{30}{\frac{25\sqrt{6}}{2}} = \frac{2\sqrt{6}}{5}$$

$$\cos \beta = \frac{40 - \left(40 - \frac{5\sqrt{6}}{2} \right)}{\frac{25\sqrt{6}}{2}} = \frac{1}{5}$$

$$\text{In } \triangle MAB, \frac{MB}{\sin \alpha} = \frac{AB}{\sin(\pi - \alpha - \beta)}.$$

$$MB = \frac{40 \sin \alpha}{\sin(\alpha + \beta)}$$

$$= \frac{40 \sin \alpha}{\sin \alpha \cos \beta + \cos \alpha \sin \beta}$$

$$= \frac{40 \sin \alpha}{\frac{1}{5} \sin \alpha + \frac{2\sqrt{6}}{5} \cos \alpha}$$

$$= \frac{200 \tan \alpha}{\tan \alpha + 2\sqrt{6}}$$


1A

1A

1M

1M

1



$$(ii) \quad \frac{dMB}{dt} = 200 \cdot \frac{(\tan \alpha + 2\sqrt{6}) \sec^2 \alpha - \tan \alpha \sec^2 \alpha}{(\tan \alpha + 2\sqrt{6})^2} \cdot \frac{d\alpha}{dt}$$

$$= \frac{400\sqrt{6} \sec^2 \alpha}{(\tan \alpha + 2\sqrt{6})^2} \cdot \frac{d\alpha}{dt}$$

$$\therefore -1.4 = \frac{400\sqrt{6} \sec^2 0.2}{(\tan 0.2 + 2\sqrt{6})^2} \cdot \frac{d\alpha}{dt}$$

$$\frac{d\alpha}{dt} \approx -0.0357 \text{ rad s}^{-1}$$

1M

1A

[HKDSE 2013 M2 #12]

5.

3. $x \ln y + y = 2$

$$\ln y + x \cdot \frac{1}{y} \frac{dy}{dx} + \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = \frac{-y \ln y}{x+y}$$

1M+1M

 1M for product rule
1M for chain rule

Alternative Solution

$$x = \frac{2-y}{\ln y}$$

$$\frac{dx}{dy} = \frac{\ln y \cdot (-1) - (2-y)\frac{1}{y}}{(\ln y)^2}$$

$$\therefore \frac{dy}{dx} = \frac{y(\ln y)^2}{y-2-y \ln y}$$

1M

For quotient rule

 For $\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$

 When the curve cuts the y -axis, $x = 0$.

$$\therefore y = 2$$

1A

[HKDSE 2014 M2 #3]

6.

4. $x = 2y + \sin y$

$$\frac{dx}{dy} = 2 + \cos y$$

$$\frac{dy}{dx} = \frac{1}{2 + \cos y}$$

1M

$$\frac{d^2y}{dx^2} = -1 \cdot (2 + \cos y)^{-2} (-\sin y) \frac{dy}{dx}$$

1M

 OR $\frac{0 - 1(-\sin y) \frac{dy}{dx}}{(2 + \cos y)^2}$
Alternative Solution

$$1 = 2 \frac{dy}{dx} + \cos y \cdot \frac{dy}{dx}$$

$$0 = 2 \frac{d^2y}{dx^2} + \left[\cos y \frac{d^2y}{dx^2} + (-\sin y) \left(\frac{dy}{dx} \right)^2 \right]$$

$$\sin y \left(\frac{1}{2 + \cos y} \right)^2 = (2 + \cos y) \frac{d^2y}{dx^2}$$

1M

1M

$$\frac{d^2y}{dx^2} = \frac{\sin y}{(2 + \cos y)^3}$$

1A

(3)

[HKDSE 2014 M2 #4]

8.

2. (a) $\frac{dy}{dx}$ $= x \cos x + \sin x - \sin x$ $= x \cos x$	1M 1A	for product rule
$\frac{d^2y}{dx^2}$ $= -x \sin x + \cos x$	1A	
(b) $x \frac{d^2y}{dx^2} + k \frac{dy}{dx} + xy$ $= x(-x \sin x + \cos x) + kx \cos x + x(x \sin x + \cos x)$ (by (a)) $= (2+k)x \cos x$ Since $x \frac{d^2y}{dx^2} + k \frac{dy}{dx} + xy = 0$ for all real values of x , we have $k = -2$.	1M 1A	for using the results of (a)
		-----(5)

[HKDSE 2015 M2 #2]

9.

9. (a) $f'(x)$

$$= \frac{(x-2)(2x)-(x^2+12)}{(x-2)^2}$$

$$= \frac{x^2-4x-12}{(x-2)^2}$$

1M for quotient rule

1A

-----(2)

(b) Note that $f'(x) = \frac{(x+2)(x-6)}{(x-2)^2}$.

So, we have $f'(x) = 0 \Leftrightarrow x = -2$ or $x = 6$.

1A

x	$(-\infty, -2)$	-2	$(-2, 2)$	$(2, 6)$	6	$(6, \infty)$
$f'(x)$	+	0	-	-	0	+
$f(x)$	\nearrow	-4	\searrow	\searrow	12	\nearrow

1M

1

1

Thus, the maximum value and the minimum value of $f(x)$ are -4 and 12 respectively.

Note that $f'(x) = \frac{(x+2)(x-6)}{(x-2)^2}$ and $f''(x) = \frac{32}{(x-2)^3}$.

1A

So, we have $f'(x) = 0 \Leftrightarrow x = -2$ or $x = 6$.

1M

Further note that $f''(-2) = \frac{-1}{2} < 0$ and $f''(6) = \frac{1}{2} > 0$.

1

Thus, the maximum value and the minimum value of $f(x)$ are -4 and 12 respectively.

1

-----(4)

(c) The equation of the vertical asymptote is $x - 2 = 0$.

1A

Note that $f(x) = x + 2 + \frac{16}{x-2}$.

1M

Thus, the equation of the oblique asymptote is $y = x + 2$.

1A

-----(3)

(d) $\frac{x^2+12}{x-2} = 14$

1A

$$x^2 - 14x + 40 = 0$$

can be absorbed

$$x = 4 \text{ or } x = 10$$

The required area

1M

$$= \int_4^{10} \left(14 - \frac{x^2+12}{x-2} \right) dx$$

$$= \int_4^{10} \left(12 - x - \frac{16}{x-2} \right) dx$$

1M

$$= \left[12x - \frac{x^2}{2} - 16 \ln(x-2) \right]_4^{10}$$

1A

$$= 30 - 32 \ln 2$$

-----(4)

[HKDSE 2015 M2 #9]

10.

4. (a) Note that the equation of the vertical asymptote is $x = 1$. $\begin{aligned} f(x) &= \frac{2x^2 + x + 1}{x - 1} \\ &= 2x + 3 + \frac{4}{x - 1} \end{aligned}$ <p>Thus, the equation of the oblique asymptote is $y = 2x + 3$.</p>	1A 1M 1A	
(b) $f'(x)$ $\begin{aligned} &= \frac{(x-1)(4x+1)-(2x^2+x+1)}{(x-1)^2} \\ &= \frac{2(x^2-2x-1)}{(x-1)^2} \end{aligned}$ $\begin{aligned} f'(2) &= \frac{2(2^2-2(2)-1)}{(2-1)^2} \\ &= -2 \end{aligned}$ <p>The slope of the normal to G at the point $(2, 11)$</p> $\begin{array}{ c c } \hline & \frac{1}{f'(2)} \\ \hline \end{array}$ $= \frac{1}{-2} = -\frac{1}{2}$	1M 1M 1A	
$f'(x)$ $\begin{aligned} &= \frac{d}{dx} \left(2x + 3 + \frac{4}{x-1} \right) \\ &= 2 - \frac{4}{(x-1)^2} \end{aligned}$ $\begin{aligned} f'(2) &= 2 - \frac{4}{(2-1)^2} \\ &= -2 \end{aligned}$ <p>The slope of the normal to G at the point $(2, 11)$</p> $\begin{array}{ c c } \hline & \frac{1}{f'(2)} \\ \hline \end{array}$ $= \frac{1}{-2} = -\frac{1}{2}$	1M 1M 1A	
-----(7)		

[HKDSE 2016 M2 #4]

11.

9. (a) Since $f(x) = x^3 + ax^2 + bx + 5$, we have $f'(x) = 3x^2 + 2ax + b$. Note that $P(-1, 10)$ is a turning point of C . So, we have $-1 + a - b + 5 = 10$ and $3 - 2a + b = 0$. Therefore, we have $a - b - 6 = 0$ and $-2a + b + 3 = 0$. Solving, we have $a = -3$ and $b = -9$.	1A 1M 1A ----- (3)	for either for both												
(b) Note that $f''(x) = 6x - 6$. $\begin{aligned} f''(-1) \\ = -12 \\ < 0 \end{aligned}$ Thus, P is a maximum point of C .	1M 1A	f.t.												
Note that $f'(x) = 3(x+1)(x-3)$. <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 2px;">x</td><td style="padding: 2px;">$(-\infty, -1)$</td><td style="padding: 2px;">-1</td><td style="padding: 2px;">$(-1, 3)$</td></tr> <tr> <td style="padding: 2px;">$f'(x)$</td><td style="padding: 2px;">+</td><td style="padding: 2px;">0</td><td style="padding: 2px;">-</td></tr> <tr> <td style="padding: 2px;">$f(x)$</td><td style="padding: 2px;">↗</td><td style="padding: 2px;">10</td><td style="padding: 2px;">↘</td></tr> </table>	x	$(-\infty, -1)$	-1	$(-1, 3)$	$f'(x)$	+	0	-	$f(x)$	↗	10	↘	1M	
x	$(-\infty, -1)$	-1	$(-1, 3)$											
$f'(x)$	+	0	-											
$f(x)$	↗	10	↘											
Thus, P is a maximum point of C .	1A	f.t.												
----- (2) (c) Note that $f'(x) = 3(x+1)(x-3)$. So, we have $f'(3) = 0$. Also note that $f''(x) = 6x - 6$. $\begin{aligned} f''(3) \\ = 12 \\ > 0 \end{aligned}$ Further note that $f(3) = -22$. Thus, the minimum value of $f(x)$ is -22 .	1M 1A													
Note that $f'(x) = 3(x+1)(x-3)$. So, we have $f'(3) = 0$. <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 2px;">x</td><td style="padding: 2px;">$(-1, 3)$</td><td style="padding: 2px;">3</td><td style="padding: 2px;">$(3, \infty)$</td></tr> <tr> <td style="padding: 2px;">$f'(x)$</td><td style="padding: 2px;">-</td><td style="padding: 2px;">0</td><td style="padding: 2px;">+</td></tr> <tr> <td style="padding: 2px;">$f(x)$</td><td style="padding: 2px;">↘</td><td style="padding: 2px;">-22</td><td style="padding: 2px;">↗</td></tr> </table>	x	$(-1, 3)$	3	$(3, \infty)$	$f'(x)$	-	0	+	$f(x)$	↘	-22	↗	1M	
x	$(-1, 3)$	3	$(3, \infty)$											
$f'(x)$	-	0	+											
$f(x)$	↘	-22	↗											
Thus, the minimum value of $f(x)$ is -22 .	1A													
----- (2) (d) Note that $f''(x) = 6(x-1)$. Therefore, we have $f''(x) = 0$ when $x = 1$.	1M													
<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 2px;">x</td><td style="padding: 2px;">$(-\infty, 1)$</td><td style="padding: 2px;">1</td><td style="padding: 2px;">$(1, \infty)$</td></tr> <tr> <td style="padding: 2px;">$f''(x)$</td><td style="padding: 2px;">-</td><td style="padding: 2px;">0</td><td style="padding: 2px;">+</td></tr> </table>	x	$(-\infty, 1)$	1	$(1, \infty)$	$f''(x)$	-	0	+	1M					
x	$(-\infty, 1)$	1	$(1, \infty)$											
$f''(x)$	-	0	+											
Thus, the point of inflection of C is $(1, -6)$.	1A													
----- (2) (e) Note that the equation of L is $y = 10$. $\begin{aligned} x^3 - 3x^2 - 9x + 5 &= 10 \\ x^3 - 3x^2 - 9x - 5 &= 0 \\ (x+1)^2(x-5) &= 0 \\ x = -1 \text{ or } x = 5 \end{aligned}$ <p>The required area</p> $\begin{aligned} &= \int_{-1}^5 (10 - (x^3 - 3x^2 - 9x + 5)) dx \\ &= \int_{-1}^5 (-x^3 + 3x^2 + 9x + 5) dx \\ &= \left[\frac{-x^4}{4} + x^3 + \frac{9x^2}{2} + 5x \right]_{-1}^5 \\ &= 108 \end{aligned}$	1M 1M 1M 1A													
		----- (4)												

[HKDSE 2016 M2 #9]

12.

6. (a) Let r cm be the radius of the water surface in the container.

$$\text{Since } \frac{r}{h} = \frac{15}{20}, \text{ we have } \frac{r}{h} = \frac{3}{4}.$$

$$\text{So, we have } r = \frac{3h}{4}.$$

$$\begin{aligned} A &= \pi \left(\frac{3h}{4} \right) \sqrt{h^2 + \left(\frac{3h}{4} \right)^2} \\ &= \pi \left(\frac{3h}{4} \right) \sqrt{\frac{25h^2}{16}} \\ &= \frac{15}{16} \pi h^2 \end{aligned}$$

1M

1M

1

- (b) Let d cm be the depth of water when the volume of water in the container is 96π cm³.

$$\text{Note that } \frac{\pi d}{3} \left(\frac{3d}{4} \right)^2 = 96\pi.$$

$$\text{So, we have } d = 8.$$

$$\text{By (a), we have } A = \frac{15}{16} \pi h^2.$$

$$\text{At time } t \text{ s, we have } \frac{dA}{dt} = \frac{15}{8} \pi h \frac{dh}{dt}.$$

$$\text{Also note that } \frac{dh}{dt} = \frac{3}{\pi}.$$

$$\text{Therefore, we have } \left. \frac{dA}{dt} \right|_{h=8} = \frac{15}{8} \pi (8) \left(\frac{3}{\pi} \right).$$

$$\text{Hence, we have } \left. \frac{dA}{dt} \right|_{h=8} = 45.$$

Thus, the required rate of change is $45 \text{ cm}^2/\text{s}$.

1M

1A

1M

1A

----- (7)

[HKDSE 2017 M2 #6]

13.

9. (a) The equation of the vertical asymptote is $x+4=0$.

$$\text{Note that } f(x) = x - 9 + \frac{36}{x+4}.$$

Thus, the equation of the oblique asymptote is $y = x - 9$.

1A

1M

1A

-----(3)

(b) $f'(x)$

$$= \frac{d}{dx} \left(x - 9 + \frac{36}{x+4} \right)$$

$$= 1 + 36(-1)(x+4)^{-2}$$

$$= 1 - \frac{36}{(x+4)^2}$$

1M

1A

$f'(x)$

$$= \frac{d}{dx} \left(\frac{x^2 - 5x}{x+4} \right)$$

$$= \frac{(x+4)(2x-5) - (x^2 - 5x)}{(x+4)^2}$$

$$= \frac{x^2 + 8x - 20}{(x+4)^2}$$

1M

1A

-----(2)

(c) Note that $f'(x) = \frac{(x+10)(x-2)}{(x+4)^2}$.

So, we have $f'(x) = 0 \Leftrightarrow x = -10$ or $x = 2$.

1A

x	$(-\infty, -10)$	-10	$(-10, -4)$	$(-4, 2)$	2	$(2, \infty)$
$f'(x)$	+	0	-	-	0	+
$f(x)$	\nearrow	-25	\searrow	\searrow	-1	\nearrow

1M

Thus, the maximum point and the minimum point of G are $(-10, -25)$ and $(2, -1)$ respectively.

1A

1A

Note that $f'(x) = \frac{(x+10)(x-2)}{(x+4)^2}$ and $f''(x) = \frac{72}{(x+4)^3}$.

So, we have $f'(x) = 0 \Leftrightarrow x = -10$ or $x = 2$.

1A

Also note that $f''(-10) = \frac{-1}{3} < 0$ and $f''(2) = \frac{1}{3} > 0$.

1M

Further note that $f(-10) = -25$ and $f(2) = -1$.

Thus, the maximum point and the minimum point of G are $(-10, -25)$ and $(2, -1)$ respectively.

1A

1A

-----(4)

(d) The required volume

$$\begin{aligned}
 &= \pi \int_0^5 \left(\frac{x^2 - 5x}{x+4} \right)^2 dx \\
 &= \pi \int_0^5 \left(x - 9 + \frac{36}{x+4} \right)^2 dx \\
 &= \pi \int_0^5 \left(x^2 - 18x + 81 + \frac{72(x-9)}{x+4} + \frac{1296}{(x+4)^2} \right) dx \\
 &= \pi \int_0^5 \left(x^2 - 18x + 153 - \frac{936}{x+4} + \frac{1296}{(x+4)^2} \right) dx \\
 &= \pi \left[\frac{x^3}{3} - 9x^2 + 153x - 936 \ln|x+4| - \frac{1296}{x+4} \right]_0^5 \\
 &= \left(\frac{2285}{3} - 1872 \ln\left(\frac{3}{2}\right) \right) \pi
 \end{aligned}$$

1M

1M

1M

1A

The required volume

$$\begin{aligned}
 &= \pi \int_0^5 \left(\frac{x^2 - 5x}{x+4} \right)^2 dx \\
 &= \pi \int_4^9 \frac{(x-4)^2(x-9)^2}{x^2} dx \\
 &= \pi \int_4^9 \left(\frac{x^4 - 26x^3 + 241x^2 - 936x + 1296}{x^2} \right) dx \\
 &= \pi \int_4^9 \left(x^2 - 26x + 241 - \frac{936}{x} + \frac{1296}{x^2} \right) dx \\
 &= \pi \left[\frac{x^3}{3} - 13x^2 + 241x - 936 \ln|x| - \frac{1296}{x} \right]_4^9 \\
 &= \left(\frac{2285}{3} - 1872 \ln\left(\frac{3}{2}\right) \right) \pi
 \end{aligned}$$

1M

1M

1M

1A

-----(4)

[HKDSE 2017 M2 #9]

14.

8. (a) Note that
- $A \neq 0$
- .

$$f'(x) = \frac{-A(2x-4)}{(x^2 - 4x + 7)^2}$$

So, we have $f'(x) = 0 \Leftrightarrow x = 2$.

Since the equation $f'(x) = 0$ has only one solution $x = 2$ and the extreme value of $f(x)$ is 4, we have $f(2) = 4$.

$$\text{Hence, we have } \frac{A}{2^2 - 4(2) + 7} = 4.$$

Therefore, we have $A = 12$.

$$\text{Thus, we have } f'(x) = \frac{24(2-x)}{(x^2 - 4x + 7)^2}.$$

1M

1M

1A

- (b) Note that
- $x^2 - 4x + 7 = (x-2)^2 + 3 > 0$
- for all real values of
- x
- .

So, there are no vertical asymptotes of the graph of $y = f(x)$.

$$\text{Also note that } f(x) = \frac{12}{x^2 - 4x + 7}.$$

Therefore, $y = 0$ is the only asymptote of the graph of $y = f(x)$.

Hence, there is only one asymptote of the graph of $y = f(x)$.

Thus, the claim is disagreed.

1M

1A f.t.

- (c)
- $f''(x)$

$$= \frac{(x^2 - 4x + 7)^2(-24) - (-24x + 48)(2)(x^2 - 4x + 7)(2x - 4)}{(x^2 - 4x + 7)^4}$$

$$= \frac{72(x-3)(x-1)}{(x^2 - 4x + 7)^3}$$

So, we have $f''(x) = 0 \Leftrightarrow x = 1$ or $x = 3$.

1M

x	$(-\infty, 1)$	1	$(1, 3)$	3	$(3, \infty)$
$f''(x)$	+	0	-	0	+

1M for testing

[HKDSE 2018 M2 #8]

Thus, the points of inflection are $(1, 3)$ and $(3, 3)$.

1A for both correct

-----(8)

15.

9. (a) $y = \frac{\ln x}{2}$
 $\frac{dy}{dx} = \frac{1}{2x}$

The slope of the tangent at P is $\frac{1}{2r}$.

The slope of the normal at P is $-2r$.
 Let a be the x -coordinate of Q .

$$\frac{0 - \ln \sqrt{r}}{a - r} = -2r$$

$$\frac{-1}{2} \ln r = 2r^2 - 2ar$$

$$2ar = 2r^2 + \frac{1}{2} \ln r$$

$$a = \frac{4r^2 + \ln r}{4r}$$

Thus, the x -coordinate of Q is $\frac{4r^2 + \ln r}{4r}$.

1M

1M

1

(3)

(b) Let A square units be the area of $\triangle PQR$.

$$A = \frac{1}{2} \left(\frac{4r^2 + \ln r}{4r} - r \right) \ln \sqrt{r}$$

$$= \frac{(\ln r)^2}{16r}$$

$$\frac{dA}{dr}$$

$$= \frac{r(2\ln r)\frac{1}{r} - (\ln r)^2}{16r^2}$$

$$= \frac{2\ln r - (\ln r)^2}{16r^2}$$

$$= \frac{(2 - \ln r)\ln r}{16r^2}$$

1M

1A

1M

So, we have $\frac{dA}{dr} = 0 \Leftrightarrow \ln r = 2$ or $\ln r = 0$ (rejected).

Hence, we have $\frac{dA}{dr} = 0 \Leftrightarrow r = e^2$.

r	$(1, e^2)$	e^2	(e^2, ∞)
$\frac{dA}{dr}$	+	0	-
A	\nearrow	$\frac{1}{4e^2}$	\searrow

1M for testing

Therefore, A attains its greatest value when $r = e^2$.

Thus, the greatest area of $\triangle PQR$ is $\frac{1}{4e^2}$ square units.

1A

(5)

$ \begin{aligned} (c) \quad & OP \\ &= \sqrt{r^2 + (\ln \sqrt{r})^2} \\ &= \frac{1}{2} \sqrt{4r^2 + (\ln r)^2} \end{aligned} $	1M
$ \begin{aligned} & \frac{dOP}{dt} \\ &= \left(\frac{4r^2 + \ln r}{2r\sqrt{4r^2 + (\ln r)^2}} \right) \left(\frac{dr}{dt} \right) \end{aligned} $	1M
$ \begin{aligned} & \frac{dA}{dt} \\ &= \left(\frac{dA}{dr} \right) \left(\frac{dr}{dt} \right) \\ &= \left(\frac{(2 - \ln r)\ln r}{16r^2} \right) \left(\frac{dr}{dt} \right) \quad (\text{by (b)}) \\ &= \left(\frac{(2 - \ln r)\ln r}{16r^2} \right) \left(\frac{2r\sqrt{4r^2 + (\ln r)^2}}{4r^2 + \ln r} \right) \left(\frac{dOP}{dt} \right) \\ &= \frac{(2 - \ln r)(\ln r)\sqrt{4r^2 + (\ln r)^2}}{8r(4r^2 + \ln r)} \left(\frac{dOP}{dt} \right) \end{aligned} $	1M
$ \begin{aligned} & \left. \frac{dA}{dt} \right _{r=e} \\ &= \frac{(2 - \ln e)(\ln e)\sqrt{4e^2 + (\ln e)^2}}{8e(4e^2 + \ln e)} \left(\left. \frac{dOP}{dt} \right _{r=e} \right) \\ &= \frac{\sqrt{4e^2 + 1}}{8e(4e^2 + 1)} \left(\left. \frac{dOP}{dt} \right _{r=e} \right) \end{aligned} $	1M
<p>Since $0 \leq \left. \frac{dOP}{dt} \right _{r=e} \leq 32e^2$, we have $0 \leq \left. \frac{dA}{dt} \right _{r=e} \leq \frac{32e^2\sqrt{4e^2 + 1}}{8e(4e^2 + 1)}$.</p> <p>So, we have $0 \leq \left. \frac{dA}{dt} \right _{r=e} \leq \frac{4e}{\sqrt{4e^2 + 1}}$.</p> <p>Therefore, we have $0 \leq \left. \frac{dA}{dt} \right _{r=e} < \frac{4e}{\sqrt{4e^2}}$.</p> <p>Hence, we have $0 \leq \left. \frac{dA}{dt} \right _{r=e} < 2$.</p> <p>Thus, the claim is correct.</p>	1A -----(4) f.t.

[HKDSE 2018 M2 #9]

16.

3. (a) $\begin{aligned} V &= \int -2t \, dt \\ &= -t^2 + C, \text{ where } C \text{ is a constant} \\ \text{Since } V &= 580 \text{ when } t = 0, \text{ we have } C = 580. \\ \text{So, we have } V &= 580 - t^2. \\ \text{When } t &= 24, \text{ we have } V = 4 > 0. \\ \text{Thus, the claim is correct.} \end{aligned}$	1M	
	1M	
	1A	f.t.
(b) Let p cm be the depth of liquid X in the vessel when $t = 18$.	1M	
Since $V _{t=18} = 580 - 18^2 = 256$, we have $p^2 + 24p = 256$.	1M	
So, we have $p^2 + 24p - 256 = 0$.	1M	
Solving, we have $p = 8$ or $p = -32$ (rejected).	1A	-0.9
Note that $\frac{dV}{dt} = (2h + 24)\frac{dh}{dt}$.	1M	
Since $\frac{dV}{dt} _{t=18} = -36$, we have $-36 = (2(8) + 24)\frac{dh}{dt} _{t=18}$.	1A	-0.9
Thus, we have $\frac{dh}{dt} _{t=18} = \frac{-9}{10}$.	-----	(6)

[HKDSE 2019 M2 #3]

Technique of Integration

1.

$$4. \quad (a) \quad \int \frac{x+1}{x} dx = \int \left(1 + \frac{1}{x}\right) dx$$

$$= x + \ln|x| + C$$

1M

1A

$$(b) \quad \begin{aligned} \text{Let } u &= x^2 - 1. \\ du &= 2x dx \\ \int \frac{x^3}{x^2 - 1} dx &= \int \frac{u+1}{u} \cdot \frac{du}{2} \\ &= \frac{1}{2}u + \frac{1}{2}\ln|u| + C \quad \text{by (a)} \\ &= \frac{1}{2}(x^2 - 1) + \frac{1}{2}\ln|x^2 - 1| + C \end{aligned}$$

1A

1M

1A

OR $\frac{1}{2}x^2 + \frac{1}{2}\ln|x^2 - 1| + C$

(5)

[HKDSE 2012 M2 #4]

2.

$$5. \quad (a) \quad \int \frac{dx}{\sqrt{9-x}} = \int -(9-x)^{-\frac{1}{2}} d(9-x)$$

1M+1A

Alternative Solution

Let $u = 9-x$.

$du = -dx$

$$\begin{aligned} \int \frac{dx}{\sqrt{9-x}} &= \int -u^{-\frac{1}{2}} du \\ &= -2u^{\frac{1}{2}} + C \end{aligned}$$

1M

1A

$$= -2\sqrt{9-x} + C$$

1A

1M

$$(b) \quad \begin{aligned} \text{Let } x &= 3 \sin \theta. \\ dx &= 3 \cos \theta d\theta \end{aligned}$$

$$\begin{aligned} \int \frac{dx}{\sqrt{9-x^2}} &= \int \frac{3 \cos \theta d\theta}{\sqrt{9-9 \sin^2 \theta}} \\ &= \int d\theta \\ &= \theta + C \\ &= \sin^{-1} \frac{x}{3} + C \end{aligned}$$

1A

1A

1A

1A

(6)

[HKDSE 2014 M2 #5]

3.

3. (a)
$$\int \frac{1}{e^{2u}} du$$

$$= \int e^{-2u} du$$

$$= \frac{-1}{2} e^{-2u} + \text{constant}$$

1M

1A

(b) Let $u = \sqrt{x}$. Then, we have $\frac{du}{dx} = \frac{1}{2\sqrt{x}}$.

1M

$$\begin{aligned} & \int_1^9 \frac{1}{\sqrt{x} e^{2\sqrt{x}}} dx \\ &= \int_1^3 \frac{2}{e^{2u}} du \\ &= 2 \int_1^3 \frac{1}{e^{2u}} du \\ &= 2 \left[\frac{-1}{2} e^{-2u} \right]_1^3 \quad (\text{by (a)}) \\ &= \frac{1}{e^2} - \frac{1}{e^6} \end{aligned}$$

1M+1A

1M

1A

for using the result of (a)

[HKDSE 2015 M2 #3]

----- (7)

4.

$$\begin{aligned}
 7. \quad (a) \quad & \int e^x \sin \pi x \, dx \\
 &= \int \sin \pi x \, de^x \\
 &= e^x \sin \pi x - \pi \int e^x \cos \pi x \, dx \\
 &= e^x \sin \pi x - \pi \int \cos \pi x \, de^x \\
 &= e^x \sin \pi x - \pi \left(e^x \cos \pi x - \pi \int -e^x \sin \pi x \, dx \right) \\
 &\int e^x \sin \pi x \, dx = e^x \sin \pi x - \pi e^x \cos \pi x - \pi^2 \int e^x \sin \pi x \, dx \\
 &\pi^2 \int e^x \sin \pi x \, dx + \int e^x \sin \pi x \, dx = e^x \sin \pi x - \pi e^x \cos \pi x + \text{constant} \\
 &(\pi^2 + 1) \int e^x \sin \pi x \, dx = e^x (\sin \pi x - \pi \cos \pi x) + \text{constant} \\
 &\int e^x \sin \pi x \, dx = \frac{1}{\pi^2 + 1} (e^x (\sin \pi x - \pi \cos \pi x)) + \text{constant}
 \end{aligned}$$

1M

1M

1M

1A

$$\begin{aligned}
 (b) \quad & \int_0^3 e^{3-x} \sin \pi x \, dx \\
 &= \int_3^0 -e^u \sin \pi(3-u) \, du \quad (\text{by letting } u = 3-x) \\
 &= \int_0^3 e^u \sin \pi u \, du \\
 &= \int_0^3 e^x \sin \pi x \, dx \\
 &= \frac{1}{\pi^2 + 1} \left[e^x (\sin \pi x - \pi \cos \pi x) \right]_0^3 \quad (\text{by (a)}) \\
 &= \frac{\pi(e^3 + 1)}{\pi^2 + 1}
 \end{aligned}$$

1M

for using the result of (a)

1A


[HKDSE 2019 M2 #7]

Applications of Integration

1.

9. (a) $\int x \sin x dx = x(-\cos x) - \int (-\cos x) dx$
 $= -x \cos x + \sin x + C$

(b) The volume $= \pi \int_0^{\pi} x \sin x dx$
 $= \pi [-x \cos x + \sin x]_0^{\pi}$
 $= \pi^2$



[HKDSE 2012 M2 #9]

2.

13. (a) (i)
$$\begin{aligned}\tan u &= \frac{-1 + \cos \frac{2\pi}{5}}{\sin \frac{2\pi}{5}} \\ &= \frac{-1 + 1 - 2 \sin^2 \frac{\pi}{5}}{2 \sin \frac{\pi}{5} \cos \frac{\pi}{5}} \\ &= -\tan \frac{\pi}{5} \\ &= \tan \frac{-\pi}{5} \\ \therefore u &= \frac{-\pi}{5} \quad \text{for } -\frac{\pi}{2} < u < \frac{\pi}{2}\end{aligned}$$

1M

1

(ii)
$$\begin{aligned}\tan v &= \frac{1 + \cos \frac{2\pi}{5}}{\sin \frac{2\pi}{5}} \\ &= \frac{1 + 2 \cos^2 \frac{\pi}{5} - 1}{2 \sin \frac{\pi}{5} \cos \frac{\pi}{5}} \\ &= \cot \frac{\pi}{5} \\ &= \tan \left(\frac{\pi}{2} - \frac{\pi}{5} \right) \\ \therefore v &= \frac{3\pi}{10} \quad \text{for } -\frac{\pi}{2} < v < \frac{\pi}{2}\end{aligned}$$

1M

1A

(4)

(b) (i)
$$\begin{aligned}x^2 + 2x \cos \frac{2\pi}{5} + 1 \\ x^2 + 2x \cos \frac{2\pi}{5} + \cos^2 \frac{2\pi}{5} + \sin^2 \frac{2\pi}{5} \\ = (x + \cos \frac{2\pi}{5})^2 + \sin^2 \frac{2\pi}{5}\end{aligned}$$

1A

(ii)
$$\int_{-1}^1 \frac{\sin \frac{2\pi}{5}}{x^2 + 2x \cos \frac{2\pi}{5} + 1} dx = \int_{-1}^1 \frac{\sin \frac{2\pi}{5}}{(x + \cos \frac{2\pi}{5})^2 + \sin^2 \frac{2\pi}{5}} dx$$

Let $x + \cos \frac{2\pi}{5} = \sin \frac{2\pi}{5} \tan \theta$
 $\therefore dx = \sin \frac{2\pi}{5} \sec^2 \theta d\theta$

1M

1A

When $x = -1$, $\tan \theta = \frac{-1 + \cos \frac{2\pi}{5}}{\sin \frac{2\pi}{5}}$ which gives $\theta = \frac{-\pi}{5}$ (by (a)(i))
When $x = 1$, $\tan \theta = \frac{1 + \cos \frac{2\pi}{5}}{\sin \frac{2\pi}{5}}$ which gives $\theta = \frac{3\pi}{10}$ (by (a)(ii))

1M

For using (a)

$$\begin{aligned}\therefore \int_{-1}^1 \frac{\sin \frac{2\pi}{5}}{x^2 + 2x \cos \frac{2\pi}{5} + 1} dx &= \int_{-\frac{\pi}{5}}^{\frac{3\pi}{10}} \frac{\sin^2 \frac{2\pi}{5} \sec^2 \theta}{\frac{5}{5} \sin^2 \frac{2\pi}{5} (\tan^2 \theta + 1)} d\theta \\ &= [\theta]_{-\frac{\pi}{5}}^{\frac{3\pi}{10}} \\ &= \frac{\pi}{2}\end{aligned}$$

1A For integrand

1A

(6)

$$(c) \int_{-1}^1 \frac{\sin \frac{7\pi}{5}}{x^2 + 2x \cos \frac{7\pi}{5} + 1} dx = \int_{-1}^1 \frac{-\sin \frac{2\pi}{5}}{x^2 - 2x \cos \frac{2\pi}{5} + 1} dx$$

1A

1M

Let $y = -x$.

$dy = -dx$

When $x = -1$, $y = 1$; when $x = 1$, $y = -1$.

$$\therefore \int_{-1}^1 \frac{\sin \frac{7\pi}{5}}{x^2 + 2x \cos \frac{7\pi}{5} + 1} dx = \int_1^{-1} \frac{-\sin \frac{2\pi}{5}}{y^2 + 2y \cos \frac{2\pi}{5} + 1} \cdot -dy$$

Alternative Solution

$$\int_{-1}^1 \frac{\sin \frac{7\pi}{5}}{x^2 + 2x \cos \frac{7\pi}{5} + 1} dx = \int_{-1}^1 \frac{\sin \frac{7\pi}{5}}{(x + \cos \frac{7\pi}{5})^2 + \sin^2 \frac{7\pi}{5}} dx$$

1M

Let $x + \cos \frac{7\pi}{5} = \sin \frac{7\pi}{5} \tan \theta$

$$\therefore dx = \sin \frac{7\pi}{5} \sec^2 \theta d\theta$$

When $x = -1$, $\theta = \frac{3\pi}{10}$; when $x = 1$, $\theta = -\frac{\pi}{5}$.

1A

$$\therefore \int_{-1}^1 \frac{\sin \frac{7\pi}{5}}{x^2 + 2x \cos \frac{7\pi}{5} + 1} dx = \int_{\frac{-\pi}{5}}^{\frac{3\pi}{10}} \frac{\sin^2 \frac{7\pi}{5} \sec^2 \theta}{\sin^2 \frac{7\pi}{5} (\tan^2 \theta + 1)} d\theta$$

1A

$$= \frac{-\pi}{2} \text{ by (b)(ii)}$$

(3)

[HKDSE 2012 M2 #13]

3.

14. (a) $y = kx^p$

$$\frac{dy}{dx} = kp x^{p-1}$$

The slope of the tangent to Γ at A is kpa^{p-1} .

$$\therefore \frac{ka^p - 0}{a - (-a)} = kpa^{p-1}$$

$$p = \frac{1}{2}$$

1A

1M

1

(3)

 (b) (i) Let $P(0, t)$ be the centre of C .

$$\therefore AP = 2$$

$$\therefore (k-t)^2 + (1-0)^2 = 2^2$$

$$k-t = -\sqrt{3} \quad [\text{or } \sqrt{3} \text{ (rejected)}]$$

$$\text{Slope of } AP = \frac{k-t}{1-0}$$

$$\text{Slope of } AB = \frac{k}{2} \quad (\text{by (a)})$$

$$\therefore (k-t)\frac{k}{2} = -1 \quad (2)$$

$$\text{Substitute (1) into (2): } (-\sqrt{3})\frac{k}{2} = -1$$

Alternative Solution

$$\text{According to the figure, } \angle ROB = \angle RAP = \frac{\pi}{2}.$$

$$\angle ORB = \angle ARP \quad (\text{vert. opp. } \angle s)$$

$$\therefore \angle RBO = \angle RPA \text{ and let the angles be } \theta.$$

$$\text{Since } PA = 2 \text{ and } QA = 1, \theta = \frac{\pi}{6}$$

$$\text{Slope of } AB = \frac{k}{2} \quad (\text{by (a)})$$

$$\therefore \tan \frac{\pi}{6} = \frac{k}{2}$$




$$k = \frac{2\sqrt{3}}{3}$$

(ii) The shaded area

$$= \text{area of } \triangle PQA + \text{area on the left of } \Gamma \text{ from } O \text{ to } A - \text{area of sector } PAS$$

$$= \frac{1}{2} \cdot 1 \cdot \sqrt{4-1} + \int_0^{\frac{2\sqrt{3}}{3}} \left(\frac{y}{\frac{2\sqrt{3}}{3}} \right)^2 dy - \frac{1}{2} (2)^2 \frac{\pi}{6}$$

$$= \frac{\sqrt{3}}{2} + \frac{3}{4} \left[\frac{y^3}{\frac{2\sqrt{3}}{3}} \right]_0^{\frac{2\sqrt{3}}{3}} - \frac{\pi}{3}$$


Alternative Solution 1

$$t = k + \sqrt{3}$$


$$= \frac{5\sqrt{3}}{3}$$

The shaded area

$$= \text{area of trapezium OFAP} - \text{area of sector PAS} - \text{area under } \Gamma \text{ from } O \text{ to } A$$

$$= \frac{1}{2} \left(\frac{2\sqrt{3}}{3} + \frac{5\sqrt{3}}{3} \right) (1) - \frac{1}{2} (2)^2 \frac{\pi}{6} - \int_0^1 \frac{2\sqrt{3}}{3} x^{\frac{1}{2}} dx$$

$$= \frac{7\sqrt{3}}{6} - \frac{\pi}{3} - \frac{4\sqrt{3}}{9} \left[x^{\frac{3}{2}} \right]_0^1$$



Alternative Solution 2

$$t = k + \sqrt{3}$$

$$= \frac{5\sqrt{3}}{3}$$

The shaded area


= area of ΔOAP + area of ΔOAF – area of sector PAS – area under Γ from O to A

$$= \frac{1}{2} \left(\frac{5\sqrt{3}}{3} \right) (1) + \frac{1}{2} (1) \left(\frac{2\sqrt{3}}{3} \right) - \frac{1}{2} (2)^2 \frac{\pi}{6} - \int_0^1 2\sqrt{3} x^{\frac{1}{2}} dx$$

$$= \frac{5\sqrt{3}}{6} + \frac{\sqrt{3}}{3} - \frac{\pi}{3} - \frac{4\sqrt{3}}{9} \left[x^{\frac{3}{2}} \right]_0^1$$

1M+1A

1M


Alternative Solution 3

$$t = k + \sqrt{3}$$

$$= \frac{5\sqrt{3}}{3}$$

The equation of C is $x^2 + \left(y - \frac{5\sqrt{3}}{3} \right)^2 = 4$.

1A

Hence, the equation of AS is $y = \frac{5\sqrt{3}}{3} - \sqrt{4-x^2}$.

$$\text{The shaded area} = \int_0^1 \left(\frac{5\sqrt{3}}{3} - \sqrt{4-x^2} - \frac{2\sqrt{3}}{3} x^{\frac{1}{2}} \right) dx$$

1M

For $\int_0^1 \sqrt{4-x^2} dx$, let $x = 2\sin\phi$.

$$\therefore dx = 2\cos\phi d\phi$$

$$\text{When } x=1, \phi = \frac{\pi}{6}; \text{ when } x=0, \phi = 0.$$

$$\begin{aligned} \int_0^1 \sqrt{4-x^2} dx &= \int_0^{\frac{\pi}{6}} \sqrt{4-4\sin^2\phi} 2\cos\phi d\phi \\ &= \int_0^{\frac{\pi}{6}} 2(1+\cos 2\phi) d\phi \end{aligned}$$

1M


$$\begin{aligned} &= [2\phi + \sin 2\phi]_0^{\frac{\pi}{6}} \\ &= \frac{\pi}{3} + \frac{\sqrt{3}}{2} \end{aligned}$$

Hence, the shaded area

$$= \left[\frac{5\sqrt{3}}{3}x - \frac{4\sqrt{3}}{9}x^{\frac{3}{2}} \right]_0^1 - \left(\frac{\pi}{3} + \frac{\sqrt{3}}{2} \right)$$

1M

For $A = \int y dx$



$$= \frac{13\sqrt{3}}{18} - \frac{\pi}{3}$$

1A

(9)

[HKDSE 2012 M2 #14]

4.

4. (a) $\frac{dy}{dx} = e^x - 1$

$$y = \int (e^x - 1) dx$$

$$= e^x - x + C$$

Since the curve passes through the point $(1, e)$, $e = e^1 - 1 + C$.

i.e. $C = 1$

$$\therefore y = e^x - x + 1$$


(b) The curve cuts the y -axis at $(0, 2)$.

$$\text{When } x = 0, \frac{dy}{dx} = 0.$$

Hence the equation of tangent to the curve at $(0, 2)$ is

$$y - 2 = 0(x - 0)$$


$$y = 2$$



[HKDSE 2013 M2 #4]

5.

6. (a) Area $= \int_0^4 \left[\left(\frac{-x^2 + 2x + 4}{2} \right) - 4 \right] dx + \int_4^5 \left[4 - \left(\frac{-x^2 + 2x + 4}{2} \right) \right] dx$
 $= \int_0^4 \left(\frac{-x^2 + 2x}{2} \right) dx + \int_4^5 \left(\frac{x^2 - 2x}{2} \right) dx$
 $= \left[\frac{-x^3}{6} + x^2 \right]_0^4 + \left[\frac{x^3}{6} - x^2 \right]_4^5$
 $= \frac{13}{2}$



[HKDSE 2013 M2 #6]

6.

11. (a)
$$\begin{aligned}\frac{d}{d\theta} \ln(\sec \theta + \tan \theta) &= \frac{\sec \theta \tan \theta + \sec^2 \theta}{\sec \theta + \tan \theta} \\ &= \sec \theta\end{aligned}$$

$$\begin{aligned}\text{Hence } \int \sec \theta d\theta &= \int \frac{d}{d\theta} \ln(\sec \theta + \tan \theta) d\theta \\ &= \ln(\sec \theta + \tan \theta) + C\end{aligned}$$

1M

1

Alternative Solution

$$\int \sec \theta d\theta = \int \frac{\sec \theta (\sec \theta + \tan \theta)}{\sec \theta + \tan \theta} d\theta$$

Let $u = \sec \theta + \tan \theta$ which gives $du = (\sec \theta \tan \theta + \sec^2 \theta)d\theta$.

$$\begin{aligned}\therefore \int \sec \theta d\theta &= \int \frac{du}{u} \\ &= \ln|u| + C\end{aligned}$$

$$= \ln(\sec \theta + \tan \theta) + C \quad \text{since } \sec \theta + \tan \theta > 0 \text{ for } 0 < \theta < \frac{\pi}{2}$$

1M

1

(2)

(b) (i) Let $u = \sec \theta$, where $0 < \theta < \frac{\pi}{2}$.

$$\therefore du = \sec \theta \tan \theta d\theta$$

$$\int \frac{du}{\sqrt{u^2 - 1}} = \int \frac{\sec \theta \tan \theta d\theta}{\sqrt{\sec^2 \theta - 1}}$$

$$= \int \sec \theta d\theta \quad \text{since } \tan \theta > 0 \text{ for } 0 < \theta < \frac{\pi}{2}$$

$$= \ln(\sec \theta + \tan \theta) + C \quad \text{by (a)}$$

$$= \ln(\sec \theta + \sqrt{\sec^2 \theta - 1}) + C \quad \text{since } \tan \theta > 0 \text{ for } 0 < \theta < \frac{\pi}{2}$$

1M

1


(ii) $\int_0^1 \frac{2x}{\sqrt{x^4 + 4x^2 + 3}} dx = \int_0^1 \frac{2x}{\sqrt{(x^2 + 2)^2 - 1}} dx$	1M
Let $u = x^2 + 2$ which gives $du = 2x dx$. When $x = 0$, $u = 2$; when $x = 1$, $u = 3$.	
$\therefore \int_0^1 \frac{2x}{\sqrt{x^4 + 4x^2 + 3}} dx = \int_2^3 \frac{du}{\sqrt{u^2 - 1}}$	1M
$= \left[\ln(u + \sqrt{u^2 - 1}) \right]_2^3$ by (i)	
$= \ln(3 + \sqrt{8}) - \ln(2 + \sqrt{3})$	
$= \ln\left(\frac{3+2\sqrt{2}}{2+\sqrt{3}} \cdot \frac{2-\sqrt{3}}{2-\sqrt{3}}\right)$	
$= \ln(6+4\sqrt{2} - 3\sqrt{3} - 2\sqrt{6})$	1
	(5)
$t = \tan \phi$	
$\frac{dt}{d\phi} = \sec^2 \phi$	
$= 1 + \tan^2 \phi$	
$\therefore \frac{d\phi}{dt} = \frac{1}{1+t^2}$	1
$\cos^2 \phi = \frac{1}{\sec^2 \phi}$	
$= \frac{1}{1+t^2}$	1A
$\int_0^{\frac{\pi}{4}} \frac{\tan \phi}{\sqrt{1+2\cos^2 \phi}} d\phi = \int_0^1 \frac{t}{\sqrt{1+\frac{2}{1+t^2}}} \cdot \frac{1}{1+t^2} dt$ where $t = \tan \phi$	1M
$= \int_0^1 \frac{t}{\sqrt{(3+t^2)(1+t^2)}} dt$	
$= \frac{1}{2} \int_0^1 \frac{2t}{\sqrt{t^4 + 4t^2 + 3}} dt$	1M
$= \frac{1}{2} \ln(6+4\sqrt{2} - 3\sqrt{3} - 2\sqrt{6})$	1A
	OR $\ln \sqrt{6+4\sqrt{2} - 3\sqrt{3} - 2\sqrt{6}}$
	(5)

[HKDSE 2013 M2 #11]

7.

6. (a) $\int xe^{-x} dx = x(-e^{-x}) - \int (-e^{-x}) dx$
 $= -xe^{-x} - e^{-x} + C$

(b) $\begin{cases} y = xe^{-x} \\ y = \frac{x}{e} \end{cases}$
 $xe^{-x} = \frac{x}{e}$
 $x(e^{-x} - \frac{1}{e}) = 0$
 $x = 0 \text{ or } 1$
 $\therefore \text{the area} = \int_0^1 \left(xe^{-x} - \frac{x}{e} \right) dx$
 $= \left[-xe^{-x} - e^{-x} - \frac{x^2}{2e} \right]_0^1$
 $= \left(-e^{-1} - e^{-1} - \frac{1}{2e} \right) - (-1)$
 $= 1 - \frac{5}{2e}$



[HKDSE 2014 M2 #6]

8.

13. (a) $1 - \cos 4\theta - 2 \cos 2\theta \sin^2 2\theta$
 $= 2\sin^2 2\theta - 2 \cos 2\theta \sin^2 2\theta$
 $= 2\sin^2 2\theta(1 - \cos 2\theta)$
 $= 2(2\sin \theta \cos \theta)^2(2\sin^2 \theta)$
 $= 16\cos^2 \theta \sin^4 \theta$

1M	For $1 - \cos 4\theta = 2\sin^2 2\theta$ OR $1 - \cos 2\theta = 2\sin^2 \theta$
1	
(2)	

(b) $\int_0^{n\pi} \cos^2 x \sin^4 x dx$
 $= \int_0^{n\pi} \frac{1 - \cos 4x - 2 \cos 2x \sin^2 2x}{16} dx \quad \text{by (a)}$
 $= \frac{1}{16} \int_0^{n\pi} (1 - \cos 4x) dx - \frac{1}{16} \int_0^{n\pi} \sin^2 2x \cdot 2 \cos 2x dx$
 $= \frac{1}{16} \left[x - \frac{\sin 4x}{4} \right]_0^{n\pi} - \frac{1}{16} \int_{x=0}^{n\pi} \sin^2 2x d(2x)$
 $= \frac{1}{16} \left[\left(n\pi - \frac{\sin 4n\pi}{4} \right) - 0 \right] - \frac{1}{16} \left[\frac{\sin^3 2x}{3} \right]_0^{n\pi}$

1M	For $d\sin 2x$
1A	For $\frac{\sin^3 2x}{3}$

Alternative Solution
 $= \frac{1}{16} \int_0^{n\pi} (1 - \cos 4x) dx - \frac{1}{16} \int_0^{n\pi} \sin 4x \sin 2x dx$
 $= \frac{1}{16} \left[x - \frac{\sin 4x}{4} \right]_0^{n\pi} - \frac{1}{16} \int_0^{n\pi} \frac{\cos 2x - \cos 6x}{2} dx$
 $= \frac{1}{16} \left[\left(n\pi - \frac{\sin 4n\pi}{4} \right) - 0 \right] - \frac{1}{32} \left[\frac{\sin 2x}{2} - \frac{\sin 6x}{6} \right]_0^{n\pi}$
 $= \frac{n\pi}{16}$

1M	For $\frac{\cos 2x - \cos 6x}{2}$
1A	For $\frac{\sin 2x}{2} - \frac{\sin 6x}{6}$
1	
(4)	

(c) Let $x = k - u$.
 $\therefore dx = -du$
 When $x = 0, u = k$; when $x = k, u = 0$.

$$\begin{aligned} \int_0^k xf(x) dx &= \int_k^0 (k-u)f(k-u)(-du) \\ &= \int_0^k (k-u)f(u) du \\ &= k \int_0^k f(u) du - \int_0^k uf(u) du \\ &= k \int_0^k f(x) dx - \int_0^k xf(x) dx \\ \therefore 2 \int_0^k xf(x) dx &= k \int_0^k f(x) dx \\ \text{i.e. } \int_0^k xf(x) dx &= \frac{k}{2} \int_0^k f(x) dx \end{aligned}$$

1M

1M+1M
1M for reversing the limits
1M for $f(k-u) = f(u)$

1

(4)

(d) Let $f(x) = \cos^2 x \sin^4 x$
 $f(\pi - x) = \cos^2(\pi - x) \sin^4(\pi - x)$
 $= (-\cos x)^2 (\sin x)^4$
 $= \cos^2 x \sin^4 x$
 $= f(x)$

$f(2\pi - x) = \cos^2(2\pi - x) \sin^4(2\pi - x)$
 $= (\cos x)^2 (-\sin x)^4$
 $= f(x)$

The volume of the solid of revolution

$$\begin{aligned} &= 2\pi \int_{\pi}^{2\pi} x \cos^2 x \sin^4 x dx \\ &= 2\pi \left(\int_0^{2\pi} x \cos^2 x \sin^4 x dx - \int_0^{\pi} x \cos^2 x \sin^4 x dx \right) \\ &= 2\pi \cdot \frac{2\pi}{2} \int_0^{2\pi} \cos^2 x \sin^4 x dx - 2\pi \cdot \frac{\pi}{2} \int_0^{\pi} \cos^2 x \sin^4 x dx \quad \text{by (c)} \\ &= 2\pi^2 \left(\frac{2\pi}{16} \right) - \pi^2 \left(\frac{\pi}{16} \right) \quad \text{by (b)} \\ &= \frac{3\pi^3}{16} \end{aligned}$$


1M

1M

1A

(4)

[HKDSE 2014 M2 #13]



9.

4. (a) $\begin{aligned} & \int x^2 \ln x \, dx \\ &= \frac{1}{3} \int \ln x \, dx^3 \\ &= \frac{1}{3} \left(x^3 \ln x - \int x^3 \, d \ln x \right) \\ &= \frac{1}{3} \left(x^3 \ln x - \int x^2 \, dx \right) \\ &= \frac{1}{3} x^3 \ln x - \frac{1}{9} x^3 + \text{constant} \end{aligned}$	1M 1A 1A	for integration by parts
(b) $\begin{aligned} & y \\ &= \int 9x^2 \ln x \, dx \\ &= 9 \left(\frac{1}{3} x^3 \ln x - \frac{1}{9} x^3 \right) + C \quad (\text{by (a)}) \\ &= 3x^3 \ln x - x^3 + C, \text{ where } C \text{ is a constant} \end{aligned}$	1M 1M	for using the result of (a)
<p>Since Γ passes through the point $(1, 4)$, we have $4 = 3 \ln 1 - 1 + C$.</p> <p>Solving, we have $C = 5$.</p> <p>Thus, the equation of Γ is $y = 3x^3 \ln x - x^3 + 5$.</p>	1M 1A	
		-----(7)

[HKDSE 2015 M2 #4]

10.

12. (a) (i) Solving $\begin{cases} 3x + y - 9 = 0 \\ x^2 - 4y + 8 = 0 \end{cases}$, we have $x^2 + 12x - 28 = 0$.

So, we have $x = 2$ or $x = -14$ (rejected).

Thus, the coordinates of B are $(2, 3)$.

1M

1A

(ii) The required capacity

$$\begin{aligned} &= \int_0^3 \pi \left(\frac{9-y}{3} \right)^2 dy + \int_3^h \pi (4y-8) dy \\ &= \pi \int_0^3 \left(9-2y+\frac{y^2}{9} \right) dy + \pi \int_3^h (4y-8) dy \\ &= \pi \left[9y - y^2 + \frac{y^3}{27} \right]_0^3 + \pi \left[2y^2 - 8y \right]_3^h \\ &= \pi(2h^2 - 8h + 25) \end{aligned}$$

1M+1M+1A

1M either one

1

-----(7)

(b) (i) Putting $x = 6$ in $x^2 - 4y + 8 = 0$, we have $y = 11$.

The required capacity

$$\begin{aligned} &= \pi(2(11)^2 - 8(11) + 25) \quad (\text{by (a)(ii)}) \\ &= 179\pi \text{ cm}^3 \end{aligned}$$

1M

1A

(ii) Let h cm be the depth of water in the cup at time t s.

Also let p cm be the depth of water when the volume of water in the cup is $35\pi \text{ cm}^3$.

Note that the volume of the frustum is $19\pi \text{ cm}^3$.

Since $35\pi > 19\pi$, we have $p > 3$.

By (a)(ii), we have $\pi(2p^2 - 8p + 25) = 35\pi$.

Simplifying, we have $p^2 - 4p - 5 = 0$.

Solving, we have $p = 5$ or $p = -1$ (rejected as $3 < p \leq 11$).

Hence, we have $p = 5$.

Let V cm 3 be the volume of water in the cup at time t s.

For $3 < h \leq 11$, we have $V = \pi(2h^2 - 8h + 25)$ (by (a)(ii)).

So, we have $\frac{dV}{dt} = \pi(4h-8)\frac{dh}{dt}$ for $3 < h \leq 11$.

1M withhold 1M if checking is omitted

1M for $k_1 p^2 + k_2 p + k_3 = 0$

Since $\frac{dV}{dt} \Big|_{h=5} = 24\pi$, we have $24\pi = \pi(4(5)-8)\frac{dh}{dt} \Big|_{h=5}$.

Therefore, we have $\frac{dh}{dt} \Big|_{h=5} = 2$.

1M

1A

Thus, the required rate of change is 2 cm/s .

-----(6)

[HKDSE 2015 M2 #12]

11.

3. (a) Note that the coordinates of Q are $(0, 2e^u)$, where $u > 0$.
The area of ΔOPQ

$$= \frac{u(2e^u)}{2} \\ = ue^u$$

1A

- (b) Let v be the y -coordinate of P .

Since $v = 2e^u$, we have $\frac{dv}{dt} = 2e^u \frac{du}{dt}$.

1M

$$\text{Therefore, we have } \frac{du}{dt} = \frac{1}{2e^u} \left(\frac{dv}{dt} \right).$$

Let A square units be the area of ΔOPQ .

By (a), we have $A = ue^u$.

$$\begin{aligned} & \frac{dA}{dt} \\ &= (e^u + ue^u) \frac{du}{dt} \\ &= \left(\frac{e^u + ue^u}{2e^u} \right) \frac{dv}{dt} \\ &= \left(\frac{1+u}{2} \right) \frac{dv}{dt} \end{aligned}$$

1M+1M

$$\text{So, we have } \frac{dA}{dt} \Big|_{u=4} = \left(\frac{1+4}{2} \right)(6) = 15.$$

1A

Thus, the required rate of change is 15 square units per second.

Let v be the y -coordinate of P .

$$\text{Since } v = 2e^u, \text{ we have } u = \ln\left(\frac{v}{2}\right).$$

1M

Let A square units be the area of ΔOPQ .

$$\text{By (a), we have } A = \frac{v}{2} \ln\left(\frac{v}{2}\right).$$

$$\begin{aligned} & \frac{dA}{dt} \\ &= \left(\left(\frac{v}{2} \right) \left(\frac{2}{v} \right) \left(\frac{1}{2} \right) + \frac{1}{2} \ln\left(\frac{v}{2}\right) \right) \frac{dv}{dt} \\ &= \left(\frac{1}{2} + \frac{1}{2} \ln\left(\frac{v}{2}\right) \right) \frac{dv}{dt} \end{aligned}$$

1M+1M

When $u = 4$, we have $v = 2e^4$.

$$\begin{aligned} & \frac{dA}{dt} \Big|_{u=4} \\ &= \left(\frac{1}{2} + \frac{1}{2} \ln\left(\frac{2e^4}{2}\right) \right)(6) \\ &= 15 \end{aligned}$$

1A

Thus, the required rate of change is 15 square units per second.

[HKDSE 2016 M2 #3]

12.

 7. (a) Let $u = \sqrt{t+1}$.

 So, we have $2u \frac{du}{dt} = 1$.

$$\begin{aligned} & \int (1+\sqrt{t+1})^2 dt \\ &= \int (1+u)^2 2u du \\ &= \int (2u + 4u^2 + 2u^3) du \\ &= u^2 + \frac{4}{3}u^3 + \frac{1}{2}u^4 + \text{constant} \\ &= (t+1) + \frac{4}{3}(t+1)^{\frac{3}{2}} + \frac{1}{2}(t+1)^2 + \text{constant} \\ &= 2t + \frac{t^2}{2} + \frac{4}{3}(t+1)^{\frac{3}{2}} + \text{constant} \end{aligned}$$

1M

1M

1A

$$\begin{aligned} & \int (1+\sqrt{t+1})^2 dt \\ &= \int (1+2\sqrt{t+1}+t+1) dt \\ &= \int (2+t+2\sqrt{t+1}) dt \\ &= 2t + \frac{t^2}{2} + \int 2\sqrt{t+1} dt \\ &= 2t + \frac{t^2}{2} + \int 2\sqrt{u} du \quad (\text{by letting } u=t+1) \\ &= 2t + \frac{t^2}{2} + \frac{4}{3}(t+1)^{\frac{3}{2}} + \text{constant} \end{aligned}$$

1M

1M

1A

 (b) Note that $y = 4x^2 - 4x$, where $1 \leq x \leq 4$.

 So, we have $x = \frac{1}{2}(1 + \sqrt{y+1})$.

1M

The required volume

$$\begin{aligned} &= \int_0^{48} \pi \left(\frac{1}{2}(1 + \sqrt{y+1}) \right)^2 dy \\ &= \frac{\pi}{4} \int_0^{48} (1 + \sqrt{y+1})^2 dy \\ &= \frac{\pi}{4} \int_0^{48} (1 + \sqrt{t+1})^2 dt \\ &= \frac{\pi}{4} \left[2t + \frac{t^2}{2} + \frac{4}{3}(t+1)^{\frac{3}{2}} \right]_0^{48} \quad (\text{by (a)}) \\ &= 426\pi \end{aligned}$$

1M

1M

for using (a)

1A

[HKDSE 2016 M2 #7]

13.

10. (a) Let
- $x = a - y$
- .

$$\text{So, we have } \frac{dx}{dy} = -1.$$

$$\int_0^a f(x) dx$$

$$= - \int_a^0 f(a-y) dy$$

$$= \int_0^a f(a-y) dy$$

$$= \int_0^a f(a-x) dx$$

1M

1M

1

-----(3)

$$(b) \quad \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx$$

$$= \int_0^{\frac{\pi}{4}} \ln\left(1 + \tan\left(\frac{\pi}{4} - x\right)\right) dx \quad (\text{by (a) with } a = \frac{\pi}{4} > 0)$$

$$= \int_0^{\frac{\pi}{4}} \ln\left(1 + \frac{1 - \tan x}{1 + \tan x}\right) dx$$

$$= \int_0^{\frac{\pi}{4}} \ln\left(\frac{2}{1 + \tan x}\right) dx$$

1M for using (a)

1M

1

-----(3)

$$(c) \quad \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \int_0^{\frac{\pi}{4}} \ln\left(\frac{2}{1 + \tan x}\right) dx \quad (\text{by (b)})$$

$$\int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \int_0^{\frac{\pi}{4}} (\ln 2 - \ln(1 + \tan x)) dx$$

$$\int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \int_0^{\frac{\pi}{4}} \ln 2 dx - \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx$$

$$2 \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \int_0^{\frac{\pi}{4}} \ln 2 dx$$

$$\int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \frac{1}{2} \int_0^{\frac{\pi}{4}} \ln 2 dx$$

$$\int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx = \frac{\pi \ln 2}{8}$$

1M

1M

1

-----(3)

$$(d) \quad \int_0^{\frac{\pi}{4}} \frac{x \sec^2 x}{1 + \tan x} dx$$

$$= [x \ln(1 + \tan x)]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \ln(1 + \tan x) dx$$

$$= \left(\frac{\pi \ln 2}{4} - 0\right) - \frac{\pi \ln 2}{8} \quad (\text{by (c)})$$

$$= \frac{\pi \ln 2}{8}$$

1M

1M for using (c)

1A

-----(3)

[HKDSE 2016 M2 #10]

14.

4. (a) $ \begin{aligned} & \int x^2 e^{-x} dx \\ &= - \int x^2 de^{-x} \\ &= -x^2 e^{-x} + \int e^{-x} dx^2 \\ &= -x^2 e^{-x} + 2 \int x e^{-x} dx \\ &= -x^2 e^{-x} - 2 \int x de^{-x} \\ &= -x^2 e^{-x} - 2(x e^{-x} - \int e^{-x} dx) \\ &= -x^2 e^{-x} - 2x e^{-x} - 2e^{-x} + \text{constant} \\ &= -e^{-x}(x^2 + 2x + 2) + \text{constant} \end{aligned} $	1M 1A 1A	for integration by parts
(b) The required area $ \begin{aligned} &= \int_0^6 x^2 e^{-x} dx \\ &= \left[-e^{-x}(x^2 + 2x + 2) \right]_0^6 \quad (\text{by (a)}) \\ &= 2 - \frac{50}{e^6} \end{aligned} $	1M 1M 1A	for using the result of (a) -----(6)

[HKDSE 2017 M2 #4]

15.

8. (a) The slope of the tangent to
- Γ
- at
- P

$$= f'(e^3)$$

$$= \frac{1}{e^3} \ln(e^3)^2$$

$$= \frac{6}{e^3}$$

1M

The equation of the tangent to Γ at P is

$$y - 7 = \frac{6}{e^3}(x - e^3)$$

$$6x - e^3y + e^3 = 0$$

1A

(b) $f(x)$

$$= \int \frac{1}{x} \ln x^2 dx$$

$$= 2 \int \ln x d \ln x$$

$$= (\ln x)^2 + C$$

Since Γ passes through P , we have $7 = (\ln e^3)^2 + C$.

Solving, we have $C = -2$.

Thus, the equation of Γ is $y = (\ln x)^2 - 2$.

1M

1A

(c) Note that $f''(x) = \frac{2 - 2 \ln x}{x^2}$.

Therefore, we have $f''(x) = 0 \Leftrightarrow x = e$.

1A

x	$(0, e)$	e	(e, ∞)
$f''(x)$	+	0	-

1M

Thus, the point of inflection of Γ is $(e, -1)$.

1A

----- (8)

[HKDSE 2017 M2 #8]

16.

11. (a) $\begin{aligned} & \int_0^1 \frac{1}{x^2 + 2x + 3} dx \\ &= \int_0^1 \frac{1}{(x+1)^2 + 2} dx \\ &= \frac{1}{\sqrt{2}} \left[\tan^{-1} \left(\frac{x+1}{\sqrt{2}} \right) \right]_0^1 \\ &= \frac{\sqrt{2}}{2} \left(\tan^{-1} \sqrt{2} - \tan^{-1} \left(\frac{\sqrt{2}}{2} \right) \right) \\ &= \frac{\sqrt{2}}{2} \tan^{-1} \left(\frac{\sqrt{2}}{4} \right) \end{aligned}$	1M 1M 1A -----(3)
(b) (i) $\begin{aligned} & \frac{2 \tan \theta}{1 + \tan^2 \theta} \\ &= \frac{2 \sin \theta}{\cos \theta} \\ &= \frac{1 - \frac{\sin^2 \theta}{\cos^2 \theta}}{1 + \frac{\sin^2 \theta}{\cos^2 \theta}} \\ &= 2 \sin \theta \cos \theta \\ &= \sin 2\theta \end{aligned}$	1
$\begin{aligned} & \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \\ &= \frac{1 - \frac{\sin^2 \theta}{\cos^2 \theta}}{1 + \frac{\sin^2 \theta}{\cos^2 \theta}} \\ &= \cos^2 \theta - \sin^2 \theta \\ &= \cos 2\theta \end{aligned}$	1
(ii) Let $t = \tan \theta$. Then, we have $\frac{d\theta}{dt} = \frac{1}{1+t^2}$.	1M
Note that $\frac{1}{\sin 2\theta + \cos 2\theta + 2} = \frac{1}{2t + \frac{1-t^2}{1+t^2} + 2} = \frac{1+t^2}{t^2 + 2t + 3}$.	
$\begin{aligned} & \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= \int_0^1 \frac{1+t^2}{t^2 + 2t + 3} \left(\frac{1}{1+t^2} \right) dt \\ &= \int_0^1 \frac{1}{t^2 + 2t + 3} dt \\ &= \int_0^1 \frac{1}{x^2 + 2x + 3} dx \\ &= \frac{\sqrt{2}}{2} \tan^{-1} \left(\frac{\sqrt{2}}{4} \right) \quad (\text{by (a)}) \end{aligned}$	1M 1M 1M 1M (a) -----(5)

(c) Let $y = \frac{\pi}{4} - \theta$. Then, we have $\frac{d\theta}{dy} = -1$.

$$\begin{aligned} & \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= - \int_{\frac{\pi}{4}}^0 \frac{\sin\left(\frac{\pi}{2} - 2y\right) + 1}{\sin\left(\frac{\pi}{2} - 2y\right) + \cos\left(\frac{\pi}{2} - 2y\right) + 2} dy \\ &= \int_0^{\frac{\pi}{4}} \frac{\cos 2y + 1}{\cos 2y + \sin 2y + 2} dy \\ &= \int_0^{\frac{\pi}{4}} \frac{\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta \end{aligned}$$

1M

1

-----(2)

$$\begin{aligned} (d) & \int_0^{\frac{\pi}{4}} \frac{8\sin 2\theta + 9}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= \int_0^{\frac{\pi}{4}} \frac{4(\sin 2\theta + 1) + 4(\sin 2\theta + 1) + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= 4 \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta + 4 \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta + \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= 4 \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta + 4 \int_0^{\frac{\pi}{4}} \frac{\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta + \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \quad (\text{by (c)}) \\ &= 4 \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + \cos 2\theta + 2}{\sin 2\theta + \cos 2\theta + 2} d\theta + \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= 4 \int_0^{\frac{\pi}{4}} d\theta + \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= \pi + \frac{\sqrt{2}}{2} \tan^{-1}\left(\frac{\sqrt{2}}{4}\right) \quad (\text{by (b)(ii)}) \end{aligned}$$

1M

for using (c)

1M

$\pi + (b)(ii)$

Let $I = \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta$ and $J = \int_0^{\frac{\pi}{4}} \frac{\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta$.

Note that $I + J = \int_0^{\frac{\pi}{4}} d\theta = \frac{\pi}{4}$.

By (c), we have $I = J = \frac{\pi}{8}$.

$$\begin{aligned} & \int_0^{\frac{\pi}{4}} \frac{8\sin 2\theta + 9}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= 8I + \int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\ &= \pi + \frac{\sqrt{2}}{2} \tan^{-1}\left(\frac{\sqrt{2}}{4}\right) \quad (\text{by (b)(ii)}) \end{aligned}$$

1M

for using (c)

1M

$\pi + (b)(ii)$

-----(3)

[HKDSE 2017 M2 #11]

17.

4. (a) $\int u(5^u) du$ $= \frac{1}{\ln 5} \left(u(5^u) - \int 5^u du \right)$ $= \frac{1}{\ln 5} \left(u(5^u) - \frac{5^u}{\ln 5} \right) + \text{constant}$ $= \frac{5^u(u \ln 5 - 1)}{(\ln 5)^2} + \text{constant}$	1M	
(b) The required area $= \int_0^1 x(5^{2x}) dx$ $= \frac{1}{4} \int_0^2 u(5^u) du \quad (\text{by letting } u = 2x)$ $= \frac{1}{4(\ln 5)^2} \left[5^u(u \ln 5 - 1) \right]_0^2 \quad (\text{by (a)})$ $= \frac{50 \ln 5 - 24}{4(\ln 5)^2}$ $= \frac{25 \ln 5 - 12}{2(\ln 5)^2}$	1M	for using the result of (a)
	1A	-----(6)

[HKDSE 2018 M2 #4]

18.

5. (a) Let $u = 1 + x^2$.
Then, we have $\frac{du}{dx} = 2x$.

$$\begin{aligned} & \int x^3 \sqrt{1+x^2} dx \\ &= \int \frac{1}{2}(u-1) u^{\frac{1}{2}} du \\ &= \frac{1}{2} \left(\int u^{\frac{3}{2}} du - \int u^{\frac{1}{2}} du \right) \\ &= \frac{1}{5} \left(\sqrt{1+x^2} \right)^5 - \frac{1}{3} \left(\sqrt{1+x^2} \right)^3 + \text{constant} \end{aligned}$$

1M

1M

1A

Let $x = \tan \theta$.
Then, we have $\frac{dx}{d\theta} = \sec^2 \theta$.

$$\begin{aligned} & \int x^3 \sqrt{1+x^2} dx \\ &= \int \tan^3 \theta \sec \theta (\sec^2 \theta) d\theta \\ &= \int \tan^3 \theta \sec^3 \theta d\theta \\ &= \int (\sec^2 \theta - 1) \sec^2 \theta d\sec \theta \\ &= \int \sec^4 \theta d\sec \theta - \int \sec^2 \theta d\sec \theta \\ &= \frac{\sec^5 \theta}{5} - \frac{\sec^3 \theta}{3} + \text{constant} \\ &= \frac{1}{5} \left(\sqrt{1+x^2} \right)^5 - \frac{1}{3} \left(\sqrt{1+x^2} \right)^3 + \text{constant} \end{aligned}$$

1M

1M

1A

(b)

$$\begin{aligned} & y \\ &= \int 15x^3 \sqrt{1+x^2} dx \\ &= 15 \int x^3 \sqrt{1+x^2} dx \\ &= 15 \left(\frac{1}{5} \left(\sqrt{1+x^2} \right)^5 - \frac{1}{3} \left(\sqrt{1+x^2} \right)^3 \right) + C \quad (\text{by (a)}) \\ &= 3 \left(\sqrt{1+x^2} \right)^5 - 5 \left(\sqrt{1+x^2} \right)^3 + C, \text{ where } C \text{ is a constant} \end{aligned}$$

1M

for using the result of (a)

Since the y -intercept of Γ is 2, we have $3-5+C=2$.
Solving, we have $C=4$.

1M

Thus, the equation of Γ is $y=3\left(\sqrt{1+x^2}\right)^5-5\left(\sqrt{1+x^2}\right)^3+4$.

1A

[HKDSE 2018 M2 #5]

19.

10. (a) (i) $\begin{aligned} & \int \sin^4 x \, dx \\ &= -\cos x \sin^3 x + \int \cos x (3 \sin^2 x \cos x) \, dx \\ &= -\cos x \sin^3 x + 3 \int (1 - \sin^2 x)(\sin^2 x) \, dx \\ &\text{So, we have } \int \sin^4 x \, dx = -\cos x \sin^3 x + 3 \int \sin^2 x \, dx - 3 \int \sin^4 x \, dx. \\ &\text{Hence, we have } 4 \int \sin^4 x \, dx = -\cos x \sin^3 x + 3 \int \sin^2 x \, dx. \\ &\text{Thus, we have } \int \sin^4 x \, dx = \frac{-\cos x \sin^3 x}{4} + \frac{3}{4} \int \sin^2 x \, dx. \end{aligned}$	1M	1
(ii) $\begin{aligned} & \int \sin^4 x \, dx \\ &= \frac{-\cos x \sin^3 x}{4} + \frac{3}{4} \int \sin^2 x \, dx \quad (\text{by (a)(i)}) \\ &= \frac{-\cos x \sin^3 x}{4} + \frac{3}{4} \int \frac{1 - \cos 2x}{2} \, dx \\ &= \frac{-\cos x \sin^3 x}{4} + \frac{3}{4} \left(\frac{x}{2} - \frac{\sin 2x}{4} \right) + \text{constant} \\ &= \frac{3x}{8} - \frac{\cos x \sin^3 x}{4} - \frac{3 \sin 2x}{16} + \text{constant} \\ &\int_0^\pi \sin^4 x \, dx \\ &= \left[\frac{3x}{8} - \frac{\cos x \sin^3 x}{4} - \frac{3 \sin 2x}{16} \right]_0^\pi \\ &= \frac{3\pi}{8} \end{aligned}$	1M	1M
$\begin{aligned} & \int_0^\pi \sin^4 x \, dx \\ &= \int_0^\pi \left(\frac{1 - \cos 2x}{2} \right)^2 \, dx \\ &= \frac{1}{4} \int_0^\pi (1 - 2 \cos 2x + \cos^2 2x) \, dx \\ &= \frac{1}{4} \int_0^\pi \left(1 - 2 \cos 2x + \frac{1 + \cos 4x}{2} \right) \, dx \\ &= \frac{1}{8} \int_0^\pi (3 - 4 \cos 2x + \cos 4x) \, dx \\ &= \frac{1}{8} \left[3x - 2 \sin 2x + \frac{\sin 4x}{4} \right]_0^\pi \\ &= \frac{3\pi}{8} \end{aligned}$	1M	1A
		-----(5)

(b) (i) Let $x = \beta - u$. Then, we have $\frac{dx}{du} = -1$.

$$\begin{aligned} & \int_0^\beta xf(x) dx \\ &= \int_\beta^0 -(\beta - u)f(\beta - u) du \\ &= \int_0^\beta (\beta f(\beta - u) - u f(\beta - u)) du \\ &= \int_0^\beta \beta f(x) dx - \int_0^\beta xf(x) dx \\ \text{So, we have } & 2 \int_0^\beta xf(x) dx = \beta \int_0^\beta f(x) dx. \\ \text{Thus, we have } & \int_0^\beta xf(x) dx = \frac{\beta}{2} \int_0^\beta f(x) dx. \end{aligned}$$

1M

1

(ii) Note that $\sin^4(\pi - x) = \sin^4 x$ for all real numbers x .

$$\begin{aligned} & \int_0^\pi x \sin^4 x dx \\ &= \frac{\pi}{2} \int_0^\pi \sin^4 x dx \quad (\text{by (b)(i)}) \\ &= \frac{\pi}{2} \left(\frac{3\pi}{8} \right) \quad (\text{by (a)(ii)}) \\ &= \frac{3\pi^2}{16} \end{aligned}$$

1M withhold 1M if checking is skipped

1M for using the result of (b)(i)

 1M for $\frac{\pi}{2}$ (a)(ii)

(c) The required volume

$$\begin{aligned} &= \int_{\pi}^{2\pi} \pi(\sqrt{x} \sin^2 x)^2 dx \\ &= \pi \int_{\pi}^{2\pi} x \sin^4 x dx \\ &= \pi \int_0^{\pi} (\pi + y) \sin^4(\pi + y) dy \quad (\text{by letting } x = \pi + y) \\ &= \pi \int_0^{\pi} (\pi \sin^4 y + y \sin^4 y) dy \\ &= \pi \int_0^{\pi} (\pi \sin^4 x + x \sin^4 x) dx \\ &= \pi \left(\pi \int_0^{\pi} \sin^4 x dx + \int_0^{\pi} x \sin^4 x dx \right) \\ &= \pi \left(\pi \left(\frac{3\pi}{8} \right) + \frac{3\pi^2}{16} \right) \quad (\text{by (a)(ii) and (b)(ii)}) \\ &= \frac{9\pi^3}{16} \end{aligned}$$

1M

 1M accept $x = 2\pi - y$

1A

[HKDSE 2018 M2 #10]

20.

$$4. \quad (a) \quad g'(x) = \frac{\sqrt{x} \left(\frac{1}{x}\right) - (\ln x) \left(\frac{1}{2\sqrt{x}}\right)}{x} = \frac{2 - \ln x}{2\sqrt{x^3}}$$

So, we have $g'(x) = 0 \Leftrightarrow x = e^2$.

x	$(0, e^2)$	e^2	$(e^2, 99)$
$g'(x)$	+	0	-

1M for quotient rule

Therefore, G attains its maximum value only at $x = e^2$.
Thus, G has only one maximum point.

1M

1

- (b) Note that $g(x) < 0$ for all $x \in (0, 1)$ and $g(x) > 0$ for all $x \in (1, 99)$.
So, we have $g(x) = 0 \Leftrightarrow x = 1$.

The required volume

$$\begin{aligned} &= \int_1^{e^2} \pi \left(\frac{\ln x}{\sqrt{x}} \right)^2 dx \\ &= \pi \int_0^2 u^2 du \quad (\text{by letting } u = \ln x) \\ &= \pi \left[\frac{u^3}{3} \right]_0^2 \\ &= \frac{8\pi}{3} \end{aligned}$$

1M

1M

1A

----- (6)

[HKDSE 2019 M2 #4]

21.

8. (a) For all $x > 0$,

$$h'(x)$$

$$= \frac{2}{x} \left(\left(x^2 - \frac{7}{2}x + \left(\frac{7}{4} \right)^2 - \left(\frac{7}{4} \right)^2 \right) + 4 \right)$$

$$= \frac{2}{x} \left(\left(x - \frac{7}{4} \right)^2 + \frac{15}{16} \right)$$

$$> 0$$

Thus, $h(x)$ is an increasing function.

1M

1A f.t.

(b) (i)

$$y = \int \left(2x - 7 + \frac{8}{x} \right) dx$$

$$= x^2 - 7x + 8 \ln x + C, \text{ where } C \text{ is a constant}$$

Note that $y = 3$ when $x = 1$.

$$\text{So, we have } 1^2 - 7(1) + 8 \ln 1 + C = 3 .$$

$$\text{Therefore, we have } C = 9 .$$

Thus, the equation of H is $y = x^2 - 7x + 8 \ln x + 9 .$

1M

1A

 (ii) $h''(x)$

$$= \frac{x(4x-7) - (2x^2 - 7x + 8)}{x^2}$$

$$= \frac{2x^2 - 8}{x^2}$$

$$= \frac{2(x-2)(x+2)}{x^2}$$

Since $x > 0$, we have $h''(x) = 0 \Leftrightarrow x = 2 .$

1M

x	$(0, 2)$	2	$(2, \infty)$
$h''(x)$	-	0	+

1M

Thus, the point of inflection of H is $(2, 8 \ln 2 - 1) .$

1A

----- (8)

[HKDSE 2019 M2 #8]

22.

9. (a) $\begin{aligned} \frac{dy}{dx} &= \left(\frac{1}{3}\right)\left(\frac{1}{2}\right)(12-x^2)^{\frac{-1}{2}}(-2x) \\ &= \frac{-x}{3\sqrt{12-x^2}} \\ \text{So, we have } \frac{dy}{dx} \Big _{x=3} &= \frac{-1}{\sqrt{3}}. \end{aligned}$ <p>The equation of L is</p> $y - \frac{\sqrt{3}}{3} = \frac{-1}{\sqrt{3}}(x-3)$ $x + \sqrt{3}y - 4 = 0$	1M	for chain rule
	1M	
	1A	
	-----(3)	
(b) (i) Putting $y = \frac{1}{\sqrt{3}}(4-x)$ in $y = \sqrt{4-x^2}$, we have $\frac{1}{\sqrt{3}}(4-x) = \sqrt{4-x^2}$ $x^2 - 2x + 1 = 0$ So, we have $x = 1$ and $y = \sqrt{3}$. Thus, the point of contact of L and C is $(1, \sqrt{3})$.	1M	1A
(ii) When $\sqrt{4-x^2} = \frac{1}{3}\sqrt{12-x^2}$, we have $36-9x^2 = 12-x^2$. So, we have $8x^2 - 24 = 0$. Since $0 < x < 2$, we have $x = \sqrt{3}$. Thus, the point of intersection of C and I' is $(\sqrt{3}, 1)$.	1M	1A
(iii) The required area $\begin{aligned} &= \int_1^{\sqrt{3}} \left(\frac{1}{\sqrt{3}}(4-x) - \sqrt{4-x^2} \right) dx + \int_{\sqrt{3}}^3 \left(\frac{1}{\sqrt{3}}(4-x) - \frac{1}{3}\sqrt{12-x^2} \right) dx \\ &= \int_1^3 \frac{1}{\sqrt{3}}(4-x) dx - \int_1^{\sqrt{3}} \sqrt{4-u^2} du - \int_{\sqrt{3}}^3 \frac{1}{3}\sqrt{12-v^2} dv \\ &= \frac{4\sqrt{3}}{3} - \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} 4\cos^2\alpha d\alpha - \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{12\cos^2\beta}{3} d\beta \quad (\text{by letting } u = 2\sin\alpha \text{ and } v = \sqrt{12}\sin\beta) \\ &= \frac{4\sqrt{3}}{3} - 8 \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\cos 2\theta + 1}{2} d\theta \\ &= \frac{4\sqrt{3}}{3} - 8 \left[\frac{\sin 2\theta}{4} + \frac{\theta}{2} \right]_{\frac{\pi}{6}}^{\frac{\pi}{3}} \\ &= \frac{4\sqrt{3}}{3} - \frac{2\pi}{3} \end{aligned}$	1M+1A	1M for either integral for either substitution 1M
	-----(9)	

[HKDSE 2019 M2 #9]

23.

$$\begin{aligned}
 10. \quad (a) \quad & \frac{1}{2 + \cos 2x} \\
 &= \frac{1}{2 + 2\cos^2 x - 1} \\
 &= \frac{1}{2\cos^2 x + 1} \\
 &= \frac{\sec^2 x}{2 + \sec^2 x}
 \end{aligned}$$

 1
-----(1)

$$\begin{aligned}
 (b) \quad & \int_0^{\frac{\pi}{4}} \frac{1}{2 + \cos 2x} dx \\
 &= \int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{2 + \sec^2 x} dx \quad (\text{by (a)}) \\
 &= \int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{3 + \tan^2 x} dx \\
 &= \int_0^1 \frac{1}{3 + t^2} dt \quad (\text{by letting } t = \tan x) \\
 &= \left[\frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{t}{\sqrt{3}} \right) \right]_0^1 \\
 &= \frac{\sqrt{3}\pi}{18}
 \end{aligned}$$

1M for using (a)

1M

1A

-----(3)

 (c) By letting $u = -x$, we have

$$\int_{-a}^0 f(x) \ln(1 + e^x) dx = \int_a^0 -f(-u) \ln(1 + e^{-u}) du .$$

1M

$$\text{So, we have } \int_{-a}^0 f(x) \ln(1 + e^x) dx = \int_0^a f(-x) \ln(1 + e^{-x}) dx .$$

$$\begin{aligned}
 & \int_{-a}^a f(x) \ln(1 + e^x) dx \\
 &= \int_{-a}^0 f(x) \ln(1 + e^x) dx + \int_0^a f(x) \ln(1 + e^x) dx \\
 &= \int_0^a f(-x) \ln(1 + e^{-x}) dx + \int_0^a f(x) \ln(1 + e^x) dx \\
 &= - \int_0^a f(x) \ln(1 + e^{-x}) dx + \int_0^a f(x) \ln(1 + e^x) dx \\
 &= - \int_0^a f(x) \ln \left(\frac{e^x + 1}{e^x} \right) dx + \int_0^a f(x) \ln(1 + e^x) dx \\
 &= - \int_0^a f(x) \ln(e^x + 1) dx + \int_0^a x f(x) dx + \int_0^a f(x) \ln(1 + e^x) dx \\
 &= \int_0^a x f(x) dx
 \end{aligned}$$

1M

1M

1M

1

-----(4)

<p>(d) Note that $\frac{\sin 2(-x)}{(2 + \cos 2(-x))^2} = \frac{-\sin 2x}{(2 + \cos 2x)^2}$ for all $x \in \mathbf{R}$.</p> <p>By (c), we have $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\sin 2x}{(2 + \cos 2x)^2} \ln(1 + e^x) dx = \int_0^{\frac{\pi}{4}} \frac{x \sin 2x}{(2 + \cos 2x)^2} dx$.</p> <p>Also note that $\frac{d}{dx} \left(\frac{1}{2(2 + \cos 2x)} \right) = \frac{\sin 2x}{(2 + \cos 2x)^2}$.</p> $\begin{aligned} & \int_0^{\frac{\pi}{4}} \frac{x \sin 2x}{(2 + \cos 2x)^2} dx \\ &= \left[\frac{x}{2(2 + \cos 2x)} \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \frac{1}{2(2 + \cos 2x)} dx \\ &= \frac{1}{2} \left(\frac{\pi}{4} \right) \left(\frac{1}{2+0} \right) - \frac{1}{2} \left(\frac{\sqrt{3}\pi}{18} \right) \quad (\text{by (b)}) \\ &= \frac{(9-4\sqrt{3})\pi}{144} \end{aligned}$ <p>Thus, we have $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\sin 2x}{(2 + \cos 2x)^2} \ln(1 + e^x) dx = \frac{(9-4\sqrt{3})\pi}{144}$.</p>	1M	withhold 1M if checking is omitted
	1M	for using (c)
	1M	for using the result of (b)
	1A	-----(5)

[HKDSE 2019 M2 #10]

Matrices

1.

$$11. \quad (a) \quad \begin{vmatrix} 1-x & 4 \\ 2 & 3-x \end{vmatrix} = 0$$

$$(1-x)(3-x) - 2 \cdot 4 = 0$$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5$$

	1M
	1A
	(2)

$$(b) \quad (i) \quad \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = -1 \cdot \begin{pmatrix} a \\ b \end{pmatrix}$$

$$\begin{cases} a + 4b = -a \\ 2a + 3b = -b \end{cases}$$

$$a + 2b = 0 \quad \dots \quad (1)$$

$$\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} c \\ 1 \end{pmatrix} = 5 \begin{pmatrix} c \\ 1 \end{pmatrix}$$

$$\begin{cases} c + 4 = 5c \\ 2c + 3 = 5 \end{cases}$$


$$c = 1 \quad \dots \quad (2)$$

$$\begin{vmatrix} a & c \\ b & 1 \end{vmatrix} = 1$$

By (2), $a - b = 1 \quad \dots \quad (3)$

Solving (1) and (3), we have $a = \frac{2}{3}$ and $b = \frac{-1}{3}$.

$$\therefore P = \begin{pmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & 1 \end{pmatrix}$$



$$(ii) \quad P^{-1} = \frac{1}{\frac{2}{3} + \frac{1}{3}} \begin{pmatrix} 1 & \frac{1}{3} \\ -1 & \frac{2}{3} \end{pmatrix}^t$$

$$= \begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix}$$

$$P^{-1} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} P = \begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & 1 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 1 \\ \frac{5}{3} & \frac{10}{3} \end{pmatrix} \begin{pmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & 1 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix}$$

$$1M$$

$$1M \quad \left| \begin{array}{l} \\ \\ \end{array} \right. \quad \text{OR} \quad \begin{pmatrix} 1 & -1 \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix} \begin{pmatrix} -2 & 5 \\ \frac{1}{3} & 5 \end{pmatrix}$$

$$1A$$

$\text{(iii)} \quad P^{-1} \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} P = \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix}$ $\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix} = P \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} P^{-1}$ $\begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}^{12} = \underbrace{P \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} P^{-1} P \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} P^{-1} \dots P \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix} P^{-1}}_{12 \text{ times}}$ $= P \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix}^{12} P^{-1}$ $= \begin{pmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 5^{12} \end{pmatrix} \begin{pmatrix} \frac{1}{3} & -\frac{1}{3} \\ \frac{2}{3} & \frac{2}{3} \end{pmatrix}$ $= \begin{pmatrix} \frac{2}{3} & 5^{12} \\ -\frac{1}{3} & 5^{12} \end{pmatrix} \begin{pmatrix} 1 & -1 \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix}$ $= \begin{pmatrix} \frac{5^{12}+2}{3} & \frac{2 \cdot 5^{12}-2}{3} \\ \frac{5^{12}-1}{3} & \frac{2 \cdot 5^{12}+1}{3} \end{pmatrix}$	1M 1M 1M For $\begin{pmatrix} 1 & 0 \\ 0 & 5^{12} \end{pmatrix}$ OR $\begin{pmatrix} \frac{2}{3} & 1 \\ -\frac{1}{3} & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ \frac{5^{12}}{3} & \frac{2 \cdot 5^{12}}{3} \end{pmatrix}$ 1A OR $\begin{pmatrix} 81380209 & 162760416 \\ 81380208 & 162760417 \end{pmatrix}$
(11)	

[HKDSE 2012 M2 #11]

2.

$$8. \quad (a) \quad \begin{pmatrix} 1 & k & 0 \\ 0 & 1 & 1 \\ k & 0 & 0 \end{pmatrix}^{-1} = \frac{1}{\begin{vmatrix} 1 & k & 0 \\ 0 & 1 & 1 \\ k & 0 & 0 \end{vmatrix}} \begin{pmatrix} 0 & k & -k \\ 0 & 0 & k^2 \\ k & -1 & 1 \end{pmatrix}^{-1}$$

$$= \frac{1}{k^2} \begin{pmatrix} 0 & 0 & k \\ k & 0 & -1 \\ -k & k^2 & 1 \end{pmatrix}$$

1M+1A

1A

$$(b) \quad \begin{pmatrix} 1 & k & 0 \\ 0 & 1 & 1 \\ k & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ 1 \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} x \\ 1 \\ z \end{pmatrix} = \frac{1}{k^2} \begin{pmatrix} 0 & 0 & k \\ k & 0 & -1 \\ -k & k^2 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} \quad \text{by (a)}$$

$$= \frac{1}{k^2} \begin{pmatrix} k \\ 2k-1 \\ 2k^2-2k+1 \end{pmatrix}$$

1M

From the second row, we have $\frac{2k-1}{k^2} = 1$.

Alternative Solution

$$\begin{pmatrix} x+k \\ 1+z \\ kx \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$$

1M

From the first and third rows, we have $x+k=2$ and $x=\frac{1}{k}$.

$$\therefore \frac{1}{k} + k = 2.$$

i.e. $k^2 - 2k + 1 = 0$
 $k = 1$


1A

(5)

[HKDSE 2013 M2 #8]

3.

$$\begin{aligned}
 13. \quad (a) \quad (i) \quad MN &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} e & f \\ g & h \end{pmatrix} \\
 &= \begin{pmatrix} ae+bg & af+bh \\ ce+dg & cf+dh \end{pmatrix} \\
 \text{tr}(MN) &= ae+bg+cf+dh \\
 NM &= \begin{pmatrix} e & f \\ g & h \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\
 &= \begin{pmatrix} ea+fc & eb+fd \\ ga+hc & gb+hd \end{pmatrix} \\
 \text{tr}(NM) &= ea+fc+gb+hd \\
 \therefore \text{tr}(MN) &= \text{tr}(NM)
 \end{aligned}$$



$$\begin{aligned}
 (ii) \quad \text{tr}(BAB^{-1}) &= 1+3 \\
 \text{tr}(AB^{-1}B) &= 4 \quad \text{by (a)(i)} \\
 \text{tr}(A) &= 4
 \end{aligned}$$

 1M
1

$$\begin{aligned}
 (iii) \quad BAB^{-1} &= \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix} \\
 |BAB^{-1}| &= \begin{vmatrix} 1 & 0 \\ 0 & 3 \end{vmatrix} \\
 |B| \cdot |A| \cdot |B^{-1}| &= 1 \cdot 3 - 0 \cdot 0 \\
 |B| \cdot |A| \cdot |B|^{-1} &= 3 \\
 |A| &= 3
 \end{aligned}$$

 1M
1A
(6)

$$\begin{aligned}
 (b) \quad (i) \quad C \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda_1 \begin{pmatrix} x \\ y \end{pmatrix} \\
 \begin{pmatrix} px+qy \\ rx+sy \end{pmatrix} &= \begin{pmatrix} \lambda_1 x \\ \lambda_1 y \end{pmatrix} \\
 \begin{cases} (p-\lambda_1)x+qy=0 \\ rx+(s-\lambda_1)y=0 \end{cases}
 \end{aligned}$$

1A

Since this system of equations has non-zero solutions $\begin{pmatrix} x \\ y \end{pmatrix}$, $\begin{vmatrix} p-\lambda_1 & q \\ r & s-\lambda_1 \end{vmatrix} = 0$.

1

Similarly, $C \begin{pmatrix} x \\ y \end{pmatrix} = \lambda_2 \begin{pmatrix} x \\ y \end{pmatrix}$ gives $\begin{vmatrix} p-\lambda_2 & q \\ r & s-\lambda_2 \end{vmatrix} = 0$.

1

(ii) By (b)(i), λ_1 and λ_2 are the roots of the equation

$$\begin{vmatrix} p-\lambda & q \\ r & s-\lambda \end{vmatrix} = 0$$

1M

$$(p-\lambda)(s-\lambda)-qr=0$$

1


$$\lambda^2 - (p+s)\lambda + ps - qr = 0$$

$$\lambda^2 - \text{tr}(C) \cdot \lambda + |C| = 0$$

1

(5)

$$\begin{aligned}
 (c) \quad A \begin{pmatrix} x \\ y \end{pmatrix} &= \lambda \begin{pmatrix} x \\ y \end{pmatrix} \quad \text{for some non-zero matrices } \begin{pmatrix} x \\ y \end{pmatrix} \\
 \lambda^2 - \text{tr}(A) \cdot \lambda + |A| &= 0 \quad \text{by (b)(ii)} \\
 \lambda^2 - 4\lambda + 3 &= 0 \quad \text{by (a)(ii) \& (a)(iii)} \\
 \lambda &= 1 \text{ or } 3
 \end{aligned}$$



[HKDSE 2013 M2 #13]

4.

$$7. \quad (a) \quad A^2 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 2 & 0 & 2 \\ 0 & 4 & 0 \\ 2 & 0 & 2 \end{pmatrix}$$

$$= 2A$$

Hence the statement is true for $n=1$.

Assume the statement is true for $n=k$, i.e. $A^{k+1} = 2^k A$.

$$\begin{aligned} A^{k+2} &= A^{k+1}A \\ &= (2^k A)A \quad \text{by assumption} \\ &= 2^k A^2 \\ &= 2^k \cdot 2A \quad \text{by the statement for } n=1 \\ &= 2^{k+1} A \end{aligned}$$

Hence the statement is also true for $n=k+1$.

By the principle of mathematical induction, the statement is true for all positive integers n .

$$(b) \quad |A| = 0$$

Hence A^{-1} does not exist and so Willy arrives at a wrong conclusion by using A^{-1} .

1

1

1

1

1

1

1A

1

(7)

[HKDSE 2014 M2 #7]

5.

$$12. \text{ (a) (i)} \quad A^{-1} = \begin{pmatrix} 1 & p \\ -1 & 1 \end{pmatrix}^{-1}$$

$$= \frac{1}{|1-p|} \begin{pmatrix} 1 & 1 \\ -p & 1 \end{pmatrix}^T$$

$$= \frac{1}{1+p} \begin{pmatrix} 1 & -p \\ 1 & 1 \end{pmatrix}$$

1M

1A

$$\text{(ii)} \quad A^{-1}MA = \frac{1}{1+p} \begin{pmatrix} 1 & -p \\ 1 & 1 \end{pmatrix} \begin{pmatrix} k-1 & k \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & p \\ -1 & 1 \end{pmatrix}$$

$$= \frac{1}{1+p} \begin{pmatrix} k-p-1 & k \\ k & k \end{pmatrix} \begin{pmatrix} 1 & p \\ -1 & 1 \end{pmatrix}$$

$$= \frac{1}{1+p} \begin{pmatrix} -1-p & k+kp-p-p^2 \\ 0 & k+kp \end{pmatrix}$$

$$= \begin{pmatrix} -1 & k-p \\ 0 & k \end{pmatrix}$$

 1M+1A OR $\frac{1}{1+p} \begin{pmatrix} 1 & -p \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & k+kp-p \\ 1 & p \end{pmatrix}$

1

$$\text{(iii)} \quad \text{By (ii), } (A^{-1}MA)^n = \begin{pmatrix} -1 & 0 \\ 0 & k \end{pmatrix}^n \quad \text{for } p=k$$

$$A^{-1}M^nA = \begin{pmatrix} (-1)^n & 0 \\ 0 & k^n \end{pmatrix}$$

$$M^n = \begin{pmatrix} 1 & k \\ -1 & 1 \end{pmatrix} \begin{pmatrix} (-1)^n & 0 \\ 0 & k^n \end{pmatrix} \cdot \frac{1}{1+k} \begin{pmatrix} 1 & -k \\ 1 & 1 \end{pmatrix}$$

$$= \frac{1}{1+k} \begin{pmatrix} (-1)^n & k^{n+1} \\ (-1)^{n+1} & k^n \end{pmatrix} \begin{pmatrix} 1 & -k \\ 1 & 1 \end{pmatrix}$$

$$= \frac{1}{1+k} \begin{pmatrix} k^{n+1} + (-1)^n & k^{n+1} + (-1)^{n+1}k \\ k^n + (-1)^{n+1} & k^n + (-1)^n k \end{pmatrix}$$

1M For either side

 1M OR $\frac{1}{1+k} \begin{pmatrix} 1 & k \\ -1 & 1 \end{pmatrix} \begin{pmatrix} (-1)^n & (-1)^{n+1}k \\ k^n & k^n \end{pmatrix}$

1A

(8)

$$\text{(b)} \quad \begin{pmatrix} x_n \\ x_{n-1} \end{pmatrix} = M \begin{pmatrix} x_{n-1} \\ x_{n-2} \end{pmatrix} \quad \text{where } M = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix} \text{ after substituting } k=2$$

1M

$$= M^2 \begin{pmatrix} x_{n-2} \\ x_{n-3} \end{pmatrix}$$

1A

$$= \dots$$

$$= M^{n-2} \begin{pmatrix} x_2 \\ x_1 \end{pmatrix}$$

1A

$$= \frac{1}{1+2} \begin{pmatrix} 2^{n-1} + (-1)^{n-2} & 2^{n-1} + (-1)^{n-1}2 \\ 2^{n-2} + (-1)^{n-1} & 2^{n-2} + (-1)^{n-2}2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{by (a)(iii)}$$

1A

$$\therefore x_n = \frac{2^{n-1} + (-1)^{n-2}}{3}$$

 OR $\frac{2^{n-1} + (-1)^n}{3}$

(3)

[HKDSE 2014 M2 #12]

6.

6. (a) Note that $|M^T| = |M|$ and $|-M| = -|M|$.
 Since $|M^T| = |-M|$, we have $|M| = -|M|$.
 So, we have $2|M| = 0$.
 Thus, we have $|M| = 0$.

1M either one

1

(b) (i)

$$A + I = \begin{pmatrix} 0 & a & b \\ -a & 0 & -8 \\ -b & 8 & 0 \end{pmatrix}$$

So, we have $(A + I)^T = \begin{pmatrix} 0 & -a & -b \\ a & 0 & 8 \\ b & -8 & 0 \end{pmatrix} = -(A + I)$.

1M

By (a), we have $|A + I| = 0$.

1

$$\begin{aligned} A + I &= \begin{pmatrix} 0 & a & b \\ -a & 0 & -8 \\ -b & 8 & 0 \end{pmatrix} \\ |A + I| &= \begin{vmatrix} 0 & a & b \\ -a & 0 & -8 \\ -b & 8 & 0 \end{vmatrix} \\ &= 0 + 8ab - 8ab - 0 - 0 - 0 \\ &= 0 \end{aligned}$$

1M

1

(ii) Note that $A^3 + I = (A + I)(A^2 - A + I)$.

$$\begin{aligned} |A^3 + I| &= |A + I||A^2 - A + I| \\ &= (0)|A^2 - A + I| \quad (\text{by (b)(i)}) \\ &= 0 \end{aligned}$$

1M

Therefore, $A^3 + I$ is a singular matrix.
 Thus, the claim is agreed.

1A
-----(6) f.t.

[HKDSE 2015 M2 #6]

7.

11. (a) (i)
$$\begin{aligned} AB &= \frac{1}{(\lambda - \mu + 2)^2} \begin{pmatrix} \lambda - \mu + 1 & 1 \\ \lambda - \mu + 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -\lambda + \mu - 1 & \lambda - \mu + 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} BA &= \frac{1}{(\lambda - \mu + 2)^2} \begin{pmatrix} 1 & -1 \\ -\lambda + \mu - 1 & \lambda - \mu + 1 \end{pmatrix} \begin{pmatrix} \lambda - \mu + 1 & 1 \\ \lambda - \mu + 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} A + B &= \frac{1}{\lambda - \mu + 2} \left[\begin{pmatrix} \lambda - \mu + 1 & 1 \\ \lambda - \mu + 1 & 1 \end{pmatrix} + \begin{pmatrix} 1 & -1 \\ -\lambda + \mu - 1 & \lambda - \mu + 1 \end{pmatrix} \right] \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

(ii)
$$\begin{aligned} A^2 &= A(I - B) && (\text{by (a)(i)}) \\ &= A - AB \\ &= A - 0 && (\text{by (a)(i)}) \\ &= A \end{aligned}$$

$$\begin{aligned} B^2 &= B(I - A) && (\text{by (a)(i)}) \\ &= B - BA \\ &= B - 0 && (\text{by (a)(i)}) \\ &= B \end{aligned}$$

(iii)
$$\begin{aligned} &(\lambda + 1)A + (\mu - 1)B \\ &= \frac{\lambda + 1}{\lambda - \mu + 2}(I - \mu I + M) + \frac{\mu - 1}{\lambda - \mu + 2}(I + \lambda I - M) \\ &= \frac{-\mu\lambda + \lambda - \mu + 1}{\lambda - \mu + 2}I + \frac{\lambda + 1}{\lambda - \mu + 2}M + \frac{\mu\lambda - \lambda + \mu - 1}{\lambda - \mu + 2}I - \frac{\mu - 1}{\lambda - \mu + 2}M \\ &= M \end{aligned}$$

So, we have $M = (\lambda + 1)A + (\mu - 1)B$.

$$\begin{aligned} M^2 &= ((\lambda + 1)A + (\mu - 1)B)((\lambda + 1)A + (\mu - 1)B) \\ &= (\lambda + 1)^2 A^2 + (\lambda + 1)(\mu - 1)AB + (\lambda + 1)(\mu - 1)BA + (\mu - 1)^2 B^2 \\ &= (\lambda + 1)^2 A + (\mu - 1)^2 B && (\text{by (a)(i) and (a)(ii)}) \\ M^3 &= M^2 M \\ &= ((\lambda + 1)^2 A + (\mu - 1)^2 B)((\lambda + 1)A + (\mu - 1)B) \\ &= (\lambda + 1)^3 A^2 + (\lambda + 1)^2 (\mu - 1)AB + (\lambda + 1)(\mu - 1)^2 BA + (\mu - 1)^3 B^2 \\ &= (\lambda + 1)^3 A + (\mu - 1)^3 B && (\text{by (a)(i) and (a)(ii)}) \end{aligned}$$

Thus, we have $M^n = (\lambda + 1)^n A + (\mu - 1)^n B$.

1M either one
for both

1A

1A

1M

1

1

1M for using (a)(i) and (a)(ii)

1

$ \begin{aligned} & (\lambda+1)A + (\mu-1)B \\ &= \frac{\lambda+1}{\lambda-\mu+2} \begin{pmatrix} \lambda-\mu+1 & 1 \\ \lambda-\mu+1 & 1 \end{pmatrix} + \frac{\mu-1}{\lambda-\mu+2} \begin{pmatrix} 1 & -1 \\ -\lambda+\mu-1 & \lambda-\mu+1 \end{pmatrix} \\ &= \frac{1}{\lambda-\mu+2} \begin{pmatrix} \lambda^2 - \lambda\mu + 2\lambda & \lambda - \mu + 2 \\ (\lambda - \mu + 1)(\lambda - \mu + 2) & \lambda\mu - \mu^2 + 2\mu \end{pmatrix} \\ &= M \end{aligned} $ <p>So, the statement is true for $n = 1$.</p> <p>Assume that $M^k = (\lambda+1)^k A + (\mu-1)^k B$, where k is a positive integer.</p> $ \begin{aligned} & M^{k+1} \\ &= MM^k \\ &= ((\lambda+1)A + (\mu-1)B)((\lambda+1)^k A + (\mu-1)^k B) \\ &= (\lambda+1)^{k+1} A^2 + (\lambda+1)(\mu-1)^k AB + (\lambda+1)^k(\mu-1)BA + (\mu-1)^{k+1} B^2 \\ &= (\lambda+1)^{k+1} A + (\mu-1)^{k+1} B \quad (\text{by (a)(i) and (a)(ii)}) \\ \end{aligned} $ <p>So, the statement is true for $n = k+1$ if it is true for $n = k$.</p> <p>By mathematical induction, we have $M^n = (\lambda+1)^n A + (\mu-1)^n B$.</p>	1 1M 1M 1 ----- (8)
---	---

<p>(b) Note that $\begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix}^{315} = 2^{315} \begin{pmatrix} 2 & 1 \\ 0 & 3 \end{pmatrix}^{315}$.</p> <p>Also note that $3-2=1 \neq 2$ and $2-3+1=0$.</p> <p>Putting $\lambda=2$, $\mu=3$ and $n=315$ in (a)(iii), we have</p> $ \begin{aligned} & \begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix}^{315} \\ &= \frac{(2^{315})(3^{315})}{1} \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} + \frac{(2^{315})(2^{315})}{1} \begin{pmatrix} 1 & -1 \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 2^{630} & 6^{315} - 2^{630} \\ 0 & 6^{315} \end{pmatrix} \end{aligned} $	1M 1M 1M withhold 1M if checking is omitted 1A ----- (8)
---	---

$ \begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix} = \begin{pmatrix} 4 & 6-4 \\ 0 & 6 \end{pmatrix} $	1M ----- (4)
$ \begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix}^2 = \begin{pmatrix} 4 & 6-4 \\ 0 & 6 \end{pmatrix} \begin{pmatrix} 4 & 6-4 \\ 0 & 6 \end{pmatrix} = \begin{pmatrix} 4^2 & 6^2 - 4^2 \\ 0 & 6^2 \end{pmatrix} $	1M ----- (4)
$ \begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix}^3 = \begin{pmatrix} 4^2 & 6^2 - 4^2 \\ 0 & 6^2 \end{pmatrix} \begin{pmatrix} 4 & 6-4 \\ 0 & 6 \end{pmatrix} = \begin{pmatrix} 4^3 & 6^3 - 4^3 \\ 0 & 6^3 \end{pmatrix} $	1M withhold 1M if the step is skipped 1A ----- (4)
$ \begin{pmatrix} 4 & 2 \\ 0 & 6 \end{pmatrix}^{315} = \begin{pmatrix} 4^{315} & 6^{315} - 4^{315} \\ 0 & 6^{315} \end{pmatrix} = \begin{pmatrix} 2^{630} & 6^{315} - 2^{630} \\ 0 & 6^{315} \end{pmatrix} $	1A ----- (4)

[HKDSE 2015 M2 #11]

8.

8. (a) (i) A

$$= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$

$$\begin{aligned} A^2 \\ = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \end{aligned}$$

1A

(ii) A^3

$$\begin{aligned} &= \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \end{aligned}$$

1M

$$\begin{aligned} A^n \\ = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} \end{aligned}$$

1A

(iii) $\det(A^n)$

$$\begin{aligned} &= \begin{vmatrix} 1 & 0 \\ n & 1 \end{vmatrix} \\ &= 1 \end{aligned}$$

$$\begin{aligned} (A^{-1})^n \\ = (A^n)^{-1} \\ = \frac{1}{\det(A^n)} \begin{pmatrix} 1 & 0 \\ -n & 1 \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 \\ -n & 1 \end{pmatrix} \end{aligned}$$

1M

1A

$A^{-1} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$		
$(A^{-1})^2 = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}$		
$(A^{-1})^3 = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -3 & 1 \end{pmatrix}$	1M	
Thus, we have $(A^{-1})^n = \begin{pmatrix} 1 & 0 \\ -n & 1 \end{pmatrix}$.	1A	

1M

1A

-----(5)

$$(b) \quad (i) \quad \sum_{k=0}^{n-1} 2^k \\ = \frac{2^n - 1}{2 - 1} \\ = 2^n - 1$$

1A

$$(ii) \quad \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^0 & 2^1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^2 = \begin{pmatrix} 1 & 0 \\ 2^0 & 2^1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2^0 & 2^1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^0 + 2^1 & 2^2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^3 = \begin{pmatrix} 1 & 0 \\ 2^0 + 2^1 & 2^2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2^0 & 2^1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^0 + 2^1 + 2^2 & 2^3 \end{pmatrix}$$

So, we have $\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^n = \left(\sum_{k=0}^{n-1} 2^k \right) \begin{pmatrix} 1 & 0 \\ 2^n & 2^n \end{pmatrix}$.

By (b)(i), we have $\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^n = \begin{pmatrix} 1 & 0 \\ 2^n - 1 & 2^n \end{pmatrix}$.

1M

1A

$$\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^1 - 1 & 2^1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^2 = \begin{pmatrix} 1 & 0 \\ 2^1 - 1 & 2^1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^2 - 1 & 2^2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^3 = \begin{pmatrix} 1 & 0 \\ 2^2 - 1 & 2^2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2^3 - 1 & 2^3 \end{pmatrix}$$

Thus, we have $\begin{pmatrix} 1 & 0 \\ 1 & 2 \end{pmatrix}^n = \left(\sum_{k=0}^{n-1} 2^k \right) \begin{pmatrix} 1 & 0 \\ 2^n - 1 & 2^n \end{pmatrix}$.

1M

1A

[HKDSE 2016 M2 #8]

-----(3)

9.

12. (a)
$$A = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

$$= 3^1 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + 3^0 (I) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

So, the statement is true for $n=1$.

Assume that $A^k = 3^k I + 3^{k-1} k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, where k is a positive integer.

$$\begin{aligned} A^{k+1} &= A^k A \\ &= A^k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix} && (\text{by induction assumption}) \\ &= \left(3^k I + 3^{k-1} k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \left(3 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \\ &= 3^{k+1} I + 3^k k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + 3^k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + 3^{k-1} k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}^2 \\ &= 3^{k+1} I + 3^k (k+1) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \\ &= 3^{k+1} I + 3^k (k+1) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \end{aligned}$$

Therefore, the statement is true for $n=k+1$ if it is true for $n=k$.

By mathematical induction, the statement is true for all positive integers n .

1

IM

for using induction assumption

(b) (i) Note that $P^{-1} = \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix}$.

$$\begin{aligned} P^{-1} B P &= \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix} \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix} \\ &= A \end{aligned}$$

1A

1A

(ii) By (b)(i), we have $P^{-1} B P = A$.

So, we have $(P^{-1} B P)^n = A^n$.

Therefore, we have $P^{-1} B^n P = A^n$.

Hence, we have $B^n = P A^n P^{-1}$.

1M

$$\begin{aligned} B^n &= \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix} \left(3^n I + 3^{n-1} n \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right) \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix} \\ &= 3^n I + 3^{n-1} n \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix} \\ &= 3^n I + 3^{n-1} n \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \end{aligned}$$

1M

1

$\begin{aligned} B &= \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \\ &= 3^1 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + 3^0 (1) \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \end{aligned}$ <p>So, the statement is true for $n=1$.</p> <p>Assume that $B^k = 3^k I + 3^{k-1} k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}$, where k is a positive integer.</p> $\begin{aligned} B^{k+1} &= B^k B \\ &= \left(3^k I + 3^{k-1} k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \right) \left(\begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \right) \quad (\text{by induction assumption}) \\ &= \left(3^k I + 3^{k-1} k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \right) \left[3I + \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \right] \\ &= 3^{k+1} I + 3^k k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} + 3^k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} + 3^{k-1} k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}^2 \\ &= 3^{k+1} I + 3^k (k+1) \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \\ &= 3^{k+1} I + 3^k (k+1) \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \end{aligned}$ <p>Therefore, the statement is true for $n=k+1$ if it is true for $n=k$. By mathematical induction, the statement is true for all positive integers n.</p>	1M	for using induction assumption	
---	----	--------------------------------	--

<p>(iii) $A^m - B^m = 4m^2$</p> $\left 3^{m-1} m \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - 3^{m-1} m \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \right = 4m^2$ $(3^{m-1})^2 m^2 \begin{vmatrix} -2 & 0 \\ 4 & 2 \end{vmatrix} = 4m^2$ $-4m^2 (3^{2(m-1)}) = 4m^2$ $3^{2(m-1)} = -1$ <p>Note that $-1 < 0 < 3^{2(m-1)}$.</p> <p>Thus, there does not exist a positive integer m such that $A^m - B^m = 4m^2$.</p>	1M	1M	1A
---	----	----	----

[HKDSE 2017 M2 #12]

10.

7. (a)
$$MX = XM$$

$$\begin{pmatrix} 7 & 3 \\ -1 & 5 \end{pmatrix} \begin{pmatrix} a & 6a \\ b & c \end{pmatrix} = \begin{pmatrix} a & 6a \\ b & c \end{pmatrix} \begin{pmatrix} 7 & 3 \\ -1 & 5 \end{pmatrix}$$

$$\begin{pmatrix} 7a+3b & 42a+3c \\ -a+5b & -6a+5c \end{pmatrix} = \begin{pmatrix} a & 33a \\ 7b-c & 3b+5c \end{pmatrix}$$

$$7a+3b = a$$

$$-a+5b = 7b-c$$

$$42a+3c = 33a$$

$$-6a+5c = 3b+5c$$

$$b = -2a$$

$$c = -3a$$

1M

1M

1A

for both correct

(b)
$$\begin{vmatrix} X \end{vmatrix}$$

$$= \begin{vmatrix} a & 6a \\ -2a & -3a \end{vmatrix}$$

$$= (a)(-3a) - (6a)(-2a)$$

$$= 9a^2$$

Note that X is a non-zero real matrix.

By (a), a is a non-zero real number.

So, we have $\begin{vmatrix} X \end{vmatrix} > 0$.

Therefore, we have $\begin{vmatrix} X \end{vmatrix} \neq 0$.

Thus, X is a non-singular matrix.

1M

for considering determinant

1

(c)
$$(X^T)^{-1}$$

$$= (X^{-1})^T$$

$$= \left(\frac{1}{\begin{vmatrix} X \end{vmatrix}} \begin{pmatrix} -3a & -6a \\ 2a & a \end{pmatrix} \right)^T$$

1M

1M

$$= \left(\frac{1}{9a} \begin{pmatrix} -3 & -6 \\ 2 & 1 \end{pmatrix} \right)^T$$

1A

$$= \frac{1}{9a} \begin{pmatrix} -3 & 2 \\ -6 & 1 \end{pmatrix}$$

$$(X^T)^{-1}$$

$$= \begin{pmatrix} a & -2a \\ 6a & -3a \end{pmatrix}^{-1}$$

$$= \frac{1}{\begin{vmatrix} X^T \end{vmatrix}} \begin{pmatrix} -3a & 2a \\ -6a & a \end{pmatrix}$$

1M

$$= \frac{1}{\begin{vmatrix} X \end{vmatrix}} \begin{pmatrix} -3a & 2a \\ -6a & a \end{pmatrix}$$

1M

$$= \frac{1}{9a} \begin{pmatrix} -3 & 2 \\ -6 & 1 \end{pmatrix}$$

1A

-----(8)

[HKDSE 2018 M2 #7]

11.

$$11. \text{ (a)} \quad M^2 = \begin{pmatrix} 2 & 7 \\ -1 & -6 \end{pmatrix} \begin{pmatrix} 2 & 7 \\ -1 & -6 \end{pmatrix} = \begin{pmatrix} -3 & -28 \\ 4 & 29 \end{pmatrix}$$

1M

$$M^2 = aM + bI \quad \begin{pmatrix} -3 & -28 \\ 4 & 29 \end{pmatrix} = \begin{pmatrix} 2a & 7a \\ -a & -6a \end{pmatrix} + \begin{pmatrix} b & 0 \\ 0 & b \end{pmatrix}$$

$$\begin{cases} -3 = 2a + b \\ -28 = 7a \\ 4 = -a \\ 29 = -6a + b \end{cases}$$

1M

Thus, we have $a = -4$ and $b = 5$.

1A for both correct

-----(3)

- (b) Note that $(1 - (-5))M + (5 + (-5))I = 6M$.

1M

So, the statement is true for $n = 1$.

Assume that $6M^k = (1 - (-5)^k)M + (5 + (-5)^k)I$, where k is a positive integer.

$$\begin{aligned} 6M^{k+1} &= M(6M^k) \\ &= M((1 - (-5)^k)M + (5 + (-5)^k)I) \\ &= (1 - (-5)^k)M^2 + (5 + (-5)^k)M \\ &= (1 - (-5)^k)((1 + (-5))M + 5I) + (5 + (-5)^k)M \\ &= (1 + (-5) - (-5)^k - (-5)^{k+1})M + (5 + (-5)^{k+1})I + (5 + (-5)^k)M \\ &= (1 - (-5)^{k+1})M + (5 + (-5)^{k+1})I \end{aligned}$$

1M

for using the result of (a)

So, the statement is true for $n = k + 1$ if it is true for $n = k$.

By mathematical induction, we have $6M^n = (1 - (-5)^n)M + (5 + (-5)^n)I$.

1

$$\begin{aligned} 6M &= (1 - (-5))M + (5 + (-5))I \\ 6M^2 &= (1 - (-5))M^2 \\ &= (1 - (-5))((1 + (-5))M + 5I) \\ &= (1 - (-5)^2)M + (5 + (-5)^2)I \\ 6M^3 &= M(6M^2) \\ &= M((1 - (-5)^2)M + (5 + (-5)^2)I) \\ &= (1 - (-5)^2)M^2 + (5 + (-5)^2)M \\ &= (1 - (-5)^2)((1 + (-5))M + 5I) + (5 + (-5)^2)M \\ &= (1 + (-5) - (-5)^2 - (-5)^3)M + (5 + (-5)^3)I + (5 + (-5)^2)M \\ &= (1 - (-5)^3)M + (5 + (-5)^3)I \end{aligned}$$

1M

for using the result of (a)

Thus, we have $6M^n = (1 - (-5)^n)M + (5 + (-5)^n)I$.

1

-----(4)

<p>(c) By (b), we have $6M^n = (1 - (-5)^n)M + (5 + (-5)^n)I$ and $6M^{n+1} = (1 - (-5)^{n+1})M + (5 + (-5)^{n+1})I$. $6(1 - (-5)^n)M^{n+1} - 6(1 - (-5)^{n+1})M^n = ((1 - (-5)^n)(5 + (-5)^{n+1}) - (1 - (-5)^{n+1})(5 + (-5)^n))I$ $M^n \left(\frac{(1 - (-5)^n)M - (1 - (-5)^{n+1})I}{-6(-5)^n} \right) = \left(\frac{(1 - (-5)^n)M - (1 - (-5)^{n+1})I}{-6(-5)^n} \right) M^n = I$ So, we have $(M^n)^{-1} = \frac{(1 - (-5)^n)M - (1 - (-5)^{n+1})I}{-6(-5)^n}$.</p> $(M^n)^{-1} = \frac{(-5)^n - 1}{6(-5)^n} M + \frac{-(-5)^{n+1} + 1}{6(-5)^n} I$ $= \left(\frac{1}{6} M + \frac{5}{6} I \right) + \frac{1}{(-5)^n} \left(\frac{-1}{6} M + \frac{1}{6} I \right)$ <p>Letting $A = \frac{1}{6} M + \frac{5}{6} I = \frac{1}{6} \begin{pmatrix} 7 & 7 \\ -1 & -1 \end{pmatrix}$ and $B = \frac{-1}{6} M + \frac{1}{6} I = \frac{1}{6} \begin{pmatrix} -1 & -7 \\ 1 & 7 \end{pmatrix}$, we have $(M^n)^{-1} = A + \frac{1}{(-5)^n} B$.</p> <p>Thus, there exists a pair of 2×2 real matrices A and B such that $(M^n)^{-1} = A + \frac{1}{(-5)^n} B$ for all positive integers n.</p>	1M	1M	1M	1A	f.t.
$6M^n = (1 - (-5)^n)M + (5 + (-5)^n)I \quad (\text{by (b)})$ $M^n = \frac{1}{6}(1 - (-5)^n) \begin{pmatrix} 2 & 7 \\ -1 & -6 \end{pmatrix} + \frac{1}{6}(5 + (-5)^n) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ $= \begin{pmatrix} \frac{-(-5)^n + 7}{6} & \frac{-7(-5)^n + 7}{6} \\ \frac{(-5)^n - 1}{6} & \frac{7(-5)^n - 1}{6} \end{pmatrix}$ $\det(M^n) = \begin{pmatrix} \frac{-(-5)^n + 7}{6} & \frac{7(-5)^n - 1}{6} \\ \frac{7(-5)^n - 1}{6} & \frac{(-5)^n - 1}{6} \end{pmatrix}$ $= (-5)^n$ $(M^n)^{-1} = \frac{1}{\det(M^n)} \begin{pmatrix} \frac{7(-5)^n - 1}{6} & \frac{7(-5)^n - 7}{6} \\ \frac{-(-5)^n + 1}{6} & \frac{-(-5)^n + 7}{6} \end{pmatrix}$ $= \begin{pmatrix} \frac{7}{6} & \frac{7}{6} \\ \frac{-1}{6} & \frac{-1}{6} \end{pmatrix} + \frac{1}{(-5)^n} \begin{pmatrix} \frac{-1}{6} & \frac{-7}{6} \\ \frac{1}{6} & \frac{7}{6} \end{pmatrix}$ <p>Letting $A = \begin{pmatrix} \frac{7}{6} & \frac{7}{6} \\ \frac{-1}{6} & \frac{-1}{6} \end{pmatrix}$ and $B = \begin{pmatrix} \frac{-1}{6} & \frac{-7}{6} \\ \frac{1}{6} & \frac{7}{6} \end{pmatrix}$, we have $(M^n)^{-1} = A + \frac{1}{(-5)^n} B$.</p> <p>Thus, there exists a pair of 2×2 real matrices A and B such that $(M^n)^{-1} = A + \frac{1}{(-5)^n} B$ for all positive integers n.</p>	1M	1M	1M	1A	f.t.

-----(5)

[HKDSE 2019 M2 #11]

Systems of Linear Equations

1.

8. (a) The augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 2 & -1 & 5 & 6 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & -3 & 3 & 6 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 1 & -1 & -2 \end{array} \right)$$

Let $z = t$, where t is a real number. Then $y = t - 2$ and $x = 2 - 2t$.

1M

1A OR Solution Set =
 $\{(2 - 2t, t - 2, t) : t \in \mathbf{R}\}$

(b) Substitute $(x, y, z) = (2 - 2t, t - 2, t)$ into the last equation:

1M

$$(2 - 2t) - (t - 2) + \lambda(t) = 4$$

$$(\lambda - 3)t = 0$$

When $\lambda \neq 3$, $t = 0$.

$$\therefore (x, y, z) = (2, -2, 0)$$

When $\lambda = 3$, t can be any real number.

$$\therefore (x, y, z) = (2 - 2t, t - 2, t)$$

1A

1A

Alternative Solution

The augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 2 & -1 & 5 & 6 \\ 1 & -1 & \lambda & 4 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & -3 & 3 & 6 \\ 0 & -2 & \lambda - 1 & 4 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & \lambda - 3 & 0 \end{array} \right)$$

1M

When $\lambda \neq 3$, $z = 0$.

$$\therefore (x, y, z) = (2, -2, 0)$$

When $\lambda = 3$, z can be any real number.

$$\therefore (x, y, z) = (2 - 2t, t - 2, t), \text{ where } t \text{ is a real number.}$$

1A

1A

(5)

[HKDSE 2012 M2 #8]

2.

9. (a) The augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & -a & 1 & 2 \\ 2 & 1-2a & 2-b & a+4 \\ 3 & 1-3a & 3-ab & 4 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & -a & 1 & 2 \\ 0 & 1 & -b & a \\ 0 & 1 & -ab & -2 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & -a & 1 & 2 \\ 0 & 1 & -b & a \\ 0 & 0 & ab-b & a+2 \end{array} \right)$$

Hence the system has infinitely many solutions when

$$\begin{cases} b(a-1)=0 \\ a+2=0 \end{cases}$$

i.e. $a = -2$ and $b = 0$

1M

1M

1A

For both

(b) The system becomes

$$\begin{cases} x + 2y + z = 2 \\ 2x + 5y + 2z = 2 \\ 3x + 7y + 3z = 4 \end{cases}$$

i.e. $\begin{cases} x+z=6 \\ y=-2 \end{cases}$

$(x, y, z) = (6-t, -2, t)$ for any real number t

1M

1A

 OR $(t, -2, 6-t)$

(5)

[HKDSE 2013 M2 #9]

3.

9. (a) The augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 100 \\ 1 & 6 & 10 & 200 \end{array} \right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 100 \\ 0 & 5 & 9 & 100 \end{array} \right)$$

Let $z = t$, where t is a real number.

$$\therefore y = 20 - \frac{9t}{5} \text{ and } x = 80 + \frac{4t}{5}$$

1M

1A+1A

Alternative Solution

Let $y = t$, where t is a real number.

$$\therefore z = \frac{100 - 5t}{9} \text{ and } x = \frac{800 - 4t}{9}$$

OR, let $x = t$ and so

$$y = 200 - \frac{9t}{4}, z = \frac{5t}{4} - 100$$

(b) $\begin{cases} m + n + k = 100 \\ 0.5m + 3n + 5k = 200 \end{cases}$

$$\therefore \begin{cases} m + n + k = 100 \\ m + 6n + 10k = 200 \end{cases}$$

By (a), if both $20 - \frac{9t}{5}$ and $80 + \frac{4t}{5}$ are integers, then t is a multiple of 5.

$m \geq 0$ gives $t \geq -100$

$$n \geq 0 \text{ gives } t \leq \frac{100}{9}$$

$k \geq 0$ gives $t \geq 0$

Combining all the conditions above, we have $t = 0, 5 \text{ or } 10$.

1A

1M

Alternative Solution (1)

By trying out different values of t , we see that m, n and k are all non-negative when $t = 0, 5 \text{ or } 10$ (OR any two of these).

} 1M

Alternative Solution (2)

By trying out different values of t , we see that (m, n, k) can be $(80, 20, 0), (84, 11, 5)$ or $(88, 2, 10)$ (OR any two of these).

} 1M

Hence there are more than one set of combination of m, n and k and so Aubrey cannot be agreed with.

1

(6)

[HKDSE 2014 M2 #9]

4.

5. (a) The augmented matrix is $\left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 2 & 3 & -3 & 4 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -5 & 0 \end{array} \right)$.

1M

Thus, the solution set is $\{(2 - 6t, 5t, t) : t \in \mathbb{R}\}$.

1A

(b) Putting $(x, y, z) = (2 - 6t, 5t, t)$ in the last equation, we have

1M

$$3(2 - 6t) + 2(5t) + kt = 6.$$

So, we have $(k - 8)t = 0$.

We now consider the cases $k = 8$ and $k \neq 8$.

1M

Case 1: $k = 8$

The system of linear equations in (b) is equivalent to the system of linear equations in (a).

1M

Thus, the solution set is $\{(2 - 6t, 5t, t) : t \in \mathbb{R}\}$.

Case 2: $k \neq 8$

So, we have $t = 0$.

1A

Thus, the solution is $(2, 0, 0)$.

The augmented matrix is

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 2 & 3 & -3 & 4 \\ 3 & 2 & k & 6 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -5 & 0 \\ 0 & 0 & k-8 & 0 \end{array} \right).$$

1M

We now consider the cases $k = 8$ and $k \neq 8$.

1M

Case 1: $k = 8$

In this case, the augmented matrix becomes $\left(\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -5 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$.

Thus, the solution set is $\{(2 - 6t, 5t, t) : t \in \mathbb{R}\}$.

1M

Case 2: $k \neq 8$

The solution is $(2, 0, 0)$.

1A

-----(6)

[HKDSE 2015 M2 #5]

5.

11. (a) (i) (1) Note that

$$\begin{vmatrix} 1 & 1 & -1 \\ 4 & 6 & a \\ 5 & 1-a & 3a-1 \end{vmatrix}$$

$$= 6(3a-1) + (a)(5) + (-1)(4)(1-a) - (a)(1-a) - 4(3a-1) - (-1)(6)(5)$$

$$= (a+2)(a+12)$$

1A

$$\text{As } (E) \text{ has a unique solution, we have } \begin{vmatrix} 1 & 1 & -1 \\ 4 & 6 & a \\ 5 & 1-a & 3a-1 \end{vmatrix} \neq 0.$$

1M

 So, we have $(a+2)(a+12) \neq 0$.

 Thus, we have $a \neq -2$ and $a \neq -12$.

1

 The augmented matrix of (E) is

$$\begin{array}{l} \left(\begin{array}{ccc|c} 1 & 1 & -1 & 3 \\ 4 & 6 & a & b \\ 5 & 1-a & 3a-1 & b-1 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & -1 & 3 \\ 0 & 2 & a+4 & b-12 \\ 0 & -a-4 & 3a+4 & b-16 \end{array} \right) \\ \sim \left(\begin{array}{ccc|c} 1 & 1 & -1 & 3 \\ 0 & 2 & a+4 & b-12 \\ 0 & 0 & (a+2)(a+12) & ab-12a+6b-80 \end{array} \right) \end{array}$$

1M

1A

 As (E) has a unique solution, we have $(a+2)(a+12) \neq 0$.

 Thus, we have $a \neq -2$ and $a \neq -12$.

1

 (2) Since (E) has a unique solution, we have

$$x = \frac{\begin{vmatrix} 3 & 1 & -1 \\ b & 6 & a \\ b-1 & 1-a & 3a-1 \end{vmatrix}}{(a+2)(a+12)}$$

$$= \frac{3a^2 - ab + 50a + 6b - 24}{(a+2)(a+12)}$$

1M

for Cramer's Rule

$$y = \frac{\begin{vmatrix} 1 & 3 & -1 \\ 4 & b & a \\ 5 & b-1 & 3a-1 \end{vmatrix}}{(a+2)(a+12)}$$

$$= \frac{2(ab-10a+8)}{(a+2)(a+12)}$$

$$z = \frac{\begin{vmatrix} 1 & 1 & 3 \\ 4 & 6 & b \\ 5 & 1-a & b-1 \end{vmatrix}}{(a+2)(a+12)}$$

$$= \frac{ab-12a+6b-80}{(a+2)(a+12)}$$

1A+1A

1A for any one + 1A for all

<p>Since (E) has a unique solution, the augmented matrix of (E)</p> $\sim \left[\begin{array}{ccc c} 1 & 1 & -1 & 3 \\ 0 & 2 & a+4 & b-12 \\ 0 & 0 & (a+2)(a+12) & ab-12a+6b-80 \end{array} \right]$ $\sim \left[\begin{array}{ccc c} 1 & 0 & 0 & \frac{3a^2-ab+50a+6b-24}{(a+2)(a+12)} \\ 0 & 1 & 0 & \frac{2(ab-10a+8)}{(a+2)(a+12)} \\ 0 & 0 & 1 & \frac{ab-12a+6b-80}{(a+2)(a+12)} \end{array} \right]$ <p>Thus, we have</p> $\begin{cases} x = \frac{3a^2-ab+50a+6b-24}{(a+2)(a+12)} \\ y = \frac{2(ab-10a+8)}{(a+2)(a+12)} \\ z = \frac{ab-12a+6b-80}{(a+2)(a+12)} \end{cases}$	1M		
<p>(ii) (1) When $a = -2$, the augmented matrix of (E) is</p> $\left[\begin{array}{ccc c} 1 & 1 & -1 & 3 \\ 4 & 6 & -2 & b \\ 5 & 3 & -7 & b-1 \end{array} \right] \sim \left[\begin{array}{ccc c} 1 & 1 & -1 & 3 \\ 0 & 2 & 2 & b-12 \\ 0 & 0 & 0 & 2b-28 \end{array} \right]$ <p>Since (E) is consistent, we have $b = 14$.</p>	1M	1A	either one
<p>(2) When $a = -2$ and $b = 14$, the augmented matrix of (E)</p> $\sim \left[\begin{array}{ccc c} 1 & 1 & -1 & 3 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right]$ <p>Thus, the solution set of (E) is $\{(2t+2, 1-t, t) : t \in \mathbb{R}\}$.</p>	1A	(9)	
<p>(b) Putting $a = -2$ and $b = 14$, (E) becomes</p> $\begin{cases} x + y - z = 3 \\ 2x + 3y - z = 7 \\ 5x + 3y - 7z = 13 \end{cases}$ <p>By (b)(ii), the solution set is $\{(2t+2, 1-t, t) : t \in \mathbb{R}\}$.</p> <p>So, we have</p> $\begin{aligned} x^2 + y^2 - 6z^2 \\ = (2t+2)^2 + (1-t)^2 - 6t^2 \\ = -t^2 + 6t + 5 \\ = -(t^2 - 6t + 3^2) + 3^2 + 5 \\ = -(t-3)^2 + 14 \end{aligned}$ <p>Therefore, the greatest value of $x^2 + y^2 - 6z^2$ is 14.</p> <p>Thus, there is no real solution of the system of linear equations satisfying $x^2 + y^2 - 6z^2 > 14$.</p>	1M	1M	f.t.

[HKDSE 2016 M2 #11]

6.

5. (a) (i)

$$\begin{vmatrix} 1 & 2 & -1 \\ 3 & 8 & -11 \\ 2 & 3 & h \end{vmatrix} \neq 0$$

$$8h - 44 - 9 + 16 + 33 - 6h \neq 0$$

$$2h - 4 \neq 0$$

$$h \neq 2$$

$$h < 2 \text{ or } h > 2$$

1M

1A

(ii) z

$$= \frac{\begin{vmatrix} 1 & 2 & 11 \\ 3 & 8 & 49 \\ 2 & 3 & k \end{vmatrix}}{2h - 4}$$

$$= \frac{k - 14}{h - 2}$$

1M

1A

(b) When $h = 2$, the augmented matrix of (E) is

$$\left(\begin{array}{ccc|c} 1 & 2 & -1 & 11 \\ 3 & 8 & -11 & 49 \\ 2 & 3 & 2 & k \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 2 & -1 & 11 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & k - 14 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 7 & -5 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & k - 14 \end{array} \right).$$

1M

Since (E) has infinitely many solutions, we have $h = 2$ and $k = 14$.
Thus, the solution set is $\{(-7t - 5, 4t + 8, t) : t \in \mathbb{R}\}$.

1A

(6)

[HKDSE 2017 M2 #5]

7.

11. (a) (i) (1) Note that

$$\begin{vmatrix} 1 & a & 4(a+1) \\ 2 & a-1 & 2(a-1) \\ 1 & -1 & -12 \end{vmatrix}$$

$$= (a-1)(-12) + a(2)(a-1) + 4(a+1)(2)(-1) - 4(a-1)(a+1) + 2(a-1) - 2a(-12)$$

$$= -2(a-3)(a+1)$$

Since (E) has a unique solution, we have $\begin{vmatrix} 1 & a & 4(a+1) \\ 2 & a-1 & 2(a-1) \\ 1 & -1 & -12 \end{vmatrix} \neq 0$.

$$\text{So, we have } -2(a-3)(a+1) \neq 0.$$

$$\text{Therefore, we have } a \neq 3 \text{ and } a \neq -1.$$

$$\text{Thus, we have } a < -1, -1 < a < 3 \text{ or } a > 3.$$

1A

1M

1A

The augmented matrix of (E) is

$$\left(\begin{array}{ccc|c} 1 & a & 4(a+1) & 18 \\ 2 & a-1 & 2(a-1) & 20 \\ 1 & -1 & -12 & b \end{array} \right) \xrightarrow{\text{R}_2 - 2\text{R}_1} \left(\begin{array}{ccc|c} 1 & a & 4(a+1) & 18 \\ 0 & -a-1 & -6a-10 & -16 \\ 1 & -1 & -12 & b-18 \end{array} \right)$$

1M

$$\xrightarrow{\text{R}_3 - \text{R}_1} \left(\begin{array}{ccc|c} 1 & a & 4(a+1) & 18 \\ 0 & -a-1 & -6a-10 & -16 \\ 0 & 0 & 2a-6 & b-2 \end{array} \right)$$

1A

Since (E) has a unique solution, we have $2a-6 \neq 0$ and $-a-1 \neq 0$.

$$\text{Therefore, we have } a \neq 3 \text{ and } a \neq -1.$$

$$\text{Thus, we have } a < -1, -1 < a < 3 \text{ or } a > 3.$$

1A

- (2) Since
- (E)
- has a unique solution, we have

$$x = \frac{\begin{vmatrix} 18 & a & 4(a+1) \\ 20 & a-1 & 2(a-1) \\ b & -1 & -12 \end{vmatrix}}{-2(a-3)(a+1)} = \frac{a^2b + ab + 10a - 2b - 50}{(a-3)(a+1)}$$

1M

for Cramer's Rule

$$y = \frac{\begin{vmatrix} 1 & 18 & 4(a+1) \\ 2 & 20 & 2(a-1) \\ 1 & b & -12 \end{vmatrix}}{-2(a-3)(a+1)} = \frac{-3ab + 22a - 5b - 38}{(a-3)(a+1)}$$

$$z = \frac{\begin{vmatrix} 1 & a & 18 \\ 2 & a-1 & 20 \\ 1 & -1 & b \end{vmatrix}}{-2(a-3)(a+1)} = \frac{b-2}{2(a-3)}$$

1A+1A

1A for any one + 1A for all

<p>Since (E) has a unique solution, the augmented matrix of (E)</p> $\sim \left[\begin{array}{ccc c} 1 & a & 4(a+1) & 18 \\ 0 & -a-1 & -6a-10 & -16 \\ 0 & 0 & 2a-6 & b-2 \end{array} \right]$ $\sim \left[\begin{array}{ccc c} 1 & 0 & 0 & \frac{a^2b+ab+10a-2b-50}{(a-3)(a+1)} \\ 0 & 1 & 0 & \frac{-3ab+22a-5b-38}{(a-3)(a+1)} \\ 0 & 0 & 1 & \frac{b-2}{2(a-3)} \end{array} \right]$ <p>Thus, we have $\begin{cases} x = \frac{a^2b+ab+10a-2b-50}{(a-3)(a+1)} \\ y = \frac{-3ab+22a-5b-38}{(a-3)(a+1)} \\ z = \frac{b-2}{2(a-3)} \end{cases}$</p>	1M	1A+1A	1A for any one + 1A for all
<p>(ii) (1) When $a=3$, the augmented matrix of (E) is</p> $\left[\begin{array}{ccc c} 1 & 3 & 16 & 18 \\ 2 & 2 & 4 & 20 \\ 1 & -1 & -12 & b \end{array} \right] \sim \left[\begin{array}{ccc c} 1 & 3 & 16 & 18 \\ 0 & 1 & 7 & 4 \\ 0 & 0 & 0 & b-2 \end{array} \right]$ <p>Since (E) is consistent, we have $b=2$.</p> <p>(2) When $a=3$ and $b=2$, the augmented matrix of (E)</p> $\sim \left[\begin{array}{ccc c} 1 & 3 & 16 & 18 \\ 0 & 1 & 7 & 4 \\ 0 & 0 & 0 & 0 \end{array} \right]$ <p>Thus, the solution set of (E) is $\{(5u+6, -7u+4, u) : u \in \mathbb{R}\}$.</p>	1M	1A	either one
<p>(b) When $a=3$ and $b=s$, (E) becomes</p> $(G): \begin{cases} x + 3y + 16z = 18 \\ x + y + 2z = 10 \\ x - y - 12z = s \end{cases}$ <p>Since (F) is consistent, (G) is consistent.</p> <p>By (a)(ii), we have $s=2$.</p> <p>When $s=2$, the solution set of (G) is $\{(5u+6, -7u+4, u) : u \in \mathbb{R}\}$.</p> <p>Therefore, we have $2(5u+6) - 5(-7u+4) - 45u = t$.</p> <p>Solving, we have $t=-8$.</p> <p>Thus, we have $s=2$ and $t=-8$.</p>	1A ---(9)	1M 1M 1A	for both correct ---(3)

[HKDSE 2018 M2 #11]

8.

6. (a) (i) Note that

$$\begin{vmatrix} 1 & -2 & -2 \\ 5 & \alpha & \alpha \\ 7 & \alpha-3 & 2\alpha+1 \end{vmatrix}$$

$$= (\alpha)(2\alpha+1) + (-2)(\alpha)(7) + (-2)(5)(\alpha-3) - (\alpha)(\alpha-3) - (-2)(5)(2\alpha+1) - (-2)(\alpha)(7)$$

$$= (\alpha+4)(\alpha+10)$$

$$\text{Since } (E) \text{ has a unique solution, we have } \begin{vmatrix} 1 & -2 & -2 \\ 5 & \alpha & \alpha \\ 7 & \alpha-3 & 2\alpha+1 \end{vmatrix} \neq 0.$$

Hence, we have $(\alpha+4)(\alpha+10) \neq 0$.

So, we have $\alpha \neq -4$ and $\alpha \neq -10$.

Thus, we have $\alpha < -10$, $-10 < \alpha < -4$ or $\alpha > -4$.

1A

1M

1A

The augmented matrix of (E) is

$$\begin{array}{l} \left(\begin{array}{ccc|c} 1 & -2 & -2 & \beta \\ 5 & \alpha & \alpha & 5\beta \\ 7 & \alpha-3 & 2\alpha+1 & 8\beta \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & -2 & -2 & \beta \\ 0 & \alpha+10 & \alpha+10 & 0 \\ 0 & \alpha+11 & 2\alpha+15 & \beta \end{array} \right) \\ \sim \left(\begin{array}{ccc|c} 1 & -2 & -2 & \beta \\ 0 & 1 & \alpha+5 & \beta \\ 0 & 0 & (\alpha+4)(\alpha+10) & (\alpha+10)\beta \end{array} \right) \end{array}$$

1M

1A

Since (E) has a unique solution, we have $(\alpha+4)(\alpha+10) \neq 0$.

So, we have $\alpha \neq -4$ and $\alpha \neq -10$.

Thus, we have $\alpha < -10$, $-10 < \alpha < -4$ or $\alpha > -4$.

1A

- (ii) Since
- (E)
- has a unique solution, we have

$$\begin{array}{l} y \\ \left| \begin{array}{ccc|c} 1 & \beta & -2 & \\ 5 & 5\beta & \alpha & \\ 7 & 8\beta & 2\alpha+1 & \end{array} \right| \\ = \frac{(\alpha+4)(\alpha+10)}{\alpha+4} \\ = \frac{-\beta}{\alpha+4} \end{array}$$

1M for Cramer's Rule

1A

- (b) When
- $\alpha = -4$
- , the augmented matrix of
- (E)
- is

$$\begin{array}{l} \left(\begin{array}{ccc|c} 1 & -2 & -2 & \beta \\ 5 & -4 & -4 & 5\beta \\ 7 & -7 & -7 & 8\beta \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & -2 & -2 & \beta \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \beta \end{array} \right) \\ \text{Since } (E) \text{ is inconsistent, we have } \beta \neq 0. \\ \text{Thus, we have } \beta < 0 \text{ or } \beta > 0. \end{array}$$

1M

1A

-----(7)

[HKDSE 2019 M2 #6]

Vectors

1.

7. (a) The area of the parallelogram
- $OADB$

$$\begin{aligned} &= |(6\mathbf{i} + 2\mathbf{j} - \mathbf{k}) \times (2\mathbf{i} + \mathbf{j})| \\ &= |\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}| \\ &= \sqrt{1^2 + 2^2 + 2^2} \\ &= 3 \end{aligned}$$

- (b) The volume of the parallelepiped
- $OADBECFG$

$$\begin{aligned} &= (6\mathbf{i} + 2\mathbf{j} - \mathbf{k}) \cdot (2\mathbf{i} + \mathbf{j}) \cdot (5\mathbf{i} - \mathbf{j} + 2\mathbf{k}) \\ &= (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) \cdot (5\mathbf{i} - \mathbf{j} + 2\mathbf{k}) \\ &= 1 \cdot 5 + (-2)(-1) + 2 \cdot 2 \\ &= 11 \end{aligned}$$

Hence, the distance between point C and plane $OADB$ is $\frac{11}{3}$.

1M 1A 1M 1M+1A (5)		1M for height = $\frac{\text{volume}}{\text{base}}$
--------------------------------	--	---

[HKDSE 2012 M2 #7]


2.

12. (a) Let $\overrightarrow{AG} = \frac{\overrightarrow{AC} + \lambda \overrightarrow{AD}}{1+\lambda}$.

Since \overrightarrow{AG} lies on a median, $\overrightarrow{AG} = k \frac{\overrightarrow{AC} + \overrightarrow{AB}}{2} = k \frac{\overrightarrow{AC} + 2\overrightarrow{AD}}{2}$ for some k .

Comparing the two expressions of \overrightarrow{AG} , we get $\lambda = 2$.

$$\begin{aligned} \overrightarrow{AG} &= \frac{\overrightarrow{AC} + 2\overrightarrow{AD}}{3} \\ &= \frac{\overrightarrow{AC} + \overrightarrow{AB}}{3} \\ &= \frac{(\mathbf{c} - \mathbf{a}) + (\mathbf{b} - \mathbf{a})}{3} \end{aligned}$$



For tip-to-tail method

Alternative Solution 1
Let M be the mid-point of BC .

$$\begin{aligned} \overrightarrow{AG} &= \frac{2}{3} \overrightarrow{AM} \\ &= \frac{2}{3} \cdot \frac{\overrightarrow{AC} + \overrightarrow{AB}}{2} \\ &= \frac{(\mathbf{c} - \mathbf{a}) + (\mathbf{b} - \mathbf{a})}{3} \end{aligned}$$

1A

1M

For tip-to-tail method

Alternative Solution 2

$$\begin{aligned} \overrightarrow{AG} &= \overrightarrow{OG} - \overrightarrow{OA} \\ &= \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3} - \mathbf{a} \\ &= \frac{\mathbf{b} + \mathbf{c} - 2\mathbf{a}}{3} \end{aligned}$$

1M

1A


 For $\frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3}$

1A

(3)

(b) (i) Since O is the circumcentre of the $\triangle ABC$, $OD \perp AB$.
 $\therefore OD \parallel CE$
 $\angle DOG = \angle CFG$ (alt. \angle s, $OD \parallel CF$)
 $\angle ODG = \angle FCG$ (alt. \angle s, $OD \parallel CF$)
 $\angle OGD = \angle FGC$ (vert. opp. \angle s)
 $\therefore \triangle DOG \sim \triangle CFG$ (A.A.A.)
 $FG : GO = CG : GD$ (corr. sides, $\sim \triangle$ s)
 $= 2 : 1$

$$\begin{aligned} \text{(ii)} \quad \overrightarrow{AG} &= \frac{\overrightarrow{AF} + 2\overrightarrow{AO}}{3} \\ \overrightarrow{AF} &= 3\overrightarrow{AG} - 2\overrightarrow{AO} \\ &= 3 \cdot \frac{\mathbf{b} + \mathbf{c} - 2\mathbf{a}}{3} - 2(-\mathbf{a}) \end{aligned}$$



1

1

1A

1M

1M

For using (b)(i)

For using (a)

Alternative Solution 1

$$\begin{aligned} \overrightarrow{AF} &\approx \overrightarrow{AG} + \overrightarrow{GF} \\ &= \overrightarrow{AG} + 2\overrightarrow{OG} \\ &= \frac{\mathbf{b} + \mathbf{c} - 2\mathbf{a}}{3} + 2 \cdot \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3} \end{aligned}$$

1M

1M

For using (b)(i)

For using (a)

Alternative Solution 2

$$\begin{aligned} \overrightarrow{AF} &= \overrightarrow{AC} + \overrightarrow{CF} \\ &= \overrightarrow{AC} + 2\overrightarrow{OD} \\ &= \overrightarrow{AC} + \overrightarrow{OA} + \overrightarrow{OB} \\ &= (\mathbf{c} - \mathbf{a}) + \mathbf{a} + \mathbf{b} \end{aligned}$$

1M

1M

Alternative Solution 3

$$\begin{aligned} \overrightarrow{AF} &= \overrightarrow{OF} - \overrightarrow{OA} \\ &= 3\overrightarrow{OG} - \overrightarrow{OA} \\ &= 3 \cdot \frac{\mathbf{a} + \mathbf{b} + \mathbf{c}}{3} - \mathbf{a} \end{aligned}$$

1M

1M

For using (b)(i)

$$= \mathbf{b} + \mathbf{c}$$

1

$$\begin{aligned} \overrightarrow{AF} \cdot \overrightarrow{BC} &= (\mathbf{b} + \mathbf{c}) \cdot (\mathbf{c} - \mathbf{b}) \\ &= |\mathbf{c}|^2 - |\mathbf{b}|^2 \\ &= 0 \quad (\because O \text{ is the circumcentre}) \end{aligned}$$

1M

1A

$$\therefore AF \perp BC$$

$\therefore AF$ is another altitude of $\triangle ABC$.

Alternative Solution

$$\begin{aligned} \overrightarrow{BF} \cdot \overrightarrow{AC} &= (\overrightarrow{BA} + \overrightarrow{AF}) \cdot \overrightarrow{AC} \\ &= (\mathbf{a} - \mathbf{b} + \mathbf{b} + \mathbf{c}) \cdot (\mathbf{c} - \mathbf{a}) \\ &= |\mathbf{c}|^2 - |\mathbf{a}|^2 \\ &= 0 \quad (\because O \text{ is the circumcentre}) \end{aligned}$$

1M

1A

$$\therefore BF \perp AC$$

$\therefore BF$ is another altitude of $\triangle ABC$.

$$\therefore F \text{ is the orthocentre of } \triangle ABC.$$

1

(9)

[HKDSE 2012 M2 #12]

3.

$$10. \text{ (a)} \quad \overrightarrow{ON} = \frac{k\overrightarrow{OA} + \overrightarrow{OB}}{k+1}$$

$$= \frac{k(2\mathbf{i}) + (\mathbf{i} + 2\mathbf{j})}{k+1}$$

$$= \frac{(2k+1)\mathbf{i} + 2\mathbf{j}}{k+1}$$

$$\text{(b)} \quad \because \overrightarrow{MB} = 2\mathbf{j}, \quad \therefore BM \perp OA$$


Since A , N , P and M are concyclic, $ON \perp AB$.

$$\therefore \overrightarrow{ON} \cdot \overrightarrow{AB} = 0$$

$$\frac{(2k+1)\mathbf{i} + 2\mathbf{j}}{k+1} \cdot (\mathbf{i} + 2\mathbf{j} - 2\mathbf{i}) = 0$$

$$-(2k+1) + 2 \cdot 2 = 0$$

$$k = \frac{3}{2}$$




[HKDSE 2013 M2 #10]

4.

14. (a) (i) $\overrightarrow{AP} \cdot \overrightarrow{BP} = (\mathbf{p} - \mathbf{a}) \cdot (\mathbf{p} - \mathbf{b})$
 $= \mathbf{p} \cdot \mathbf{p} - \mathbf{a} \cdot \mathbf{p} - \mathbf{b} \cdot \mathbf{p} + \mathbf{a} \cdot \mathbf{b}$
 $= \mathbf{p} \cdot \mathbf{p} - (\mathbf{a} + \mathbf{b}) \cdot \mathbf{p} \quad (\because \mathbf{a} \cdot \mathbf{b} = 0)$

(ii) $\overrightarrow{AP} \cdot \overrightarrow{BP} + \overrightarrow{BP} \cdot \overrightarrow{CP} + \overrightarrow{CP} \cdot \overrightarrow{AP} = 0$
 By (i) and some similar results, we have
 $\mathbf{p} \cdot \mathbf{p} - (\mathbf{a} + \mathbf{b}) \cdot \mathbf{p} + \mathbf{p} \cdot \mathbf{p} - (\mathbf{b} + \mathbf{c}) \cdot \mathbf{p} + \mathbf{p} \cdot \mathbf{p} - (\mathbf{c} + \mathbf{a}) \cdot \mathbf{p} = 0$
 $3\mathbf{p} \cdot \mathbf{p} - 2(\mathbf{a} + \mathbf{b} + \mathbf{c}) \cdot \mathbf{p} = 0$
 $3\mathbf{p} \cdot \mathbf{p} - 2(3\mathbf{d}) \cdot \mathbf{p} = 0$
 $\mathbf{p} \cdot \mathbf{p} = 2\mathbf{p} \cdot \mathbf{d} \quad \text{----- (*)}$



(iii) $|\mathbf{p} - \mathbf{d}|^2 = (\mathbf{p} - \mathbf{d}) \cdot (\mathbf{p} - \mathbf{d})$
 $= \mathbf{p} \cdot \mathbf{p} - 2\mathbf{p} \cdot \mathbf{d} + \mathbf{d} \cdot \mathbf{d}$
 $= \mathbf{d} \cdot \mathbf{d} \quad \text{by (*)}$
 $= |\mathbf{d}|^2$

Hence $|\mathbf{p} - \mathbf{d}| = |\mathbf{d}|$.

$\therefore |\overrightarrow{DP}| = |\overrightarrow{OD}|$

$\therefore PD = OD \quad \text{----- (**)}$

Thus, the distance between P and D is a constant and therefore P lies on the sphere centred at D with fixed radius.

1M	
1	
	1
	(8)

(b) (i) Yes. Since O satisfies (**), O lies on the sphere mentioned in (i).

(ii) Yes. Since $\overrightarrow{DP_1} \times \overrightarrow{DP_2} = \overrightarrow{DP_2} \times \overrightarrow{DP_3}$, the normal vector of the plane containing D, P_1 and P_2 equals the normal vector of the plane containing D, P_2 and P_3 . Thus the plane containing D, P_1 and P_2 is parallel to the plane containing D, P_2 and P_3 . Since D and P_2 are common points of the planes, D, P_1, P_2 and P_3 are on the same plane. Since D is the centre of the sphere and P_1, P_2 and P_3 lie on the largest circle on the sphere, the radius of the circle equals the radius of the sphere, which is OD .

Follow through

1A	
	1M
	1M
	1A

[HKDSE 2013 M2 #14]

5.

8. (a) $\overrightarrow{OP} \times \overrightarrow{OQ} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 2 & 2 \\ 1 & -1 & 2 \end{vmatrix}$
 $= 6\mathbf{i} + 4\mathbf{j} - \mathbf{k}$

1A

$$\text{The volume of tetrahedron } OPQR = \frac{1}{6} |\overrightarrow{OP} \times \overrightarrow{OQ}| \cdot OR$$
 $= \frac{1}{6} |(6\mathbf{i} + 4\mathbf{j} - \mathbf{k}) \cdot (2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k})|$
 $= 1$

1M

1A

(b) $OR = \sqrt{2^2 + (-3)^2 + 6^2}$
 $= 7$

1A

$$\text{The area of } \triangle OPQ = \frac{1}{2} |\overrightarrow{OP} \times \overrightarrow{OQ}|$$
 $= \frac{1}{2} \sqrt{6^2 + 4^2 + (-1)^2}$
 $= \frac{\sqrt{53}}{2}$

1A

Let h be the height of the tetrahedron with OPQ as base.

$$\therefore \frac{1}{3} \cdot \frac{\sqrt{53}}{2} h = 1$$

1M

$$h = \frac{6}{\sqrt{53}}$$

Let θ be the angle between the plane OPQ and the line OR .

$$\therefore \sin \theta = \frac{6}{\sqrt{53}}$$

1M

Alternative Solution

$$|\overrightarrow{OP} \times \overrightarrow{OQ}| = \sqrt{6^2 + 4^2 + (-1)^2}$$
 $= \sqrt{53}$



1A

Let θ be the angle between the plane OPQ and the line OR .

$$\overrightarrow{OR} \cdot (\overrightarrow{OP} \times \overrightarrow{OQ}) = |\overrightarrow{OR}| \cdot |\overrightarrow{OP} \times \overrightarrow{OQ}| \cos(\theta + 90^\circ)$$

1M+1M

$$\cos(\theta + 90^\circ) = \frac{2 \cdot 6 - 3 \cdot 4 + 6(-1)}{7\sqrt{53}}$$



$$\theta \approx 6.8^\circ$$

1A


(8)

[HKDSE 2014 M2 #8]

6.

11. (a) (i) $\overrightarrow{OC} = t\mathbf{b}$
 $\therefore \overrightarrow{OE} = \frac{\mathbf{a} + mt\mathbf{b}}{1+m}$

(ii) $\overrightarrow{OD} = (1-t)\mathbf{a}$
 $\therefore \overrightarrow{OE} = \frac{n(1-t)\mathbf{a} + \mathbf{b}}{1+n}$



(iii) Comparing (i) and (ii), we have

$$\begin{cases} \frac{1}{1+m} = \frac{n(1-t)}{1+n} \\ \frac{mt}{1+m} = \frac{1}{1+n} \end{cases}$$

(2)+(1):

$$mt = \frac{1}{n(1-t)} \quad \text{(3)}$$

$$\text{By (1), } \frac{1}{1+\frac{1}{nt(1-t)}} = \frac{n(1-t)}{1+n}$$

$$t(1+n) = nt(1-t) + 1$$

$$t = -nt^2 + 1$$

$$n = \frac{1-t}{t^2}$$

$$\text{By (3), } mt = \frac{1}{\frac{1-t}{t^2}(1-t)}$$

$$m = \frac{t}{(1-t)^2}$$

(iv) If $m=n$, then $\frac{t}{(1-t)^2} = \frac{1-t}{t^2}$.

$$t^3 = (1-t)^3$$

$$t = \frac{1}{2}$$

Hence C and D are the mid-points of OB and OA respectively.
 Therefore, E is the centroid of $\triangle OAB$ and Chris is agreed with.

1M

1M

1

1

1A

1A

(9)

(b) $\overrightarrow{AC} = t\mathbf{b} - \mathbf{a}$

$$\overrightarrow{AC} \cdot \overrightarrow{OB} = (t\mathbf{b} - \mathbf{a}) \cdot \mathbf{b}$$

$$= 4t - \mathbf{a} \cdot \mathbf{b}$$

When $AC \perp OB$, $\overrightarrow{AC} \cdot \overrightarrow{OB} = 0$ which gives $\mathbf{a} \cdot \mathbf{b} = 4t$ (4)

$$\overrightarrow{BD} = (1-t)\mathbf{a} - \mathbf{b}$$

$$\overrightarrow{BD} \cdot \overrightarrow{OA} = [(1-t)\mathbf{a} - \mathbf{b}] \cdot \mathbf{a}$$

$$= (1-t) - \mathbf{a} \cdot \mathbf{b}$$

By (4), $\overrightarrow{BD} \cdot \overrightarrow{OA} = 1 - 5t$.

So, $\overrightarrow{BD} \cdot \overrightarrow{OA} \neq 0$ in general.
 i.e. BD is not always perpendicular to OA and Francis is not agreed with.

1A

1M

1A

1A

1A

(4)

[HKDSE 2014 M2 #11]

Since $\overrightarrow{OC} = \frac{3}{8}\overrightarrow{OA} + \frac{3}{16}\overrightarrow{OB}$, we have $\overrightarrow{OC} = \frac{21}{2}\mathbf{i} - \frac{21}{4}\mathbf{j} - \frac{9}{2}\mathbf{k}$.

$$\overrightarrow{CA} = \frac{19}{2}\mathbf{i} - \frac{3}{4}\mathbf{j} - \frac{15}{2}\mathbf{k}$$

$$\overrightarrow{CB} = \frac{11}{2}\mathbf{i} - \frac{43}{4}\mathbf{j} + \frac{9}{2}\mathbf{k}$$

$$\overrightarrow{CD} = \frac{-19}{2}\mathbf{i} + \frac{33}{4}\mathbf{j} - \frac{3}{2}\mathbf{k}$$

The volume of the tetrahedron $ABCD$

$$= \frac{1}{6} \left| \overrightarrow{CD} \cdot (\overrightarrow{CA} \times \overrightarrow{CB}) \right|$$

$$= \frac{1}{6} \begin{vmatrix} -19 & 33 & -3 \\ 2 & 4 & 2 \\ 19 & -3 & -15 \\ 2 & 4 & 2 \\ 11 & -43 & 9 \\ 2 & 4 & 2 \end{vmatrix}$$

$$= \frac{252}{6}$$

$$= 42$$

1M

1M

1A

-----(5)

[HKDSE 2015 M2 #10]

8.

12. (a) $|\overrightarrow{PA}| = |\overrightarrow{PB}|$

$$|\overrightarrow{OA} - \overrightarrow{OP}| = |\overrightarrow{OB} - \overrightarrow{OP}|$$

$$|- \mathbf{i} + (2-t)\mathbf{j} + 2\mathbf{k}| = |3\mathbf{i} + (1-t)\mathbf{j} + \mathbf{k}|$$

$$\sqrt{(-1)^2 + (2-t)^2 + 2^2} = \sqrt{3^2 + (1-t)^2 + 1^2}$$

$$t^2 - 4t + 9 = t^2 - 2t + 11$$

$$t = -1$$

1M

1M

1A

-----(3)

(b) (i) $\overrightarrow{CA} \times \overrightarrow{CB}$

$$= (-2\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}) \times (2\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})$$

$$= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 3 & -2 \\ 2 & 2 & -3 \end{vmatrix}$$

$$= -5\mathbf{i} - 10\mathbf{j} - 10\mathbf{k}$$

1M

$$|\overrightarrow{CA} \times \overrightarrow{CB}|$$

$$= \sqrt{(-5)^2 + (-10)^2 + (-10)^2}$$

$$= 15$$

A unit vector which is perpendicular to Π

$$= \frac{\overrightarrow{CA} \times \overrightarrow{CB}}{|\overrightarrow{CA} \times \overrightarrow{CB}|}$$

$$= \frac{-1}{3}\mathbf{i} - \frac{2}{3}\mathbf{j} - \frac{2}{3}\mathbf{k}$$

1M

1A

accept $\frac{1}{3}\mathbf{i} + \frac{2}{3}\mathbf{j} + \frac{2}{3}\mathbf{k}$

(ii) Note that $\overrightarrow{CD} = \mathbf{i} + 3\mathbf{j} + \mathbf{k}$.

$$\text{Let } \mathbf{n} = \frac{-1}{3}\mathbf{i} - \frac{2}{3}\mathbf{j} - \frac{2}{3}\mathbf{k}.$$

$$\overrightarrow{CD} \cdot \mathbf{n}$$

$$= (\mathbf{i} + 3\mathbf{j} + \mathbf{k}) \cdot \left(\frac{-1}{3}\mathbf{i} - \frac{2}{3}\mathbf{j} - \frac{2}{3}\mathbf{k} \right)$$

$$= \frac{-1}{3} - 2 - \frac{2}{3}$$

$$= -3$$

1M

Let θ be the angle between CD and Π .

Since $\overrightarrow{CD} \cdot \mathbf{n} < 0$, the angle between \overrightarrow{CD} and \mathbf{n} is $\frac{\pi}{2} + \theta$.

$$\overrightarrow{CD} \cdot \mathbf{n} = |\overrightarrow{CD}| |\mathbf{n}| \cos\left(\frac{\pi}{2} + \theta\right)$$

$$3 = \sqrt{1^2 + 3^2 + 1^2} \sin \theta$$

$$\sin \theta = \frac{3\sqrt{11}}{11}$$

$$\theta = \sin^{-1}\left(\frac{3\sqrt{11}}{11}\right)$$

1M

1A

Thus, the angle between CD and Π is $\sin^{-1}\left(\frac{3\sqrt{11}}{11}\right)$.

$$(iii) \quad \overrightarrow{DE}$$

$$= (\overrightarrow{DC} \cdot \mathbf{n}) \mathbf{n}$$

$$= (3)\left(\frac{-1}{3}\mathbf{i} - \frac{2}{3}\mathbf{j} - \frac{2}{3}\mathbf{k}\right)$$

$$= -\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}$$

1M

$$\overrightarrow{PF}$$

$$= \overrightarrow{PA} + \overrightarrow{PB} + \overrightarrow{PC}$$

$$= (-\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}) + (3\mathbf{i} + 2\mathbf{j} + \mathbf{k}) + (\mathbf{i} + 4\mathbf{k})$$

$$= 3\mathbf{i} + 5\mathbf{j} + 7\mathbf{k}$$

$$\overrightarrow{FD}$$

$$= \overrightarrow{OD} - (\overrightarrow{OP} + \overrightarrow{PF})$$

$$= (3\mathbf{i} + 2\mathbf{j} + 5\mathbf{k}) - ((\mathbf{i} - \mathbf{j}) + (3\mathbf{i} + 5\mathbf{j} + 7\mathbf{k}))$$

$$= -\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}$$

1M

Therefore, we have $\overrightarrow{DE} = \overrightarrow{FD}$.

So, we have $DE \parallel FD$ and $DE = DF$.

Hence, D , E and F are collinear and $DE = DF$.

Thus, D is the mid-point of the line segment joining E and F .

1A+1A f.t.

[HKDSE 2016 M2 #12]

-----(10)

9.

3. (a) \overrightarrow{OP}

$$= \frac{2}{2+3} \mathbf{a} + \frac{3}{2+3} \mathbf{b}$$

$$= \frac{2}{5} \mathbf{a} + \frac{3}{5} \mathbf{b}$$

1A

(b) (i) $\mathbf{a} \cdot \mathbf{b}$
 $= |\mathbf{a}| |\mathbf{b}| \cos \angle AOB$
 $= (45)(20)\left(\frac{1}{4}\right)$
 $= 225$

1M

1A

(ii) $|\overrightarrow{OP}|^2$
 $= \left(\frac{2}{5} \mathbf{a} + \frac{3}{5} \mathbf{b}\right) \cdot \left(\frac{2}{5} \mathbf{a} + \frac{3}{5} \mathbf{b}\right)$
 $= \frac{4}{25} |\mathbf{a}|^2 + 2\left(\frac{2}{5}\right)\left(\frac{3}{5}\right) \mathbf{a} \cdot \mathbf{b} + \frac{9}{25} |\mathbf{b}|^2$
 $= 324 + 108 + 144$
 $= 576$

1M

for using (b)(i)

$$\begin{aligned} &|\overrightarrow{OP}| \\ &= \sqrt{576} \\ &= 24 \end{aligned}$$

1A

[HKDSE 2017 M2 #3]

10.

10. (a) Note that $\overrightarrow{AB} = 2\mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\overrightarrow{AC} = 6\mathbf{i} - 6\mathbf{j}$.

$$\begin{aligned}\overrightarrow{AE} &= \frac{1}{1+r} \overrightarrow{AC} + \frac{r}{1+r} \overrightarrow{AB} \\ &= \frac{2r+6}{r+1} \mathbf{i} + \frac{r-6}{r+1} \mathbf{j} + \frac{r}{r+1} \mathbf{k} \\ \text{Also note that } \overrightarrow{AE} &= \frac{1}{11} \overrightarrow{AF} + \frac{10}{11} \overrightarrow{AD} \text{ and } \overrightarrow{AC} = 2 \overrightarrow{AD}.\end{aligned}$$

$$\begin{aligned}\overrightarrow{AF} &= 11\overrightarrow{AE} - 5\overrightarrow{AC} \\ &= \frac{-8r+36}{r+1} \mathbf{i} + \frac{41r-36}{r+1} \mathbf{j} + \frac{11r}{r+1} \mathbf{k}\end{aligned}$$

Since A , B and F are collinear, we have $\frac{2}{-8r+36} = \frac{1}{41r-36} = \frac{1}{11r}$.

Solving, we have $r = \frac{6}{5}$.

1M	any one -----
1A	for both -----
1M	
1A	1.2
-----	(4)

- (b) (i) Note that $\overrightarrow{AD} = \frac{1}{2} \overrightarrow{AC} = 3\mathbf{i} - 3\mathbf{j}$.

$$\text{By (a), we have } \overrightarrow{AE} = \frac{1}{11}(42\mathbf{i} - 24\mathbf{j} + 6\mathbf{k}) .$$

$$\begin{aligned}\overrightarrow{AD} \cdot \overrightarrow{DE} &= \overrightarrow{AD} \cdot (\overrightarrow{AE} - \overrightarrow{AD}) \\ &= (3\mathbf{i} - 3\mathbf{j}) \cdot \left(\frac{1}{11}(42\mathbf{i} - 24\mathbf{j} + 6\mathbf{k}) \right) \\ &= 0\end{aligned}$$

$$(ii) \quad \overrightarrow{AB} \cdot \overrightarrow{BC}$$

$$\begin{aligned}&= \overrightarrow{AB} \cdot (\overrightarrow{AC} - \overrightarrow{AB}) \\ &= (2\mathbf{i} + \mathbf{j} + \mathbf{k}) \cdot (4\mathbf{i} - 7\mathbf{j} - \mathbf{k}) \\ &= 0\end{aligned}$$

Therefore, we have $\angle ABC = 90^\circ = \angle ADE$.

So, we have $\angle CBF = 90^\circ = \angle CDF$.

Thus, B , D , C and F are concyclic.

1M	for using (a)
1A	
1M	
1M	
-----	(5)

- (c) Note that $\overrightarrow{AF} = 12\mathbf{i} + 6\mathbf{j} + 6\mathbf{k}$ and $\overrightarrow{AP} = \mathbf{i} + 7\mathbf{j} - 2\mathbf{k}$.

Since $\angle CBF = 90^\circ$, Q is the mid-point of CF .

$$\text{Therefore, we have } \overrightarrow{AQ} = \frac{1}{2}(\overrightarrow{AC} + \overrightarrow{AF}) = 9\mathbf{i} + 3\mathbf{k} .$$

The volume of the tetrahedron $ABPQ$

$$\begin{aligned}&= \frac{1}{6} \left| \overrightarrow{AQ} \cdot (\overrightarrow{AB} \times \overrightarrow{AP}) \right| \\ &= \frac{1}{6} \begin{vmatrix} 9 & 0 & 3 \\ 2 & 1 & 1 \\ 1 & 7 & -2 \end{vmatrix} \\ &= 7\end{aligned}$$

1M

1M

1A

----- (3)

[HKDSE 2017 M2 #10]

11.

12. (a) (i) Note that $\overrightarrow{AB} = -5\mathbf{i} + 6\mathbf{j} - 4\mathbf{k}$ and $\overrightarrow{AC} = 3\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}$.

$$\begin{aligned}\overrightarrow{AB} \times \overrightarrow{AC} \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -5 & 6 & -4 \\ 3 & 2 & 4 \end{vmatrix} \\ &= 32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}\end{aligned}$$

1A

- (ii) Note that $\overrightarrow{AD} = -\mathbf{i} + \mathbf{j} - 6\mathbf{k}$.

The required volume

$$\begin{aligned}&= \frac{1}{6} |(\overrightarrow{AB} \times \overrightarrow{AC}) \cdot \overrightarrow{AD}| \\ &= \frac{1}{6} |(32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}) \cdot (-\mathbf{i} + \mathbf{j} - 6\mathbf{k})| \\ &= \frac{1}{6} |(32)(-1) + (8)(1) + (-28)(-6)| \\ &= 24\end{aligned}$$

1M

1A

- (iii) \overrightarrow{DE}

$$\begin{aligned}&= \left(\overrightarrow{DA} \cdot \frac{32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}}{\sqrt{32^2 + 8^2 + (-28)^2}} \right) \left(\frac{32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}}{\sqrt{32^2 + 8^2 + (-28)^2}} \right) \\ &= \left((\mathbf{i} - \mathbf{j} + 6\mathbf{k}) \cdot \frac{32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}}{\sqrt{32^2 + 8^2 + (-28)^2}} \right) \left(\frac{32\mathbf{i} + 8\mathbf{j} - 28\mathbf{k}}{\sqrt{32^2 + 8^2 + (-28)^2}} \right) \\ &= \frac{-32}{13}\mathbf{i} - \frac{8}{13}\mathbf{j} + \frac{28}{13}\mathbf{k}\end{aligned}$$

1M

1A

----- (5)

- (b) (i) Let $\overrightarrow{BF} = t\overrightarrow{BC}$, where $0 < t < 1$.

$$\begin{aligned}\overrightarrow{DF} \\ &= (1-t)\overrightarrow{DB} + t\overrightarrow{DC} \\ &= (1-t)(-4\mathbf{i} + 5\mathbf{j} + 2\mathbf{k}) + t(4\mathbf{i} + \mathbf{j} + 10\mathbf{k}) \\ &= (8t-4)\mathbf{i} + (5-4t)\mathbf{j} + (8t+2)\mathbf{k}\end{aligned}$$

1M

Since $DF \perp BC$, we have $\overrightarrow{DF} \cdot \overrightarrow{BC} = 0$.

1M

Note that $\overrightarrow{BC} = 8\mathbf{i} - 4\mathbf{j} + 8\mathbf{k}$.

Hence, we have $(8t-4)(8) + (5-4t)(-4) + (8t+2)(8) = 0$.

So, we have $144t - 36 = 0$.

Solving, we have $t = \frac{1}{4}$.

Thus, we have $\overrightarrow{DF} = -2\mathbf{i} + 4\mathbf{j} + 4\mathbf{k}$.

1A

$$\begin{aligned}
 \text{(ii)} \quad & \overrightarrow{EF} \\
 &= \overrightarrow{DF} - \overrightarrow{DE} \\
 &= -2\mathbf{i} + 4\mathbf{j} + 4\mathbf{k} - \left(\frac{-32}{13}\mathbf{i} - \frac{8}{13}\mathbf{j} + \frac{28}{13}\mathbf{k} \right) \\
 &= \frac{6}{13}\mathbf{i} + \frac{60}{13}\mathbf{j} + \frac{24}{13}\mathbf{k}
 \end{aligned}$$

1M

$$\begin{aligned}
 & \overrightarrow{BC} \cdot \overrightarrow{EF} \\
 &= (8\mathbf{i} - 4\mathbf{j} + 8\mathbf{k}) \cdot \left(\frac{6}{13}\mathbf{i} + \frac{60}{13}\mathbf{j} + \frac{24}{13}\mathbf{k} \right) \\
 &= 8\left(\frac{6}{13}\right) - 4\left(\frac{60}{13}\right) + 8\left(\frac{24}{13}\right) \\
 &= 0
 \end{aligned}$$

Thus, \overrightarrow{BC} is perpendicular to \overrightarrow{EF} .

 1A f.t.
-----(5)

- (c) Note that the required angle is $\angle DFE$.
 $\cos \angle DFE$

1M for identifying the required angle

$$\begin{aligned}
 & = \frac{\overrightarrow{DF} \cdot \overrightarrow{EF}}{\|\overrightarrow{DF}\| \|\overrightarrow{EF}\|} \\
 &= \frac{(-2\mathbf{i} + 4\mathbf{j} + 4\mathbf{k}) \cdot \left(\frac{6}{13}\mathbf{i} + \frac{60}{13}\mathbf{j} + \frac{24}{13}\mathbf{k} \right)}{\sqrt{(-2)^2 + 4^2 + 4^2} \sqrt{\left(\frac{6}{13}\right)^2 + \left(\frac{60}{13}\right)^2 + \left(\frac{24}{13}\right)^2}} \\
 &= \frac{324}{(6)(18\sqrt{13})} \\
 &= \frac{3}{\sqrt{13}} \\
 &= \frac{3\sqrt{13}}{13}
 \end{aligned}$$

1M

Thus, the required angle is $\cos^{-1}\left(\frac{3\sqrt{13}}{13}\right)$.

 1A
-----(3)

[HKDSE 2018 M2 #12]

12.

12. (a) $\begin{aligned} \overrightarrow{AC} &= \overrightarrow{BC} \\ \overrightarrow{OC} - \overrightarrow{OA} &= \overrightarrow{OC} - \overrightarrow{OB} \\ -6\mathbf{i} - 8\mathbf{j} + (t-2)\mathbf{k} &= -8\mathbf{j} + (t-8)\mathbf{k} \\ \sqrt{(-6)^2 + (-8)^2 + (t-2)^2} &= \sqrt{(-8)^2 + (t-8)^2} \\ t^2 - 4t + 104 &= t^2 - 16t + 128 \\ 12t &= 24 \\ t &= 2 \end{aligned}$	1M 1M 1A -----(3)
(b) $\begin{aligned} &\overrightarrow{AB} \times \overrightarrow{AC} \\ &= (-6\mathbf{i} + 6\mathbf{k}) \times (-6\mathbf{i} - 8\mathbf{j}) \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -6 & 0 & 6 \\ -6 & -8 & 0 \end{vmatrix} \\ &= ((0)(0) - (6)(-8))\mathbf{i} + ((6)(-6) - (6)(0))\mathbf{j} + ((-6)(-8) - (0)(-6))\mathbf{k} \\ &= 48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k} \end{aligned}$	1M 1A -----(2)
(c) The volume of the pyramid $OABC$ $\begin{aligned} &= \frac{1}{6} \overrightarrow{OA} \cdot (\overrightarrow{AB} \times \overrightarrow{AC}) \\ &= \frac{1}{6} (\mathbf{i} - 4\mathbf{j} + 2\mathbf{k}) \cdot (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) \\ &= \frac{1}{6} (1)(48) + (-4)(-36) + (2)(48) \\ &= 48 \end{aligned}$	1M 1A -----(2)
The volume of the pyramid $OABC$ $\begin{aligned} &= \frac{1}{6} \overrightarrow{OA} \cdot (\overrightarrow{OB} \times \overrightarrow{OC}) \\ &= \frac{1}{6} \begin{vmatrix} 1 & -4 & 2 \\ -5 & -4 & 8 \\ -5 & -12 & 2 \end{vmatrix} \\ &= \frac{1}{6} (1)(-4)(2) + (-4)(8)(-5) + (2)(-5)(-12) - (1)(8)(-12) - (-4)(-5)(2) - (2)(-4)(-5) \\ &= 48 \end{aligned}$	1M 1A -----(2)
(d) (i) By (c), the volume of the pyramid $OABC$ is not equal to 0. So, O does not lie on Π . Therefore, \overrightarrow{OP} , \overrightarrow{OQ} and \overrightarrow{OR} are non-zero vectors. Hence, we have $p \neq 0$, $q \neq 0$ and $r \neq 0$. Thus, we have $pqr \neq 0$.	1

$$\begin{aligned}
 \text{(ii)} \quad & \overrightarrow{OD} \\
 &= \overrightarrow{OA} \cdot \frac{\overrightarrow{AB} \times \overrightarrow{AC}}{|\overrightarrow{AB} \times \overrightarrow{AC}|^2} (\overrightarrow{AB} \times \overrightarrow{AC}) \\
 &= \frac{(6)(48)}{48^2 + (-36)^2 + 48^2} (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) \\
 &= \frac{288}{5904} (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) \\
 &= \frac{96}{41} \mathbf{i} - \frac{72}{41} \mathbf{j} + \frac{96}{41} \mathbf{k}
 \end{aligned}$$

1M

$$\begin{aligned}
 \text{(iii)} \quad & \overrightarrow{AP} \cdot (\overrightarrow{AB} \times \overrightarrow{AC}) = 0 \\
 & ((p-1)\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}) \cdot (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) = 0 \\
 & 48p - 48 - 144 - 96 = 0 \\
 & p = 6
 \end{aligned}$$

1A

$$\begin{aligned}
 & \overrightarrow{AQ} \cdot (\overrightarrow{AB} \times \overrightarrow{AC}) = 0 \\
 & (-\mathbf{i} + (q+4)\mathbf{j} - 2\mathbf{k}) \cdot (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) = 0 \\
 & -48 - 36q - 144 - 96 = 0 \\
 & q = -8
 \end{aligned}$$

1M

$$\begin{aligned}
 & \overrightarrow{AR} \cdot (\overrightarrow{AB} \times \overrightarrow{AC}) = 0 \\
 & (-\mathbf{i} + 4\mathbf{j} + (r-2)\mathbf{k}) \cdot (48\mathbf{i} - 36\mathbf{j} + 48\mathbf{k}) = 0 \\
 & -48 - 144 + 48r - 96 = 0 \\
 & r = 6
 \end{aligned}$$

1A

So, we have $\overrightarrow{OE} = \frac{1}{6}\mathbf{i} - \frac{1}{8}\mathbf{j} + \frac{1}{6}\mathbf{k}$.

By (b)(ii), we have $\overrightarrow{OE} = \frac{41}{576} \overrightarrow{OD}$.
Thus, D , E and O are collinear.

1A ft.

(6)

[HKDSE 2019 M2 #12]

(6)