HKDSE MATH EP

M2

HONG KONG EDUCATIONAL PUBLISHING COMPANY

HONG KONG DIPLOMA OF
SECONDARY EDUCATION EXAMINATION

## **MATHEMATICS Extended Part**

## Module 2 (Algebra and Calculus) MOCK EXAM 1 Question-Answer Book

Time allowed: 2½ hours

This paper must be answered in English

## INSTRUCTIONS

- 1. After the announcement of the start of the examination, you should first write your Candidate Number in the space provided on Page 1 and stick barcode labels in the spaces provided on Pages 1, 3, 5, 7, 9 and 11.
- 2. This paper consists of TWO sections, A and B.
- 3. Attempt ALL questions in this paper. Write your answers in the spaces provided in this Question-Answer Book. Do not write in the margins. Answers written in the margins will not be marked.
- 4. Graph paper and supplementary answer sheets will be supplied on request. Write your Candidate Number, mark the question number box and stick a barcode label on each sheet, and fasten them with string INSIDE this book.
- 5. Unless otherwise specified, all working must be clearly shown.
- 6. Unless otherwise specified, numerical answers must be exact.
- 7. In this paper, vectors may be represented by bold-type letters such as  $\mathbf{u}$ , but candidates are expected to use appropriate symbols such as  $\mathbf{u}$  in their working.
- 8. The diagrams in this paper are not necessarily drawn to scale.
- 9. No extra time will be given to candidates for sticking on the barcode labels or filling in the question number boxes after the 'Time is up' announcement.
- © 香港教育圖書公司 保留版權 Hong Kong Educational Publishing Company All Rights Reserved 2016

·	Please	stick	the	barc	ode	lab	el h	ere.
·								
				_				•
	Candidate Number							

## FORMULAS FOR REFERENCE

$$\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$cos(A \pm B) = cos A cos B \mp sin A sin B$$

$$\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$2 \sin A \cos B = \sin (A + B) + \sin (A - B)$$

$$2\cos A\cos B = \cos (A+B) + \cos (A-B)$$

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B)$$

$$\sin A + \sin B = 2\sin \frac{A+B}{2}\cos \frac{A-B}{2}$$

$$\sin A - \sin B = 2\cos \frac{A+B}{2}\sin \frac{A-B}{2}$$

$$\sin A - \sin B = 2\cos\frac{A+B}{2}\sin\frac{A-B}{2}$$

$$\cos A + \cos B = 2\cos\frac{A+B}{2}\cos\frac{A-B}{2}$$

$$\cos A - \cos B = -2\sin\frac{A+B}{2}\sin\frac{A-B}{2}$$

Answers written in the margins will not be marked.

Expand  $(4-x)^3$ . Hence, find the constant term in the expansion of  $(4-x)^3 \left(1+\frac{6}{x}\right)^4$ .

(5 marks)

Please stick the barcode label here.

	<u> </u>			 -			
	-			<del></del>			
		-116				••	
	**						
	1.02					<del></del>	_
	_						
			<del> </del>		•		
	<u>,</u> ,	<del>a.</del>				•••	
•				 			_
_		•					
-	<u> </u>						_
			~~~				
							_

Answers written in the margins will not be marked.

(4 marks)

2013

2. Find  $\frac{d}{dx}(\sin x)$  from first principles.

2016

Please stick the barcode label here.

	angents to the curve at these two points.	(7 mar
-		
-		
, <del>,-</del>		
-		
_		<del></del>
-		
-		
-	(	
-	•	
-		
-		
		,
		<del></del>
	•	
	·	

swers written in the margins will not be marked	
written in the margins	rked
written in the margins	ma
written in the margins	<u>8</u>
written in the margins	no
written in the	Will
written in the	margins
written	the
written	.⊟
swers	written
Å	Answers

<b>016</b> (a)	Using mathematica	l induction,	prove that	$\sum_{k=1}^{n} \left(-1\right)^{k} k =$	$=\frac{(-1)^n(2n+1)}{4}$	$\frac{-1}{}$ for all positions	tive
	integers $n$ .			K-1			
(b)	Using (a), evaluate		+1).				
		k=3	• • •	·		(6 ma	ırks)
	•		`				
	, ,						
	,	-		*			
				<del>.</del>			<u> </u>
		3 tu-					
						·	
	,					,, <u>,</u>	
	<del>,</del>			.,,-			
	·•	<u></u>	<u> </u>		<u></u>	ν <del>.,</del>	
					<del>-</del>		
							<del></del>
				·	<u> </u>		

Answers written in the margins will not be marked.

Please stick the barcode label here.

Answers written in the margins will not be marked.

Please stick the barcode label here.

	volume of the solid of revolution generated by revolving R about-the x-a	(8 marks)
_	· · · · · · · · · · · · · · · · · · ·	
	· · · · · · · · · · · · · · · · · · ·	
_		
•	,	
_		
-		
-		
-		
-		
-		
-		
-	<u> </u>	
-	· ·	
	·	
		· 
	,	
		74

Answers written in the margins will not be marked.

Please stick the barcode label here.

- 1**2016** 8. I Let n be a positive integer.
  - (a) Define  $M = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ . Evaluate
    - (i)  $M^2$ ,
    - (ii)  $M^n$ ,
    - (iii)  $\left(\begin{array}{cc} 1 & 0 \\ 2 & 1 \end{array}\right)^n$ .
  - (b) Evaluate
    - (i)

(8 marks)

Answers written in the margins will not be marked.

marke
þ
not
will
margins
the
.⊑
written
Answers

	We $y = f(x)$ by C. It is given that $P(-2, 9)$ is a stationary point of C. Find p and q.	(3 marks	
	Is P a minimum point of C? Explain your answer.	(2 marks)	
(c)	Find the relative extreme points(s) of $C$ .	(2 marks	
	Find the point(s) of inflexion of $C$ .	(2 marks	
	Let $L$ be the tangent to $C$ at $P$ . Find the area of the region bounded by $C$ and $L$ .		
(0)	Lot 2 be the tangent to 6 at 111 me the them of the 19	`	
	,	•	
		<del></del>	
		<u> </u>	
		<del>.</del>	
_		<del>,</del>	

· · · · · · · · · · · · · · · · · · ·

- 10. (a) Suppose  $\frac{\pi}{2} < k < \pi$ .
  - (i) Show that  $\int_{\frac{\pi}{4}}^{k-\frac{\pi}{4}} \ln \sin(k-x) \, dx = \int_{\frac{\pi}{4}}^{k-\frac{\pi}{4}} \ln \sin x \, dx$ .
  - (ii) Hence show that  $\int_{\frac{\pi}{4}}^{k-\frac{\pi}{4}} \ln(\sin k \cot x \cos k) dx = 0.$

(6 marks)

- (b) (i) Show that  $\int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{x \csc^2 x}{\cot x + 1} dx = \frac{\pi \ln 2}{4} + \int_{-\frac{\pi}{4}}^{\frac{\pi}{2}} \ln(\cot x + 1) dx.$ 
  - (ii) Using the results of (a)(ii) and (b)(i), evaluate  $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{x \csc^2 x}{\cot x + 1} dx$ .

(6 marks)

Answers written in the margins will not be marked.

4			
	·		
		<u>.</u>	
*			
	· · · · · · · · · · · · · · · · · · ·		
		<b>.</b>	
		r .	
		÷	
	·		
		······································	
	· ·		

Answers written in the margins will not be marked.

EP(M2) MOCK 1-16

		•						·		
		,				14	,			
_			<del></del> *			-				
				,		-				
				-						
			,						_	
_	-				_					-
_										_ <del>-</del> *
_	. 10				-					
_	·			***			,	,		
		-	. <del></del>		,				-	•
_	,		· ·		<u> </u>					•
-	A.		•					·		
-	.,	,						· .		
_						-	,			
_		· · · · · · · · · · · · · · · · · · ·			-					
-		<del></del> -					<del>•</del>	· ·		
_					·					
_	٠.				,	,			,	
-				·	-					
-			,	41						
-								· · ·		
-	<u>-</u>	···					,			
-		•	-			**				
-	<del>.</del>	_	<del></del>	,						

11. (a) Consider the system of linear equations in real variables x, y, z

(E): 
$$\begin{cases} x - y + z = 2\\ 2x - y + pz = q\\ 4x + (2-p)y + (4p-5)z = q-8 \end{cases}$$

where p and q are real numbers.

- (i) Assume that (E) has a unique solution.
  - (1) Prove that  $p \neq 1$  and  $p \neq 3$ .
  - (2) Solve (E).
- (ii) Assume that p = 3 and (E) is consistent.
  - (1) Find q.
  - (2) Solve (E).

(9 marks)

(b) Is there a real solution of the system of linear equations

$$\begin{cases} x - y + z = 2\\ 8x - 4y + 12z = -8\\ 8x - 2y + 14z = -20 \end{cases}$$

satisfying xy = 3z? Explain your answer.


(3 marks)

Answers written in the margins will not be marked

			_		
				3	:
	·				***************************************
				<u>.</u>	
			:		
,					
				<del></del>	
			·		
	· · · · · · · · · · · · · · · · · · ·		<b>4.</b> ·	<u>-</u>	
			<del></del>		
	•				
		<u>_</u>			
			·		
				<del></del>	1
			_		

				_			
			· · · ·				
				<del></del>			
							_
•							
,				-			
		· · · · · · · · · · · · · · · · · · ·		•	**		
	·				·		
			**	<u></u> .		<u>.</u>	
				<u>.</u>	<u>.</u>		
				<u> </u>			
			1-11				
			.,				
						1	
				•		·	
			*	<u> </u>	<del>.</del>		
				<u> </u>			
			.,				
		<u> </u>	<u>.</u>				
				±u.		·	

	`	
ļ		
ked.		
be mai		
Il not		
gins wi		
e marg		
n in th		
writte		
Answers written in the margins will not be marked.		
Y)		
	ı	
	1	



In Figure 1, **a** and **b** are unit vectors with  $\mathbf{a} \cdot \mathbf{b} = \frac{1}{3}$ . Let  $\overrightarrow{OP} = 4\mathbf{a}$ ,  $\overrightarrow{OQ} = 2\mathbf{b}$  and  $\overrightarrow{OC} = r\mathbf{a} + s\mathbf{b}$ , where C is the circumcentre of  $\triangle OPQ$ , r and s are constants.

(a) By considering  $\overrightarrow{OP} \cdot \overrightarrow{CM}$ , where M is the mid-point of OP, show that 3r + s = 6.

(3 marks)

(b) Find the values of r and s.

(3 marks)

(c)

Answers written in the margins will not be marked.




Figure 2

In Figure 2, H is the orthocentre of  $\triangle OPQ$  and N is the mid-point of PQ. Let OH : CN = k : 1, where k is a constant. By expressing  $\overrightarrow{OH}$  in terms of k, k and k, find k in terms of k and k. (7 marks)