

MATHEMATICS Extended Part
Module 2 (Algebra and Calculus)
Question-Answer Book

This paper must be answered in English

©香港考試及評核局 保留版權
Hong Kong Examinations and Assessment Authority
All Rights Reserved 2017

Candidate Number

--	--	--	--	--	--	--	--	--



FORMULAS FOR REFERENCE

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$	$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$
$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$	$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$
$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$	$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$
$2 \sin A \cos B = \sin(A+B) + \sin(A-B)$	$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$
$2 \cos A \cos B = \cos(A+B) + \cos(A-B)$	
$2 \sin A \sin B = \cos(A-B) - \cos(A+B)$	

SECTION A (50 marks)

1. Find $\frac{d}{d\theta} \sec 6\theta$ from first principles.

(5 marks)

$$\begin{aligned}
 \frac{d}{d\theta} \sec 6\theta &= \lim_{h \rightarrow 0} \frac{\sec(6(\theta+h)) - \sec 6\theta}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sec(6\theta+6h) - \sec 6\theta}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{\cos(6\theta+6h)} - \frac{1}{\cos 6\theta}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos 6\theta - \cos(6\theta+6h)}{h \cos(6\theta+6h) \cos 6\theta} \\
 &= \lim_{h \rightarrow 0} \frac{1}{h} \cdot \frac{-2 \sin(6\theta+3h) \sin(-3h)}{\frac{1}{2}(\cos 12\theta + \cos 6h)} \\
 &= \lim_{h \rightarrow 0} \frac{4}{h} \cdot \frac{\sin(6\theta+3h) \sin 3h}{2 \cos(6\theta+3h) \cos(6\theta-3h)} \\
 &= 2 \cdot 3 \lim_{h \rightarrow 0} \frac{\sin 3h}{3h} \lim_{h \rightarrow 0} \frac{\sin(6\theta+3h)}{\cos(6\theta+3h) \cos(6\theta-3h)} \\
 &= 6 \sec 6\theta \tan 6\theta
 \end{aligned}$$

Answers written in the margins will not be marked.

2. Let $(1+ax)^8 = \sum_{k=0}^8 \lambda_k x^k$ and $(b+x)^9 = \sum_{k=0}^9 \mu_k x^k$, where a and b are constants. It is given that $\lambda_2 : \mu_7 = 7 : 4$ and $\lambda_1 + \mu_8 + 6 = 0$. Find a . (5 marks)

For $(1+ax)^8$ general term $= C_8^d (ax)^d = C_8^d a^d x^d$

$$\lambda_2 = C_8^2 a^2 \quad \lambda_1 = C_8^1 a$$

For $(b+x)^9$ general term $= C_9^f b^{9-f} x^f$

$$\mu_7 = C_9^7 b^{9-7} = C_9^7 b^2$$

$$\mu_8 = C_9^8 b^{9-8} = C_9^8 b$$

$$\frac{\lambda_2}{\mu_7} = \frac{7}{4}$$

$$\frac{C_8^2 a^2}{C_9^7 b^2} = \frac{7}{4}$$

$$\frac{7a^2}{2b^2} = \frac{7}{4} \quad \frac{a}{\sqrt{2}b} = \frac{1}{2}$$

$$28a^2 = 14b^2 \quad (ii) \quad a = \frac{1}{2}(\sqrt{2}b) \quad (i)$$

Put (i) into (ii)

$$28a^2 = 14 \left(\frac{1}{2}(\sqrt{2}b) \right)^2$$

$$28a^2 = 14 \left(\frac{1}{4} \cdot 2b^2 \right)$$

$$28a^2 = 14 \left(\frac{1}{2}b^2 \right)$$

$$28a^2 = 7b^2$$

$$4a^2 = b^2$$

$$a = \frac{b}{2}$$

$$a = \frac{-3\sqrt{2}}{4+4\sqrt{2}} = \frac{-3\sqrt{2}(4-4\sqrt{2})}{4(4-16)} = \frac{-12\sqrt{2}+24}{-48} = \frac{12\sqrt{2}-24}{48}$$

3. P is a point lying on AB such that $AP:PB=3:2$. Let $\vec{OA}=\mathbf{a}$ and $\vec{OB}=\mathbf{b}$, where O is the origin.

(a) Express \vec{OP} in terms of \mathbf{a} and \mathbf{b} .

(b) It is given that $|\mathbf{a}|=45$, $|\mathbf{b}|=20$ and $\cos\angle AOB=\frac{1}{4}$. Find

(i) $\mathbf{a} \cdot \mathbf{b}$,

(ii) $|\vec{OP}|$.

3a) $\vec{OP} = \frac{3\vec{b} + 2\vec{a}}{3+2} = \frac{2}{5}\vec{a} + \frac{3}{5}\vec{b}$ (5 marks)

b i) $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \angle AOB$
 $\vec{a} \cdot \vec{b} = (45)(20)(\frac{1}{4})$
 $\vec{a} \cdot \vec{b} = 225$

b ii) $|\vec{OP}|^2 = (\frac{2}{5}\vec{a} + \frac{3}{5}\vec{b}) \cdot (\frac{2}{5}\vec{a} + \frac{3}{5}\vec{b})$
 $= \frac{4}{25}|\vec{a}|^2 + \frac{6}{25}\vec{a} \cdot \vec{b} + \frac{6}{25}\vec{b} \cdot \vec{a} + \frac{9}{25}|\vec{b}|^2$
 $= \frac{4}{25}(45)^2 + 2(\frac{6}{25})(225) + \frac{9}{25}(20)^2$
 $= 576$
 $|\vec{OP}| = \sqrt{576}$
 $= 24$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

4. (a) Using integration by parts, find $\int x^2 e^{-x} dx$.

(b) Find the area of the region bounded by the graph of $y = x^2 e^{-x}$, the x -axis and the straight line $x = 6$.

(6 marks)

$$\begin{aligned}
 \text{a) } \int x^2 e^{-x} dx &= -\int x^2 de^{-x} \\
 &= -x^2 e^{-x} + \int e^{-x} dx \\
 &= -x^2 e^{-x} + 2 \int x e^{-x} dx \\
 &= -x^2 e^{-x} - 2 \int x de^{-x} \\
 &= -x^2 e^{-x} - 2x e^{-x} + 2 \int e^{-x} dx \\
 &= -x^2 e^{-x} - 2x e^{-x} - 2e^{-x} + C
 \end{aligned}$$

$$\begin{aligned}
 \text{b) When } y=0, x^2 e^{-x} &= 0 \\
 x^2 &= 0 \text{ or } e^{-x} = 0 \text{ (neg.)} \\
 x &= 0
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{Required area} &= \int_0^6 x^2 e^{-x} dx \\
 &= \left[-x^2 e^{-x} - 2x e^{-x} - 2e^{-x} \right]_0^6 \\
 &= -36e^{-6} - 12e^{-6} - 2e^{-6} + 2 \\
 &= -\frac{50}{e^6} + 2
 \end{aligned}$$

5. Consider the following system of linear equations in real variables x, y, z

$$(E): \begin{cases} x + 2y - z = 11 \\ 3x + 8y - 11z = 49, \text{ where } h, k \in \mathbb{R} \\ 2x + 3y + hz = k \end{cases}$$

(a) Assume that (E) has a unique solution.

(i) Find the range of values of h .

(ii) Express z in terms of h and k .

(b) Assume that (E) has infinitely many solutions. Solve (E) .

5ai)

$$\Delta = \begin{vmatrix} 1 & 2 & -1 \\ 3 & 8 & -11 \\ 2 & 3 & h \end{vmatrix} = \begin{vmatrix} 1 & 2 & -1 \\ 0 & -2 & 8 \\ 0 & 1 & -2h \end{vmatrix}$$

$$= (-2)(-2h) - 8 = 4 + 2h - 8$$

$$= 2h - 4$$

$\therefore (E)$ has a unique solution

$$\therefore \Delta \neq 0$$

$$2h - 4 \neq 0$$

$$h \neq 2$$

aii)

$$\Delta_x = \begin{vmatrix} 11 & 2 & -1 \\ 49 & 8 & -11 \\ k & 3 & h \end{vmatrix} = \begin{vmatrix} -5 & 0 & 7 \\ 33 & 0 & -3-2h \\ k & 3 & h \end{vmatrix}$$

$$= -3[(-5)(-3-2h) - 7(33-2h)]$$

$$= -3(15 + 10h - 231 + 14h)$$

$$= -3(10h + 14h - 216)$$

$$= -6(5h + 7k - 108)$$

$$\Delta_y = \begin{vmatrix} 1 & 1 & -1 \\ 3 & 49 & -11 \\ 2 & k & h \end{vmatrix} = \begin{vmatrix} 1 & 1 & -1 \\ 0 & -16 & -22 \\ 0 & 22-k & 2-h \end{vmatrix}$$

$$= (-16)(-2-h) + 22(22-k)$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$\Delta y = 32 + 16h + 484 - 22k$$

$$J = (6h - 22k + 516)$$

$$= 2(8h - 11k + 258)$$

$$\Delta z = \begin{vmatrix} 1 & 2 & 11 \\ 3 & 8 & 49 \\ 2 & 3 & k \end{vmatrix} = \begin{vmatrix} 1 & 2 & 11 \\ 0 & -2 & -16 \\ 0 & 1 & 22-k \end{vmatrix}$$

$$= (-2)(22-k) + 16$$

$$= -44 + 2k + 16$$

$$= 2k - 28$$

$$= 2(k-14) \quad \therefore z = \frac{k-14}{h-2}$$

$$(x, y, z) = \left(\frac{-3(5h+7k-108)}{h-2}, \frac{8h+1k+258}{h-2}, \frac{k-14}{h-2} \right)$$

$$b) \begin{pmatrix} 1 & 2 & -1 & 11 \\ 3 & 8 & -1 & 49 \\ 2 & 3 & h & k \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & -1 & 11 \\ 0 & -2 & 8 & -16 \\ 0 & 1 & 2h & 22k \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 2 & -1 & 11 \\ 0 & 1 & -4 & 8 \\ 0 & 1 & 2h & 22k \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & -1 & 11 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 2+h & -14+k \end{pmatrix}$$

$\therefore (E)$ has infinitely many solutions

$$\therefore -2+h=0 \text{ and } -14+k=0$$

$$h=2 \text{ and } k=14$$

$$\begin{pmatrix} 1 & 2 & -1 & 11 \\ 3 & 8 & -1 & 49 \\ 2 & 3 & 2 & 14 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & -1 & 11 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

6. A container in the form of an inverted right circular cone is held vertically. The height and the base radius of the container are 20 cm and 15 cm respectively. Water is now poured into the container.

- (a) Let $A \text{ cm}^2$ be the wet curved surface area of the container and $h \text{ cm}$ be the depth of water in the container. Prove that $A = \frac{15}{16} \pi h^2$.
- (b) The depth of water in the container increases at a constant rate of $\frac{3}{\pi} \text{ cm/s}$. Find the rate of change of the wet curved surface area of the container when the volume of water in the container is $96\pi \text{ cm}^3$.

a) Let $a \text{ cm}$ be the radius of water surface (7 marks)

$$\begin{aligned} \frac{r}{h} &= \frac{15}{20} \\ r &= \frac{3}{4}h \\ A &= \pi \left(\frac{3}{4}h\right) \sqrt{1^2 + \left(\frac{3}{4}\right)^2} \\ &= \pi \left(\frac{3}{4}h\right) \sqrt{25/16} \\ &= \pi \left(\frac{3}{4}h\right) \left(\frac{5}{4}\right) \\ &= \frac{15}{16} \pi h^2 \end{aligned}$$

b) Let V be the volume of water.

$$\begin{aligned} V &= \frac{1}{3} \pi \left(\frac{3}{4}h\right)^2 h \\ &= \frac{1}{3} \pi \left(\frac{9}{16}h^2\right) h \\ &= \frac{9}{48} h^3 \pi \end{aligned}$$

$$\begin{aligned} \text{When } V &= 96\pi, \quad 96\pi = \frac{9}{48} h^3 \pi \\ h^3 &= 512 \\ h &= 8 \end{aligned}$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$A = \frac{15}{16} \pi h^2$$

$$\frac{dA}{dt} = \left(\frac{15}{16}\right)(2h)(\pi) \frac{dh}{dt}$$

$$\frac{dA}{dt} = \left(\frac{15}{8}\right)(8)(\pi) \left(-\frac{3}{\pi}\right)$$

$$\frac{dA}{dt} = -45$$

∴ Required rate of change is
 $45 \text{ cm}^2/\text{s}$

7. (a) Prove that $\sin 3x = 3\sin x - 4\sin^3 x$.

(b) Let $\frac{\pi}{4} < x < \frac{\pi}{2}$.

(i) Prove that $\frac{\sin 3\left(x - \frac{\pi}{4}\right)}{\sin\left(x - \frac{\pi}{4}\right)} = \frac{\cos 3x + \sin 3x}{\cos x - \sin x}$.

(ii) Solve the equation $\frac{\cos 3x + \sin 3x}{\cos x - \sin x} = 2$.

(8 marks)

7(a) $\sin 3x = \sin(2x + x)$
 $= \sin 2x \cos x + \cos 2x \sin x$
 $= 2\sin x \cos^2 x + (1 - 2\sin^2 x) \sin x$
 $= 2\sin x (1 - \sin^2 x) + \sin x - 2\sin^3 x$
 $= 2\sin x - 2\sin^3 x + \sin x - 2\sin^3 x$
 $= 3\sin x - 4\sin^3 x$

7(b) $\frac{\sin 3\left(x - \frac{\pi}{4}\right)}{\sin\left(x - \frac{\pi}{4}\right)}$
 $= \frac{\sin 3x \cos \frac{3\pi}{4} - \cos 3x \sin \frac{3\pi}{4}}{\sin x \cos \frac{\pi}{4} - \cos x \sin \frac{\pi}{4}}$
 $= \frac{-\frac{\sqrt{2}}{2} \sin 3x - \frac{\sqrt{2}}{2} \cos 3x}{\frac{\sqrt{2}}{2} (\cos x - \sin x)}$
 $= \frac{-\frac{\sqrt{2}}{2} (\sin 3x + \cos 3x)}{\frac{\sqrt{2}}{2} (\cos x - \sin x)}$
 $= \frac{-(\sin 3x + \cos 3x)}{(\cos x - \sin x)}$
 $= \frac{\cos 3x + \sin 3x}{\cos x - \sin x}$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$\begin{aligned}
 \text{7bii)} \quad & \frac{\cos 3x + \sin 3x}{\cos x - \sin x} = 2 \\
 & \frac{\sin 3(x - \frac{\pi}{4})}{\sin(x - \frac{\pi}{4})} = 2 \\
 & \sin 3(x - \frac{\pi}{4}) = 2 \sin(x - \frac{\pi}{4}) \\
 & 3 \sin(x - \frac{\pi}{4}) - 4 \sin^3(x - \frac{\pi}{4}) = 2 \sin(x - \frac{\pi}{4}) \\
 & \sin(x - \frac{\pi}{4}) = 0 \text{ or } 3 - 4 \sin^2(x - \frac{\pi}{4}) = 2 \\
 & \sin(x - \frac{\pi}{4}) = 0 \text{ or } 4 \sin^2(x - \frac{\pi}{4}) = 1 \\
 & \text{(rej.) } \sin(x - \frac{\pi}{4}) = 0 \text{ or } \sin(x - \frac{\pi}{4}) = \pm \frac{1}{2} \\
 & \quad \quad \quad x - \frac{\pi}{4} = \frac{\pi}{6} \quad (\sin(x - \frac{\pi}{4}) = -\frac{1}{2} \text{ is rej.}) \\
 & \quad \quad \quad x = \frac{5\pi}{12}
 \end{aligned}$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

8. Let $f(x)$ be a continuous function defined on \mathbf{R}^+ , where \mathbf{R}^+ is the set of positive real numbers. Denote the curve $y=f(x)$ by Γ . It is given that Γ passes through the point $P(e^3, 7)$ and $f'(x) = \frac{1}{x} \ln x^2$ for all $x > 0$. Find

(a) the equation of the tangent to Γ at P ,

(b) the equation of Γ ,

(c) the point(s) of inflexion of Γ .

8a) $f'(e^3) = \frac{1}{e^3} \ln e^{3(2)} = \frac{1}{e^3} \ln e^6 = \frac{6}{e^3}$ (8 marks)

Equation of the tangent to Γ at P :

$$y - 7 = \frac{6}{e^3} (x - e^3)$$

$$ye^3 - 7e^3 = 6x - 6e^3$$

$$ye^3 = 6x + e^3$$

$$y = \frac{6}{e^3} x + 1$$

b) $f(x) = \int \frac{1}{x} \ln x^2 dx$
 $= 2 \int \frac{1}{x} \ln x dx$
 $= 2 \int \ln x d \ln x$
 $= 2 \left(\frac{1}{2} (\ln x)^2 \right) + C$
 $= (\ln x)^2 + C$

Put $P(e^3, 7)$ into $y = f(x)$,

$$7 = (\ln e^3)^2 + C$$

$$7 = 9 + C$$

$$C = -2$$

\therefore Equation of Γ :

$$f(x) = (\ln x)^2 - 2$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$8c) f'(x) = \frac{1}{x} \ln x^2$$

$$f''(x) = -\frac{1}{x^2} \ln x^2 + \left(\frac{1}{x}\right)\left(\frac{2x}{x^2}\right)$$

$$= -\frac{1}{x^2} \ln x^2 + \frac{2}{x^2}$$

$$f''(x) = 0$$

$$-\frac{1}{x^2} \ln x^2 + \frac{2}{x^2} = 0$$

$$-\ln x^2 + 2 = 0$$

$$\ln x^2 = 2$$

$$\ln x^2 = \ln e^2$$

$$x = e$$

x	$x < 0$	$x = 0$	$0 < x < e$	$x = e$	$x > e$
$f''(x)$	-	X	+	0	-

\therefore Point of inflexion
 $= (e, -1)$

SECTION B (50 marks)

9. Define $f(x) = \frac{x^2 - 5x}{x + 4}$ for all $x \neq -4$. Denote the graph of $y = f(x)$ by G .

(a) Find the asymptote(s) of G . (3 marks)

(b) Find $f'(x)$. (2 marks)

(c) Find the maximum point(s) and the minimum point(s) of G . (4 marks)

(d) Let R be the region bounded by G and the x -axis. Find the volume of the solid of revolution generated by revolving R about the x -axis. (4 marks)

9a) $f(x) = \frac{x^2 - 5x}{x + 4}$

$$= x - 9 + \frac{36}{x + 4}$$

Oblique asymptote: $y = x - 9$

Vertical asymptote: $x = -4$

b) $f'(x) = 1 - \frac{36}{(x + 4)^2}$
 $(f(x) = x - 9 + \frac{36}{x + 4})$

c) $f'(x) = 0$

$$1 - \frac{36}{(x + 4)^2} = 0$$

$$(x + 4)^2 - 36 = 0$$

$$(x + 4)^2 = 36$$

$$x + 4 = 6 \text{ or } x + 4 = -6$$

$$x = 2 \text{ or } x = -10$$

x	$x < -10$	$x = -10$	$-10 < x < -4$	$x = -4$	$-4 < x < 2$	$x = 2$	$x > 2$
$f'(x)$	+	0	-	X	-	0	+
∴ Maximum point = $(-10, -25)$							
Minimum point = $(2, -1)$							

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

9d) When $g=0$,

$$\frac{x^2-5x}{x+4} = 0$$

$$x^2-5x=0$$

$$x(x-5)=0$$

$$x=0 \text{ or } 5$$

Required Volume

$$= \pi \int_0^5 \left(x - 9 + \frac{36}{x+4} \right)^2 dx$$

$$= \pi \int_0^5 \left(x^2 - 18x + 81 + \frac{72(x-9)}{x+4} + \frac{1296}{(x+4)^2} \right) dx$$

$$= \pi \int_0^5 \left(x^2 - 18x + 81 + \frac{72x}{x+4} - \frac{648}{x+4} + \frac{1296}{(x+4)^2} \right) dx$$

$$= \pi \int_0^5 \left(x^2 - 18x + 81 + 72 - \frac{936}{x+4} + \frac{1296}{(x+4)^2} \right) dx$$

$$= \pi \left[\frac{1}{3}x^3 - 9x^2 + 153x - 936 \ln|x+4| - \frac{1296}{x+4} \right]_0^5$$

$$= \pi \left(\frac{125}{3} - 750 + 765 - 936 \ln 9 - 144 + 936 \ln 4 + \frac{1296}{4} \right)$$

$$= \pi \left(\frac{710}{3} + 936 \ln \frac{4}{9} \right)$$

$$= \pi \left(\frac{710}{3} + 1872 \ln \frac{2}{3} \right)$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

A large rectangular area with horizontal ruling lines, intended for writing answers.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. ABC is a triangle. D is the mid-point of AC . E is a point lying on BC such that $BE:EC = 1:r$. AB produced and DE produced meet at the point F . It is given that $DE:EF = 1:10$. Let $\vec{OA} = 2\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}$, $\vec{OB} = 4\mathbf{i} + 4\mathbf{j} - \mathbf{k}$ and $\vec{OC} = 8\mathbf{i} - 3\mathbf{j} - 2\mathbf{k}$, where O is the origin.

(a) By expressing \vec{AE} and \vec{AF} in terms of r , find r . (4 marks)

(b) (i) Find $\vec{AD} \cdot \vec{DE}$.

(ii) Are B , D , C and F concyclic? Explain your answer.

(5 marks)

(c) Let $\vec{OP} = 3\mathbf{i} + 10\mathbf{j} - 4\mathbf{k}$. Denote the circumcentre of $\triangle BCF$ by Q . Find the volume of the tetrahedron $ABPQ$. (3 marks)

(10a) $\vec{AE} = \frac{\vec{AC} + r\vec{AB}}{1+r}$

$$= \frac{(8\mathbf{i} - 3\mathbf{j} - 2\mathbf{k}) - (2\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}) + r(4\mathbf{i} + 4\mathbf{j} - \mathbf{k})}{1+r}$$

$$= \frac{6\mathbf{i} - 6\mathbf{j} + r(2\mathbf{i} + 2\mathbf{j} + \mathbf{k})}{1+r}$$

$$= \frac{2(r+3)}{1+r}\mathbf{i} + \frac{2(r-3)}{1+r}\mathbf{j} + \frac{r}{1+r}\mathbf{k}$$

$\vec{AF} =$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$\begin{aligned} 100) \quad & \vec{AB} \times \vec{AP} \\ & = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 2 & 1 & 1 \\ 1 & 7 & 2 \end{vmatrix} = -9\vec{i} + 11\vec{j} + 13\vec{k} \\ & \vec{AQ} \cdot \vec{AB} \times \vec{AP} \\ & = \end{aligned}$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

A large rectangular area with horizontal lines for writing answers.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

11. (a) Using $\tan^{-1} \sqrt{2} - \tan^{-1} \left(\frac{\sqrt{2}}{2} \right) = \tan^{-1} \left(\frac{\sqrt{2}}{4} \right)$, evaluate $\int_0^1 \frac{1}{x^2 + 2x + 3} dx$. (3 marks)

(b) (i) Let $0 \leq \theta \leq \frac{\pi}{4}$. Prove that $\frac{2 \tan \theta}{1 + \tan^2 \theta} = \sin 2\theta$ and $\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \cos 2\theta$.

(ii) Using the substitution $t = \tan \theta$, evaluate $\int_0^{\frac{\pi}{4}} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta$. (5 marks)

(c) Prove that $\int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta = \int_0^{\frac{\pi}{4}} \frac{\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta$. (2 marks)

(d) Evaluate $\int_0^{\frac{\pi}{4}} \frac{8 \sin 2\theta + 9}{\sin 2\theta + \cos 2\theta + 2} d\theta$. (3 marks)

11(a) $\int_0^1 \frac{1}{x^2 + 2x + 3} dx$
 $= \int_0^1 \frac{1}{(x+1)^2 + 2} dx$
 $= \int_{\tan^{-1}(\frac{\sqrt{2}}{2})}^{\tan^{-1}(\sqrt{2})} \frac{1}{(\sqrt{2} \tan \theta + 1)^2 + 2} d(\sqrt{2} \tan \theta + 1)$
 $= \sqrt{2} \int_{\tan^{-1}(\frac{\sqrt{2}}{2})}^{\tan^{-1}(\sqrt{2})} \frac{\sec^2 \theta}{2 \tan^2 \theta + 2} d\theta$
 $= 2\sqrt{2} \int_{\tan^{-1}(\frac{\sqrt{2}}{2})}^{\tan^{-1}(\sqrt{2})} \frac{\sec^2 \theta}{\sec^2 \theta} d\theta$
 $= 2\sqrt{2} \int_{\tan^{-1}(\frac{\sqrt{2}}{2})}^{\tan^{-1}(\sqrt{2})} d\theta$
 $= 2\sqrt{2} \left[\theta \right]_{\tan^{-1}(\frac{\sqrt{2}}{2})}^{\tan^{-1}(\sqrt{2})}$
 $= 2\sqrt{2} \left(\tan^{-1}(\sqrt{2}) - \tan^{-1}(\frac{\sqrt{2}}{2}) \right)$
 $= 2\sqrt{2} \tan^{-1} \left(\frac{\sqrt{2}}{4} \right)$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$\begin{aligned}
 \text{11b i)} \quad \frac{2 \tan \theta}{1 + \tan^2 \theta} &= \frac{2 \tan \theta}{\sec^2 \theta} \\
 &= \frac{2 \sin \theta \cdot \cos^2 \theta}{\cos \theta} \\
 &= 2 \sin \theta \cos \theta \\
 &= \sin 2\theta
 \end{aligned}$$

$$\begin{aligned}
 \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} &= \frac{1 - (\sec^2 \theta - 1)}{\sec^2 \theta} \\
 &= \frac{2 - \sec^2 \theta}{\sec^2 \theta} \\
 &= 2 \cos^2 \theta - 1 \\
 &= \cos 2\theta
 \end{aligned}$$

$$\begin{aligned}
 \text{bii)} \quad &\int_{-\pi/4}^{\pi/4} \frac{1}{\sin 2\theta + \cos 2\theta + 2} d\theta \\
 &= \int_{-\pi/4}^{\pi/4} \frac{1}{\frac{2 \tan \theta}{1 + \tan^2 \theta} + \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} + 2} d\theta \\
 &= \int_{-\pi/4}^{\pi/4} \frac{1}{\frac{2 \tan \theta + 1 - \tan^2 \theta + 2(1 + \tan^2 \theta)}{\sec^2 \theta}} d\theta \\
 &= \int_{-\pi/4}^{\pi/4} \frac{\sec^2 \theta}{2 \tan \theta + 1 - \tan^2 \theta + 2 + 2 \tan^2 \theta} d\theta \\
 &= \int_{-\pi/4}^{\pi/4} \frac{1}{\tan^2 \theta + 2 \tan \theta + 3} d \tan \theta \\
 &= \int_0^1 \frac{1}{t^2 + 2t + 3} dt \\
 &= 2\sqrt{2} \tan^{-1} \left(\frac{\sqrt{2}}{4} \right)
 \end{aligned}$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$\begin{aligned}
 & 11c) \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sqrt{\sin 2\theta + \cos 2\theta + 2}} d\theta + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\cos 2\theta + 1}{\sqrt{\sin 2\theta + \cos 2\theta + 2}} d\theta \\
 &= \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + \cos 2\theta + 1}{\sqrt{\sin 2\theta + \cos 2\theta + 2}} d\theta \\
 &= \int_0^{\frac{\pi}{4}} \frac{\sin 2\theta + 1}{\sqrt{\sin 2\theta + \cos 2\theta + 2}} d\theta
 \end{aligned}$$

$$\begin{aligned}
 & d) \int_0^{\frac{\pi}{4}} \frac{8\sin 2\theta + 9}{\sin 2\theta + \cos 2\theta + 2} d\theta \\
 &= \int_0^{\frac{\pi}{4}} \left(8 \frac{\sin 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} + \frac{9}{\sin 2\theta + \cos 2\theta + 2} \right) d\theta \\
 &= 8 \left[\theta \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \frac{8\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta + 9(2\sqrt{2}) \tan^{-1}\left(\frac{\sqrt{2}}{4}\right) \\
 &= 2\pi + 18\sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}}{4}\right) - \int_0^{\frac{\pi}{4}} \frac{8\cos 2\theta + 1}{\sin 2\theta + \cos 2\theta + 2} d\theta
 \end{aligned}$$

Answers written in the margins will not be marked.

A large rectangular area with horizontal ruling lines for writing answers.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

12. Let $A = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$. Denote the 2×2 identity matrix by I .

(a) Using mathematical induction, prove that $A^n = 3^n I + 3^{n-1} n \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ for all positive integers n .

(4 marks)

(b) Let $B = \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix}$.

(i) Define $P = \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}$. Evaluate $P^{-1}BP$.

(ii) Prove that $B^n = 3^n I + 3^{n-1} n \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}$ for any positive integer n .

(iii) Does there exist a positive integer m such that $|A^m - B^m| = 4m^2$? Explain your answer.

(8 marks)

Q) Let $P(n): A^n = 3^n I + 3^{n-1} n \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$

For $n=1$

LHS = $A = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$

RHS = $3I + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$

$P(1)$ is true.

Assume $P(k)$ is true. i.e. $A^k = 3^k I + 3^{k-1} k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$

For $n=k+1$

$A^{k+1} = A^k A$

$$= [3^k I + 3^{k-1} k \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}] \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 3^k & 0 \\ 0 & 3^k \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix} + \begin{pmatrix} 0 & 3^{k-1} k \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 3^{k+1} & 0 \\ 0 & 3^{k+1} \end{pmatrix} + \begin{pmatrix} 0 & 3^k + 3^k k \\ 0 & 0 \end{pmatrix}$$

$$= 3^{k+1} I + \begin{pmatrix} 0 & 3^{k-1} + 3^{k-1} k \\ 0 & 0 \end{pmatrix}$$

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

$$= 3^{k+1}I + 3^k(k+1)\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$\therefore P(k+1)$ is true.

$\therefore P(n)$ is true for all positive integers n .

$$\text{bii) } P^{-1} = \frac{1}{\det P} \begin{pmatrix} -1 & -2 \\ 0 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix}$$

$$P^{-1}BP$$

$$= \begin{pmatrix} -1 & 0 \\ -2 & -1 \end{pmatrix} \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} -5 & -1 \\ -6 & -3 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

$$\text{bii) let } Q(n): B^n = 3^n I + 3^{n-1}n \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}$$

For $n=1$,

$$\text{LHS} = B = \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix}$$

$$\text{RHS} = 3I + \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} = \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix}$$

$\therefore Q(1)$ is true.

Assume $Q(k)$ is true. i.e. $B^k = 3^k I + 3^{k-1}k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}$

For $n=k+1$,

$$B^{k+1} = B^k B$$

$$= \left[3^k I + 3^{k-1}k \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix} \right] \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix}$$

$$\begin{aligned}
 P_{k+1} &= (3^k + 2k(3^{k-1}) - 3^{k-1}k) \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \\
 &= (-4k)3^{k-1} - 3^k + 2k(3^k) \begin{pmatrix} 5 & 1 \\ -4 & 1 \end{pmatrix} \\
 &= (5(3^k) + 10k(3^{k-1}) - 4(3^k) - 3^k + 3)3^{k-1} \\
 &= (-20k3^{k-1} - 4(3^k) + 8k(3^{k-1}) - 4k)3^{k-1} + 3^k - 2k(3^{k-1}) \\
 &= (5(3^k) + 6k(3^{k-1}) - 4(3^k) - 6k(3^{k-1}) + 3^k) \\
 &= (12k3^{k-1} - 4(3^k) + 3^k) \\
 &= 3^{k+1} + 3^k(k+1) \begin{pmatrix} 2 & 1 \\ -4 & -2 \end{pmatrix}
 \end{aligned}$$

$Q(k+1)$ is true

$Q(n)$ is true for any positive integers n

12biii)

END OF PAPER

考生編號 Candidate Number

試題編號 Question No.

1	2	3	4	5	6	7	8	9	10	11	12	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
13	14	15	16	17	18	19	20	21	22	23	24	≥25

1. 每題另起新頁作答。

Start each question on a new page.

2. 補充答題紙不可撕開使用。

Do not tear the supplementary answer sheet apart.

$$5b) \sim \begin{pmatrix} 1 & 0 & 7 & -5 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$(x, y, z) = (-7t - 5, 4t + 8, t) \text{ for all } t \in \mathbb{R}.$$

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

本頁積分 Page total

試題編號 Question No.

1 2 3 4 5 6 7 8 9 10 11 12

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

13 14 15 16 17 18 19 20 21 22 23 24 ≥ 25

1. 每題另起新頁作答。

Start each question on a new page.

2. 補充答題紙不可撕開使用。

Do not tear the supplementary answer sheet apart.

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

寫於邊界以外的答案，將不予評閱。

Answers written in the margins will not be marked.

本頁積分 Page total