HKDSE MATH EP

M2

HONG KONG EDUCATIONAL PUBLISHING COMPANY

HONG KONG DIPLOMA OF SECONDARY EDUCATION EXAMINATION

MATHEMATICS Extended Part

Module 2 (Algebra and Calculus) MOCK EXAM 3 Question-Answer Book

Time allowed: 2½ hours

This paper must be answered in English

INSTRUCTIONS

- 1. After the announcement of the start of the examination, you should first write your Candidate Number in the space provided on Page 1 and stick barcode labels in the spaces provided on Pages 1, 3, 5, 7, 9 and 11.
- 2. This paper consists of TWO sections, A and B.
- 3. Attempt ALL questions in this paper. Write your answers in the spaces provided in this Question-Answer Book. Do not write in the margins. Answers written in the margins will not be marked.
- 4. Graph paper and supplementary answer sheets will be supplied on request. Write your Candidate Number, mark the question number box and stick a barcode label on each sheet, and fasten them with string INSIDE this book.
- 5. Unless otherwise specified, all working must be clearly shown.
- 6. Unless otherwise specified, numerical answers must be exact.
- 7. In this paper, vectors may be represented by bold-type letters such as \mathbf{u} , but candidates are expected to use appropriate symbols such as \mathbf{u} in their working.
- 8. The diagrams in this paper are not necessarily drawn to scale.
- 9. No extra time will be given to candidates for sticking on the barcode labels or filling in the question number boxes after the 'Time is up' announcement.
- © 香港教育圖書公司 保留版權 Hong Kong Educational Publishing Company All Rights Reserved 2016

Please	stick	the	bar	code	la l	bel	her	e.
•	(
Candidate Number								

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$2\sin A\cos B = \sin (A+B) + \sin (A-B)$$

$$2\cos A\cos B = \cos (A+B) + \cos (A-B)$$

$$2 \sin A \sin B = \cos (A - B) - \cos (A + B)$$

$$\sin A + \sin B = 2\sin\frac{A+B}{2}\cos\frac{A-B}{2}$$

$$\sin A - \sin B = 2\cos\frac{A+B}{2}\sin\frac{A-B}{2}$$

$$\cos A + \cos B = 2\cos\frac{A+B}{2}\cos\frac{A-B}{2}$$

$$\cos A - \cos B = -2\sin\frac{A+B}{2}\sin\frac{A-B}{2}$$

SECTION A (50 marks)

2014

In the expansion of $(1 + 2x)^2(1 - x)^n$, the coefficient of x is -7.

- (a) Find the value of n.
- (b) Find the coefficient of x^2 .

(4 marks)

Answers written in the margins will not be marked.

				· .		
				-		
					•	
			-	4		
			•			
				*		
				•		
		•				
		•		٠,		
-		 				
	- -					
			- 			

Please stick the barcode label here.

· · · · · · · · · · · · · · · · · · ·	
·	· · · · · · · · · · · · · · · · · · ·
· · · · · · · · · · · · · · · · · · ·	
	.
· · · · · · · · · · · · · · · · · · ·	

Answers written in the margins will not be marked.

3.	(a)	Let $x = \cot \theta$. Show that $\frac{x^2 - 1}{x^2 + 1} = \cos 2\theta$.	
		Using (a), find the least value of $\frac{2(x+1)(x-1)}{x^2+1}$, where x is real.	(5 marks)
		:	
-			
:			,
			
	.—		
:			
			·
		1	

Please stick the barcode label here.

	Find $f'(x)$		t is a cons				,		
				0 for all	real values	of x , find	the value(s	s) of k .	
(-)	- J ()	y ()	, , ,				·		(5 mar
		-	•	**					
								÷	
									<u></u>
			-	···	•				
-		,					·		,
				•				,	
						•			-
	<u>-</u> .					. .			
	· · ·								
	<u>.</u>		·						 _
	4.14			γ	•			•	
•	_	<u>-</u>				-		<u>-</u>	400
								· · · · · · · · · · · · · · · · · · ·	
		• 1							
		ı		****					
		.					· <u>-</u>		
		,			· · · · · · · · · · · · · · · · · · ·				
					<u>_</u>				
<u>-</u>			,						

	Using integration by substitution, find $\int \frac{dx}{x^2 + 9}$.
(b)	At any point (x, y) on the curve Γ , the slope of the tangent to Γ is $\frac{x^2}{x^2 + 9}$. It is given that Γ passes through the point $(0, 6)$. Find the equation of Γ . (7 marks
,	
	
	· · · · · · · · · · · · · · · · · · ·
_	
<u>~</u>	

Please stick the barcode label here.

6. Let f(x) be a polynomial. Figure 1 shows a sketch of the curve y = f'(x), where $a \le x \le c$. The curve cuts the x-axis at the origin and (b, 0), where 0 < b < c.




Figure 1

- (a) Write down the x-coordinates of the maximum and minimum points of the curve $y \neq f(x)$ for a < x < c.
- (b) It is known that f(0) = 1 and the area of the shaded region R_2 as shown in Figure 1 is 6.
 - (i) By considering $\int_0^b f'(x) dx$, find the value of f(b).
 - (ii) If f(a) = f(c) and the areas of the shaded regions R_1 and R_3 as shown in Figure 1 are equal, find the value of f(a).

(7 marks)

	•	•				•		
 •			•		-17-		·	
_		,						
 1.	,				<u> </u>			
,	-			-				
			-					•

ਠ
Ž.
ਜ਼
Ħ
Ξ.
8
_
ਰ
П
_
ᇃ
will
S
.⊑
.ஹ
nargins
Ĕ
_
异
≒
□
-=
⊏
ritten
.=
7
>
S
<u>5</u>
≥
35
٦
7

If th	e system has	non-trivial s	solutions, fi	nd the po	ssible va	alues of	k.		(4	marks)
	·				·····			<u> </u>		
										
		<u></u>			<u></u>		_ .		<u> </u>	
				i		<u></u>				
					· <u>-</u>				_	
						>				
					-			-		
	·	- 								
	· .					***				
		-			•			1		
	, 					<u>.</u>			<u></u>	
		·			<u>.</u>		. , ,			
	· · · · · · · · · · · · · · · · · · ·		·			-		.,-		
									_ .	
					-					
									· .	
		<u> </u>								

Please stick the barcode label here.

2015

- 8. (a) Let A be a 3×3 matrix such that $A^T = -A$, where A^T is the transpose of A. Prove that |A| = 0.
 - (b) Let $M = \begin{pmatrix} 1 & x & y \\ -x & 1 & z \\ -y & -z & 1 \end{pmatrix}$, where x, y and z are real numbers. It is given that M is a non-

singular matrix. Denote the 3×3 identity matrix by I.

- (i) Using (a), or otherwise, prove that |I M| = 0.
- (ii) Someone claims that $I M^{-1}$ is a singular matrix. Do you agree? Explain your answer. (6 marks)

Answers written in the margins will not be marked.

	-				•
		,			
				49:	
					<u></u>
	<u></u>	- ALM, 12		· .	
<u></u>		<u>.</u> ,	· ***		
			,		

Answers written in the margins will not be marked.

 (a) Find the unit vector in the direction of OR. (b) Find ∠ROP correct to the nearest degree. 		
o) That Zhor confect to the homest degree.	·	(8 marks)
		,
	,	
	•	
·		
	 .	
		-
	·	
	<u> </u>	<u> </u>
		
		•
		<u> </u>
•		
		-
7		
		
•		
		·
		•

Answers written in the margins will not be marked.

Please stick the barcode label here.

10. Let
$$A = \begin{pmatrix} 0 & 0 & 1 \\ 4 & 4 & 4 \\ 5 & 4 & 5 \end{pmatrix}$$
 and $B = \begin{pmatrix} 4 & 4 & -4 \\ 0 & -5 & 4 \\ -4 & 0 & 0 \end{pmatrix}$ and $C = \begin{pmatrix} -3 & -4 & -4 \\ 5 & 6 & 5 \\ 0 & 0 & 1 \end{pmatrix}$

- (a) (i) Find AB and BA.
 - (ii) Using the result of (i), or otherwise, find A^{-1} .

(4 marks)

- (b) (i) Find ACA^{-1} .
 - (ii) Show that C is invertible.
 - (iii) Find $(D^{-1})^{2016}$, where $D = ACA^{-1}$. Hence, or otherwise, find $(C^{-1})^{2016}$.

(7 marks)

Answers written in the margins will not be marked.

Š	ļ
not	
WIII	
inswers written in the margins will not be	
the the	
Ξ	l
written	
Answers	
•	1




Figure 2

Let $\overrightarrow{OA} = -\mathbf{i} - \mathbf{k}$, $\overrightarrow{OB} = 3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$, $\overrightarrow{OC} = -\mathbf{i} + 3\mathbf{j} - \mathbf{k}$ and $\overrightarrow{OD} = \mathbf{i} + \mathbf{j} + 3\mathbf{k}$ (see Figure 2). Let P and Q be points on the straight lines AB and CD respectively such that AP : PB = r : (1 - r) and CQ : QD = s : (1 - s), where 0 < r < 1 and 0 < s < 1. Suppose that PQ is perpendicular to both AB and CD.

- (a) (i) Express \overrightarrow{PQ} in terms of r, s, i, j and k.
 - (ii) Hence find the shortest distance between the straight lines AB and CD.

(8 marks).

in the margins will not be marked.

Answers written

- (b) (i) Find $\overrightarrow{AB} \times \overrightarrow{AC}$.
 - (ii) Let R be the projection of D on the plane ABC. Find the coordinates of R by considering $\overrightarrow{AR} \cdot (\overrightarrow{AB} \times \overrightarrow{AC})$.

(5 marks)

- 12. Let $f(x) = \frac{x^4}{x^3 + 2}$, where $x \neq -2^{\frac{1}{3}}$.
 - . (a) Find the x- and y-intercept(s) of the graph of y = f(x).

(1 mark)

(b) Find f'(x) and prove that $f''(x) = \frac{12x^2(4-x^3)}{(x^3+2)^3}$.

(4 marks)


- (c) (i) For the graph of y = f(x), find all the extreme point(s) and point(s) of inflexion.
 - (ii) Find all the asymptote(s) of the graph of y = f(x).
 - (iii) Sketch the graph of y = f(x).

(8 marks)

		·			
					-
*		•			
		,			.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
		4.4.49		,**	
		4,00			
-			-	,	· ·
				-	
					<u>.</u>
•	<u>.</u>			 	
	· ,				,
	·				

Answers written in the margins will not be marked.

13. (a)




In Figure 3, the region enclosed by the curve $y = x^2$, the straight lines y = 4 and y = 4 + h(where $h \ge 0$) is revolved about the y-axis. Show the the volume of the solid of revolution is $\frac{\pi}{2}(h^2 + 8h)$.

. (2 marks)

(b)

Answers written in the margins will not be marked.



In Figure 4, an empty cup consists of two portions. The upper portion is in the shape of the solid described in (a) with height 8 cm. The lower portion is a frustum of a right circular cone. The height of the frustum is 4 cm and the radii of the bases of the frustum are 4 cm and 2 cm respectively. Milk is poured into the cup to a depth H cm at a rate of $8 \text{ cm}^3 \text{s}^{-1}$, where $0 \le H \le 12$.

Let $V \text{ cm}^3$ be the volume of milk in the cup.

- Find the rate of increase of the depth of milk when the depth is 7 cm.
- (ii) Show that $V = \frac{128\pi}{3} \left[1 \left(1 \frac{H}{8} \right)^3 \right]$ for $0 \le H \le 4$.
- (iii) After the cup is fully filled, suddenly it cracks at the bottom and the milk is leaking out. It is given that the rate of leaking is $\frac{\pi}{2}$ cm³s⁻¹. Find the rate of decrease of the depth of milk after 130 seconds of leaking. (Give the answer correct to 3 significant figures.)

(11 marks)