

Social Quantum Field Theory 31~40



7月 19, 2026

Chapter 31

Optimal Control and Decision Theory for Relational Fields

31.1 Introduction

The Digital Twin Social Field developed in the previous chapter provides a continuously updated mathematical representation of a complex relational system. Such a representation enables not only observation and prediction but also **decision-making**.

A central question naturally follows:

Given a relational field and a desired objective, how should interventions be designed to steer the system toward that objective while respecting its intrinsic dynamics?

Classical control theory addresses analogous questions for engineering systems by determining optimal control inputs. Social Quantum Field Theory (SQFT) extends this philosophy to relational systems by treating policies, institutional reforms, resource allocations, and strategic interventions as **control operators** acting upon an evolving relational field.

31.2 Controlled Relational Dynamics

Let

$$u(t)$$

denote an externally applied control.

The controlled Hamiltonian becomes

$$H(t) = H_0 + H_{\text{int}} + H_c(u(t)), \quad (31.1)$$

where

- H_0 represents intrinsic relational dynamics,
- H_{int} represents endogenous interactions,
- H_c represents external interventions.

Unlike previous chapters, the evolution of the relational field is now influenced by deliberate actions.

Definition 31.1 (Control Operator)

A **Control Operator** is an operator acting on the relational field that modifies its evolution according to an externally specified intervention.

Depending upon the application, control operators may represent

- public policy,
 - organizational restructuring,
 - investment strategies,
 - communication campaigns,
 - technological deployment,
 - institutional reform.
-

31.3 Controlled Lindblad Dynamics

For open relational systems,

the evolution equation becomes

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H(u), \rho] + \sum_k \left(L_k \rho L_k^\dagger - \frac{1}{2}\{L_k^\dagger L_k, \rho\} \right). \quad (31.2)$$

The control affects only the Hamiltonian,

while environmental influences continue to be represented by the Lindblad operators.

31.4 Objective Functional

Suppose the desired relational outcome is represented by the functional

$$J(u).$$

$J(u)$.

A general optimal control problem seeks

$$u^* = \operatorname{argmin} J(u). \quad (31.3)$$

A typical objective functional is

$$J(u) = \Phi(\rho(T)) + \int_0^T C(\rho, u) dt, \quad (31.4)$$

where

- Φ evaluates the final relational state,
- C represents the running cost of intervention.

Interpretation

Possible optimization goals include

- maximizing institutional resilience,
- minimizing systemic risk,
- reducing communication delay,
- increasing organizational coordination,
- improving innovation capacity.

31.5 Constraints

Real interventions are subject to practical limitations.

Examples include

$$u_{\min} \leq u(t) \leq u_{\max}, \quad (31.5)$$

budget constraints,

legal constraints,

ethical requirements,

resource limitations,

and political feasibility.

Accordingly,

optimal control is performed over the admissible set

$$u \in U. \quad (31.6)$$

$u \in U.$ (31.6)

31.6 Feedback Control

Rather than prescribing interventions in advance,
control may depend upon the observed relational state,

$$u(t) = K(\rho(t)). \quad (31.7)$$

Such **feedback control** continuously adapts to changes in the relational field.

Digital Twin Social Fields provide precisely the state estimates required for this approach.

Definition 31.2 (Adaptive Control)

An adaptive control policy continuously modifies interventions according to updated estimates of the relational field obtained through sequential observation and Bayesian inference.

31.7 Model Predictive Control

Modern engineering frequently employs **Model Predictive Control (MPC)**.

Within SQFT,

at each decision time

$$t_k,$$

one solves

$$\min_u J(u), \quad (31.8)$$

over a finite prediction horizon,

implements only the first control action,

updates the relational state,

and repeats the optimization.

This rolling-horizon strategy naturally integrates with Digital Twin Social Fields.

31.8 Multi-Objective Optimization

Many institutional decisions involve competing objectives.

For example,

one may simultaneously seek

- maximum economic efficiency,

- minimum systemic inequality,
- maximum institutional stability,
- minimum implementation cost.

The optimization becomes

$$\min(J_1, J_2, \dots, J_m). \quad (31.9)$$

Solutions are characterized by the **Pareto frontier** rather than by a single optimum.

31.9 Robust Control

Because parameter estimates remain uncertain,

control policies should remain effective across a range of plausible models.

Suppose

$$\Theta \sim P(\Theta|D).$$

The robust objective becomes

$$\min_u E_{\Theta}[J(u, \Theta)]. \quad (31.10)$$

This formulation naturally incorporates Bayesian uncertainty into decision-making.

31.10 Ethical Constraints

Unlike physical control systems,

relational systems involve autonomous individuals and institutions.

Therefore,

optimization should satisfy additional ethical constraints.

Examples include

- transparency,
- accountability,
- fairness,
- legal compliance,
- respect for individual rights.

These considerations are not merely normative additions but define admissible regions of the control space.

Theorem 31.1 (Existence of an Optimal Control)

Suppose

1. the admissible control set

$$U$$

is compact;

2. the objective functional

$$J(u)$$

is continuous;

3. the controlled SQFT evolution admits unique solutions for every admissible control.

Then there exists at least one optimal control

$$u^* \in U$$

that minimizes the objective functional.

Proof

Because the admissible control set is compact and the objective functional is continuous, the Weierstrass Extreme Value Theorem guarantees that the minimum of $J(u)$ is attained within U . The uniqueness of the system evolution ensures that the objective is well-defined for every admissible control. Therefore, at least one optimal control exists.

Corollary 31.1

When combined with Bayesian parameter estimation and Digital Twin Social Fields, optimal control provides a mathematically consistent framework for evaluating intervention strategies under uncertainty.

31.11 Discussion

Optimal control transforms SQFT from a predictive theory into a prescriptive one. Rather than merely forecasting the evolution of relational systems, the framework provides mathematical tools for identifying interventions that achieve specified objectives while respecting structural constraints and explicitly accounting for uncertainty.

Importantly, the optimal solution produced by the theory should be interpreted as a **decision-support recommendation** rather than an automatic prescription. Human judgment, domain expertise, legal requirements, and ethical considerations remain essential in selecting and implementing real-world interventions.

The next chapter develops the concept of **Inverse Social Field Theory**, addressing the inverse problem: given observed relational dynamics, how can the underlying Hamiltonian, interaction operators, and latent structural laws be reconstructed from data?

Chapter 32

Inverse Social Field Theory: Reconstructing Hidden Relational Dynamics

32.1 Introduction

The preceding chapter addressed the **forward problem** of Social Quantum Field Theory (SQFT): given a structural Hamiltonian and an initial relational state, predict the future evolution of the relational field.

Many scientific investigations, however, begin with the opposite question.

Rather than asking

What will happen given a known model?

we ask

What underlying relational dynamics generated the observed data?

This constitutes the **inverse problem**.

Inverse problems arise throughout science.

Examples include

- reconstructing Earth's interior from seismic waves,
- recovering medical images from projection measurements,
- inferring molecular structures from scattering experiments,
- estimating cosmological parameters from astronomical observations.

Analogously, SQFT seeks to reconstruct latent relational structures, interaction operators, and governing Hamiltonians from empirical observations.

32.2 Forward and Inverse Problems

The forward problem is expressed as

$$(H, \rho_0) \rightarrow D, \quad (32.1)$$

where

- H is the relational Hamiltonian,
- ρ_0 is the initial relational state,
- D denotes observable data.

The inverse problem reverses this mapping,

$$D \rightarrow (H, \rho_0, \Theta). \quad (32.2)$$

Unlike the forward problem,

the inverse problem is generally non-unique and often ill-posed, requiring additional assumptions or regularization.

Definition 32.1 (Inverse Social Field Problem)

An **Inverse Social Field Problem** seeks to infer the latent operators, parameters, and relational states that best explain a given collection of empirical observations under the governing equations of SQFT.

32.3 Observation Operator

Empirical measurements rarely observe the complete relational state.

Instead,

they are generated through an observation operator

$$O,$$

such that

$$D = O(\rho) + \varepsilon, \quad (32.3)$$

where

$$\varepsilon$$

represents measurement uncertainty.

Examples include

- sampled network interactions,
- aggregate economic indicators,
- communication statistics,
- organizational surveys,
- institutional performance measures.

32.4 Parameter Identification

Suppose the Hamiltonian depends upon parameters

$$\Theta.$$

The inverse problem seeks

$$\hat{\Theta} = \operatorname{argmin}_{\Theta} \|D - O(\rho(\Theta))\|^2. \quad (32.4)$$

This optimization compares simulated observations with empirical measurements.

Interpretation

Estimated parameters may correspond to

- interaction strengths,
 - institutional resilience,
 - communication efficiency,
 - adaptation rates,
 - diffusion coefficients.
-

32.5 State Reconstruction

In some applications,

the governing Hamiltonian is assumed known,

while the latent relational state remains unknown.

The inverse problem then becomes

$$\hat{\rho} = \operatorname{argmin}_{\rho} \|D - O(\rho)\|^2. \quad (32.5)$$

This procedure reconstructs the hidden relational configuration most consistent with available observations.

32.6 Hamiltonian Learning

A more ambitious objective is to reconstruct the Hamiltonian itself.

Suppose

$$H = \sum_i \theta_i H_i, \quad (32.6)$$

where

$$H_i$$

are basis operators.

Learning the Hamiltonian consists of estimating

$$\Theta = (\theta_1, \dots, \theta_n).$$

The resulting Hamiltonian provides a mechanistic description of the relational dynamics rather than merely a statistical fit.

32.7 Sparse Reconstruction

Complex relational systems often contain only a limited number of dominant interactions.

Accordingly,

one may solve

$$\min_{\Theta} \|D - O(\Theta)\|^2 + \lambda \|\Theta\|_1, \quad (32.7)$$

where

the L_1 penalty promotes sparse interaction structures.

Sparse reconstruction improves interpretability while reducing overfitting.

32.8 Identifiability

Not every parameter can necessarily be recovered from observations.

Definition 32.2 (Identifiability)

A parameter is **identifiable** if distinct parameter values generate distinguishable observational distributions under the assumed SQFT model.

When multiple parameter sets produce equivalent observations,

the inverse problem admits multiple solutions.

Additional observations,

experimental interventions,

or stronger prior information may then be required.

32.9 Bayesian Inverse Theory

Combining Chapter 28 with inverse reconstruction,

the posterior distribution becomes

$$P(H, \Theta, \rho|D) \propto P(D|H, \Theta, \rho)P(H, \Theta, \rho). \quad (32.8)$$

Rather than selecting a single solution,

Bayesian inversion quantifies uncertainty over all admissible reconstructions.

This probabilistic treatment is especially valuable when observations are noisy or incomplete.

32.10 Sequential Inverse Learning

As additional observations become available,

the reconstruction improves recursively.

Given new data

$$D_{n+1},$$

the posterior satisfies

$$P(\Theta|D_{1:n+1}) \propto P(D_{n+1}|\Theta)P(\Theta|D_{1:n}). \quad (32.9)$$

Consequently,

the inferred relational dynamics evolve continuously with accumulating evidence.

This recursive formulation naturally integrates with the Digital Twin framework introduced in Chapter 30.

32.11 Information-Theoretic Limits

The quality of inverse reconstruction depends fundamentally upon the information contained in the observations.

If measurements contain insufficient information,

no algorithm can uniquely recover the hidden relational dynamics.

The Fisher Information Matrix

$$I(\Theta) = E \left[\left(\frac{\partial \ln P(D|\Theta)}{\partial \Theta} \right) \left(\frac{\partial \ln P(D|\Theta)}{\partial \Theta} \right)^T \right] \quad (32.10)$$

quantifies the sensitivity of observations to parameter changes.

A nearly singular Fisher Information Matrix indicates poor identifiability and large estimation uncertainty.

Theorem 32.1 (Consistency of Bayesian Inverse Reconstruction)

Suppose

1. the SQFT model is correctly specified;
2. the true parameters are identifiable;
3. observational data increase without bound;
4. the prior assigns positive probability to the true parameter neighborhood.

Then the Bayesian posterior distribution converges toward the true relational Hamiltonian and parameter set.

Proof

The result follows from standard Bayesian consistency theory under identifiable statistical models. As additional observations accumulate, the likelihood increasingly dominates the posterior distribution, while the positivity of the prior near the true parameters ensures that the correct solution remains admissible. Consequently, posterior probability concentrates around the true Hamiltonian and associated parameters.

Corollary 32.1

When identifiability conditions are satisfied and sufficient observational information is available, Digital Twin Social Fields can progressively recover increasingly accurate representations of latent relational dynamics.

32.12 Discussion

Inverse Social Field Theory complements the predictive framework developed throughout previous chapters by addressing one of the central challenges of empirical science: inferring hidden mechanisms from observable outcomes. Within SQFT, inverse methods provide systematic procedures for reconstructing relational states, interaction operators, and structural Hamiltonians from incomplete and uncertain observations.

Together with Bayesian inference, tensor-network computation, and physics-informed artificial intelligence, inverse reconstruction closes the scientific loop between observation, explanation, prediction, and intervention. The framework therefore supports not only

forecasting but also the discovery of the latent principles governing complex relational systems.

The next chapter introduces **Variational Principles and the Principle of Least Action for Relational Fields**, establishing an action-based formulation that unifies the dynamical equations of SQFT within a single variational framework.

Chapter 33

Variational Principles and the Principle of Least Action for Relational Fields

33.1 Introduction

Previous chapters formulated the dynamics of Social Quantum Field Theory (SQFT) through Hamiltonians, density operators, Lindblad evolution, Bayesian inference, and optimal control. These formulations describe how relational systems evolve once the governing equations are known.

Modern theoretical physics, however, rests upon an even deeper principle.

Rather than postulating equations of motion directly, one begins with a single scalar quantity—the **action**—from which the equations of motion emerge through a variational principle.

Examples include

- Newtonian mechanics,
- classical electrodynamics,
- general relativity,
- quantum field theory,
- gauge theories,
- string theory.

This chapter develops an analogous formulation for SQFT, demonstrating that relational dynamics may be derived from a stationary-action principle rather than introduced axiomatically.

33.2 The Relational Action

Let

$$\Phi(x, t)$$

denote the relational field introduced in previous chapters.

We define the **Relational Action**

$$S[\Phi] = \int_{\Omega} L(\Phi, \partial_{\mu}\Phi) d^4x, \quad (33.1)$$

where

- S is the action,
- L is the relational Lagrangian density,
- d^4x denotes integration over spacetime.

The action summarizes the complete dynamical information of the relational system.

Definition 33.1 (Relational Action)

The **Relational Action** is a scalar functional whose stationary points determine the physically admissible evolution of a relational field.

Unlike the Hamiltonian, which emphasizes temporal evolution, the action provides a global description over an entire spacetime history.

33.3 The Relational Lagrangian

The Lagrangian density is decomposed as

$$L = L_{\text{kin}} - V + L_{\text{int}} + L_{\text{env}}, \quad (33.2)$$

where

- L_{kin} describes intrinsic relational dynamics,
- V is the relational potential,
- L_{int} represents interactions,
- L_{env} models environmental influences.

This decomposition parallels that used in conventional quantum field theory while accommodating the open-system nature of relational dynamics.

33.4 Stationary Action Principle

The central postulate is

$$\delta S = 0. \quad (33.3)$$

This does **not** imply that the action is always minimized.

Rather, the physical trajectory renders the action **stationary**, meaning that infinitesimal variations produce no first-order change.

Accordingly, the resulting solution may correspond to a minimum, maximum, or saddle point.

Definition 33.2 (Stationary Relational Evolution)

A relational evolution is called **stationary** if it satisfies

$$\delta S = 0.$$

Such trajectories satisfy the Euler–Lagrange equations derived below.

33.5 Euler–Lagrange Equations

Applying the calculus of variations yields

$$\frac{\partial L}{\partial \Phi} - \partial_\mu \left(\frac{\partial L}{\partial (\partial_\mu \Phi)} \right) = 0. \quad (33.4)$$

These equations constitute the fundamental field equations of SQFT in Lagrangian form.

Interpretation

Equation (33.4) governs the evolution of relational fields in exactly the same way that the Euler–Lagrange equations govern

- classical mechanics,
- electromagnetism,
- quantum field theory.

Thus, the dynamical laws of SQFT emerge from a single variational principle rather than being independently postulated.

33.6 Symmetry of the Action

Suppose the action remains invariant under the continuous transformation

$$\Phi \rightarrow \Phi + \epsilon X(\Phi), \quad (33.5)$$

where

$$\epsilon$$

is infinitesimal.

Such invariance implies the existence of conserved quantities through Noether's theorem.

Symmetry therefore serves as the fundamental origin of conservation laws within SQFT.

33.7 Relational Potential

The potential

$$V(\Phi)$$

determines stable relational configurations.

A common example is

$$V(\Phi) = \alpha\Phi^2 + \beta\Phi^4. \quad (33.6)$$

Depending upon the parameters,

the potential may possess

- a single equilibrium,
- multiple equilibria,
- spontaneous symmetry breaking,
- metastable states.

These structures correspond to alternative institutional configurations within the relational field.

33.8 Action and Hamiltonian

The canonical momentum is defined as

$$\Pi = \frac{\partial L}{\partial(\partial_t\Phi)}. \quad (33.7)$$

The Hamiltonian density follows from the Legendre transformation,

$$H = \Pi\partial_t\Phi - L. \quad (33.8)$$

Consequently,

the Hamiltonian formulation developed in earlier chapters is mathematically equivalent to the variational formulation whenever the required regularity conditions are satisfied.

33.9 Dissipative Systems

Open relational systems exchange information and resources with their environment.

Pure action principles generally describe conservative dynamics.

To incorporate dissipation,

one introduces either

- generalized variational principles,
- influence functionals,
- effective actions,

or

- non-Hermitian contributions.

The resulting equations reproduce the Lindblad dynamics discussed in Chapter 9 while retaining an underlying variational interpretation where applicable.

33.10 Effective Action

Microscopic relational variables are often inaccessible.

Integrating over hidden degrees of freedom produces the **effective action**

$$\Gamma[\Phi]. \quad (33.9)$$

The effective action governs macroscopic relational behavior after unresolved microscopic fluctuations have been incorporated.

This concept naturally complements the renormalization methods developed in Chapter 23.

33.11 Principle of Structural Economy

The stationary-action principle admits a broader interpretation within SQFT.

Rather than suggesting that institutions literally minimize effort, it states that observed relational evolution follows trajectories that satisfy the governing variational equations implied by the chosen action functional.

In many applications, these trajectories may correspond to efficient resolutions of competing structural constraints, but such interpretations depend upon the specific model and should not be regarded as universal laws of social behavior.

Theorem 33.1 (Variational Equivalence)

Suppose

1. the relational action is differentiable;
2. admissible variations vanish on the boundary;
3. the Legendre transformation is regular.

Then every stationary point of the action satisfies the Euler–Lagrange equations, and these equations are locally equivalent to the Hamiltonian equations introduced in previous chapters.

Proof

Taking the first variation of the action and integrating by parts while imposing vanishing boundary variations yields the Euler–Lagrange equations. Under the regularity condition ensuring an invertible Legendre transformation, the canonical momenta are uniquely defined, and the Lagrangian and Hamiltonian formulations describe the same local dynamics. Therefore, stationary-action trajectories are equivalent to solutions of the Hamiltonian equations.

Corollary 33.1

The Hamiltonian formalism, operator formulation, and variational formulation of SQFT constitute mathematically equivalent descriptions of the same underlying relational dynamics whenever the assumptions of Theorem 33.1 are satisfied.

33.12 Discussion

The introduction of a variational formulation significantly strengthens the theoretical foundations of Social Quantum Field Theory. By deriving the equations of motion from a single action functional, SQFT joins the family of modern field theories whose dynamics arise from stationary-action principles rather than independent postulates.

This formulation also establishes a natural bridge to gauge theory, differential geometry, and advanced quantum field theory, providing a common mathematical language for subsequent developments.

In the next chapter, we introduce **Noether's Theorem and Conservation Laws in Relational Fields**, demonstrating how continuous symmetries of the relational action generate conserved quantities that govern the persistence, stability, and transport of structure within complex relational systems.

Chapter 34

Noether's Theorem and Conservation Laws in Relational Fields

34.1 Introduction

The previous chapter established that the dynamics of Social Quantum Field Theory (SQFT) may be derived from a stationary-action principle. Once a Lagrangian formulation exists, an immediate and profound consequence follows.

One of the most fundamental results in modern theoretical physics is **Noether's Theorem**, proved by Emmy Noether in 1918. The theorem demonstrates that every continuous symmetry of the action corresponds to a conserved quantity.

This principle underlies virtually every conservation law in physics:

- time translation symmetry $\rightarrow$ energy conservation,
- spatial translation symmetry $\rightarrow$ momentum conservation,
- rotational symmetry $\rightarrow$ angular momentum conservation,
- gauge symmetry $\rightarrow$ conserved charges.

Within SQFT, Noether's Theorem provides the mathematical connection between **structural symmetries** and **persistent relational quantities**, revealing why certain institutional structures, information flows, and organizational patterns remain stable despite continual local fluctuations.

34.2 Continuous Symmetry

Consider the relational field

$$\Phi(x).$$

Suppose the action remains invariant under the infinitesimal transformation

$$\Phi \rightarrow \Phi + \varepsilon X(\Phi), \quad (34.1)$$

where

- ε is infinitesimal,
- $X(\Phi)$ is the generator of the transformation.

If

$$\delta S = 0,$$

the action possesses a continuous symmetry.

Definition 34.1 (Continuous Relational Symmetry)

A **continuous relational symmetry** is a smooth transformation of the relational field that leaves the action invariant.

Such symmetries characterize structural invariances rather than static configurations.

34.3 Noether Current

Associated with every continuous symmetry is a conserved current,

$$J^\mu = \frac{\partial L}{\partial(\partial_\mu \Phi)} \delta\Phi. \quad (34.2)$$

The current satisfies the continuity equation

$$\partial_\mu J^\mu = 0. \quad (34.3)$$

Equation (34.3) expresses local conservation.

Interpretation

The Noether current measures the transport of conserved relational quantities across the relational field.

Depending on the symmetry considered, the current may represent

- information flow,
 - organizational influence,
 - institutional continuity,
 - resource transmission,
 - structural coherence.
-

34.4 Conserved Charges

Integrating the temporal component of the current gives the conserved quantity

$$Q = \int J^0 d^3x. \quad (34.4)$$

Differentiating with respect to time yields

$$\frac{dQ}{dt} = 0, \quad (34.5)$$

provided no external sources or sinks break the underlying symmetry.

Definition 34.2 (Relational Conserved Quantity)

A **Relational Conserved Quantity** is a scalar derived from a continuous symmetry of the relational action that remains invariant throughout the system's evolution under the assumptions of the governing model.

34.5 Temporal Symmetry

If the relational action is invariant under

$$t \rightarrow t + \varepsilon, \quad (34.6)$$

the associated conserved quantity is

$$E.$$

Within SQFT,

this quantity represents the conservation of the system's effective relational energy under time-independent dynamics.

Examples include

- stable institutional commitments,
 - persistent organizational resources,
 - invariant structural capacities.
-

34.6 Spatial Symmetry

If the action is invariant under

$$x \rightarrow x + \varepsilon, \quad (34.7)$$

the conserved quantity is momentum.

Within relational systems,

this corresponds to the preservation of directional information propagation through homogeneous relational environments.

Examples include

- stable communication channels,
 - consistent supply-chain transmission,
 - uninterrupted knowledge diffusion.
-

34.7 Internal Symmetry

Many relational transformations leave observable behavior unchanged while altering internal representations.

Suppose

$$\Phi \rightarrow e^{i\alpha}\Phi. \quad (34.8)$$

Such internal symmetries generate conserved relational charges.

These quantities may characterize

- institutional identity,
- organizational membership,
- network affiliation,
- persistent classification structures.

Importantly, the specific interpretation depends upon the chosen model rather than following universally from the mathematics.

34.8 Broken Symmetry

Real relational systems frequently violate exact symmetry.

Suppose

$$\delta S \neq 0. \quad (34.9)$$

Then

$$\partial_\mu J^\mu \neq 0, \quad (34.10)$$

and the corresponding quantity is no longer strictly conserved.

Instead,

its rate of change reflects the magnitude of symmetry breaking.

Examples include

- institutional reform,
- regulatory intervention,
- organizational restructuring,
- technological disruption.

34.9 Approximate Conservation

Perfect symmetry is uncommon in empirical systems.

Nevertheless,

small symmetry-breaking terms produce approximately conserved quantities.

Suppose

$$\partial_\mu J^\mu = \epsilon, \quad (34.11)$$

where

$$|\epsilon| \ll 1.$$

Then

$$Q(t)$$

changes only gradually,

providing useful long-term structural invariants.

This approximation explains why many organizations exhibit remarkable stability despite continual local adaptation.

34.10 Conservation in Open Systems

Earlier chapters introduced Lindblad dynamics for open relational systems.

In such systems,

environmental exchange modifies the continuity equation,

$$\partial_\mu J^\mu = \Sigma, \quad (34.12)$$

where

$$\Sigma$$

represents external sources or sinks.

Consequently,

conservation laws become balance equations rather than exact invariants.

Interpretation

Examples include

- migration into or out of organizations,
- external investment,
- information injection,
- regulatory influence,
- environmental disturbances.

The conserved quantity evolves according to both internal dynamics and external exchange.

34.11 Hierarchies of Symmetry

Complex relational systems often possess multiple symmetry levels.

For example,

- microscopic interaction symmetry,
- organizational symmetry,
- institutional symmetry,
- societal symmetry.

Renormalization, discussed in Chapter 23, explains how symmetries may emerge, disappear, or transform across observational scales.

Thus,

Noether's Theorem naturally complements the multiscale architecture of SQFT.

Theorem 34.1 (Relational Noether Theorem)

Suppose

1. the relational action is differentiable;
2. the action is invariant under a continuous one-parameter transformation;
3. the relational field satisfies the Euler–Lagrange equations.

Then there exists a conserved current

$$J^\mu$$

satisfying

$$\partial_\mu J^\mu = 0.$$

Proof

The proof follows the standard variational derivation of Noether's Theorem. Under an infinitesimal continuous symmetry, the first-order variation of the action vanishes. Applying the Euler–Lagrange equations and integrating by parts isolates a divergence term, which defines the Noether current. Since the action is invariant, this divergence must vanish, yielding the continuity equation

$$\partial_\mu J^\mu = 0.$$

Thus every continuous symmetry generates a conserved current.

Corollary 34.1

Whenever a continuous symmetry of the relational action is only weakly broken, the associated conserved quantity evolves slowly, providing a natural explanation for long-lived institutional structures and persistent organizational patterns within the assumptions of the SQFT model.

34.12 Discussion

Noether's Theorem elevates symmetry from a descriptive property to a generative principle within Social Quantum Field Theory. Rather than postulating conservation laws independently, SQFT derives them directly from the invariance of the relational action. This unifies the treatment of information flow, organizational persistence, and structural stability within a single mathematical framework.

Combined with the variational formulation developed in the previous chapter, Noether's Theorem completes the Lagrangian foundations of SQFT and provides the conceptual bridge to modern gauge theories, where interactions themselves arise from local symmetry principles.

The next chapter introduces **Gauge Symmetry and Gauge Fields in Relational Systems**, extending SQFT by showing how local relational symmetries naturally give rise to interaction fields, relational connections, and covariant dynamics analogous to those found in contemporary quantum field theory.

Chapter 35

Gauge Symmetry and Gauge Fields in Relational Systems

35.1 Introduction

The previous chapter demonstrated that continuous symmetries give rise to conserved quantities through Noether's Theorem. A deeper insight from modern physics is that **interactions themselves** can emerge from symmetry.

In quantum field theory, requiring a theory to remain invariant under **local** rather than merely **global** symmetry transformations necessitates the introduction of gauge fields. Electromagnetism, the weak interaction, and the strong interaction all arise from this principle.

Social Quantum Field Theory (SQFT) adopts an analogous mathematical framework. Local relational symmetries imply the existence of **Relational Gauge Fields**, which mediate interactions among agents, institutions, and organizations while preserving structural consistency under local transformations.

Importantly, the term *gauge field* in SQFT is a mathematical analogy. It denotes a field that preserves local structural consistency within the relational model and should not be interpreted as implying the existence of a physical force analogous to electromagnetism.

35.2 Global and Local Symmetry

Consider the relational field

$$\Phi(x).$$

A global transformation is

$$\Phi(x) \rightarrow e^{i\alpha}\Phi(x), \quad (35.1)$$

where

$$\alpha$$

is constant throughout the relational domain.

The transformation affects every point identically.

A local transformation allows

$$\alpha = \alpha(x),$$

so that

$$\Phi(x) \rightarrow e^{i\alpha(x)}\Phi(x). \quad (35.2)$$

Different regions of the relational system may therefore undergo distinct local transformations.

Definition 35.1 (Local Relational Symmetry)

A **local relational symmetry** is a transformation whose parameters vary continuously across the relational manifold while leaving observable predictions invariant under the adopted SQFT model.

35.3 Failure of Ordinary Derivatives

Ordinary derivatives transform as

$$\partial_\mu \Phi,$$

which generally fail to preserve local symmetry because

$$\partial_\mu e^{i\alpha(x)} \neq e^{i\alpha(x)} \partial_\mu.$$

Consequently,

the ordinary derivative is not locally invariant.

A new mathematical object must therefore be introduced.

35.4 Covariant Derivative

The required object is the **covariant derivative**

$$D_\mu = \partial_\mu + igA_\mu, \quad (35.3)$$

where

- A_μ is the relational gauge field,
- g is the coupling constant.

The covariant derivative satisfies

$$D_\mu \Phi \rightarrow e^{i\alpha(x)} D_\mu \Phi, \quad (35.4)$$

preserving local symmetry.

Definition 35.2 (Relational Gauge Field)

A **Relational Gauge Field**

$$A_\mu$$

is a mathematical connection that compensates for local relational transformations, ensuring that the governing equations remain invariant under the specified symmetry group.

35.5 Geometric Interpretation

Gauge fields define how relational states are compared across different regions.

Rather than representing physical forces,

they specify the **connection** between neighboring relational configurations.

The relational gauge field therefore describes

- institutional coordination,
- communication protocols,
- regulatory consistency,
- organizational compatibility,
- shared operational standards.

These interpretations depend on the application and serve as mathematical representations of relational connections rather than literal physical fields.

35.6 Gauge Connection

The gauge field defines parallel transport along the relational manifold.

If

$$\Gamma$$

denotes a path,

the transported relational state becomes

$$\Phi_f = P \exp \left(i g \int_{\Gamma} A_{\mu} dx^{\mu} \right) \Phi_i, \quad (35.5)$$

where

$$P$$

denotes path ordering.

Parallel transport preserves local consistency while accounting for accumulated relational transformations.

35.7 Gauge Curvature

The field strength is defined by

$$F_{\mu\nu} = \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu} + i g [A_{\mu}, A_{\nu}]. \quad (35.6)$$

For Abelian symmetries,

the commutator vanishes,

leaving

$$F_{\mu\nu} = \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu}. \quad (35.7)$$

The curvature measures the extent to which relational transport depends upon the chosen path.

Interpretation

Within SQFT,

large gauge curvature may indicate

- inconsistent regulations,
- conflicting communication standards,
- incompatible organizational procedures,
- institutional friction.

Small curvature corresponds to relatively coherent relational coordination.

35.8 Gauge-Invariant Lagrangian

The gauge-field contribution to the action is

$$L_G = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (35.8)$$

The complete relational Lagrangian becomes

$$L = (D_\mu\Phi)^\dagger(D^\mu\Phi) - V(\Phi) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (35.9)$$

Gauge invariance determines the interaction structure without introducing arbitrary coupling terms.

35.9 Abelian and Non-Abelian Structures

The simplest symmetry group is

$$U(1),$$

whose generators commute.

More complex relational systems require non-Abelian groups,

such as

$$SU(N),$$

whose generators satisfy

$$[T_a, T_b] = if_{abc}T_c. \quad (35.10)$$

Non-Abelian gauge fields possess self-interactions,

making their dynamics substantially richer than those of Abelian fields.

Interpretation

Possible modeling examples include

Gauge Structure	Illustrative Relational Interpretation
$U(1)$	Uniform communication standard

Gauge Structure	Illustrative Relational Interpretation
$SU(2)$	Two-level institutional coordination
$SU(3)$	Multi-department interaction network
Higher Lie groups	Large-scale heterogeneous organizational systems

These examples are illustrative mathematical analogies rather than unique correspondences.

35.10 Gauge Fixing

Gauge symmetry introduces redundant mathematical descriptions.

A gauge condition,

such as

$$\partial_\mu A^\mu = 0, \tag{35.11}$$

removes this redundancy without changing observable predictions.

Gauge fixing therefore simplifies both analytical calculations and numerical simulations.

35.11 Emergent Interactions

A remarkable consequence of gauge theory is that interactions need not be postulated independently.

Instead,

they emerge naturally from the requirement of local symmetry.

Within SQFT,

this implies that many relational interactions may be represented as consequences of maintaining local structural consistency across the relational manifold.

This perspective shifts the emphasis from viewing interactions as isolated mechanisms to understanding them as manifestations of deeper symmetry principles within the theoretical framework.

Theorem 35.1 (Gauge Covariance of Relational Dynamics)

Suppose

- the relational field transforms under a continuous local symmetry group;
- ordinary derivatives are replaced by covariant derivatives;
- the gauge field transforms according to the corresponding connection law.

Then the relational Lagrangian (35.9) remains invariant under local gauge transformations.

Proof

Under a local transformation, the ordinary derivative acquires additional terms proportional to the gradient of the transformation parameter. The transformation law of the gauge field exactly cancels these additional contributions, ensuring that the covariant derivative transforms homogeneously. Since both the kinetic term and the gauge-field term are constructed from gauge-covariant quantities, the full Lagrangian remains invariant under local gauge transformations.

Corollary 35.1

Within the SQFT formalism, interaction fields may be introduced systematically from local symmetry requirements, providing a mathematically unified description of relational coupling across multiple organizational scales.

35.12 Discussion

Gauge symmetry represents one of the deepest organizing principles in modern theoretical physics, and its adaptation within SQFT significantly enriches the theory's mathematical structure. By replacing ordinary derivatives with covariant derivatives and introducing relational gauge fields, SQFT obtains a geometrically consistent description of local interactions that extends naturally across heterogeneous relational systems.

The gauge formulation also establishes direct connections with differential geometry, fiber bundles, and modern field theory, preparing the framework for more advanced geometric developments.

The next chapter introduces **Fiber Bundles and Differential Geometry of Relational Fields**, providing the geometric language in which gauge fields, covariant derivatives, and relational manifolds are unified into a single mathematical structure.

Chapter 36

Fiber Bundles and Differential Geometry of Relational Fields

36.1 Introduction

The previous chapter introduced gauge symmetry and demonstrated that local relational invariance naturally gives rise to gauge fields and covariant derivatives. Although these concepts may first appear algebraic, their deepest interpretation is fundamentally geometric.

Modern gauge theories are formulated not simply on spacetime itself, but on **fiber bundles**, mathematical structures that distinguish between the base space of observable events and the internal spaces attached to every point.

This chapter extends Social Quantum Field Theory (SQFT) into the language of differential geometry by interpreting relational systems as sections of fiber bundles endowed with connections and curvature. This geometric formulation unifies relational fields, gauge interactions, and information geometry within a common mathematical framework.

36.2 Relational Manifold

Let

M
 M

denote the **Relational Manifold**.

Unlike ordinary Euclidean space,
points of

M
represent relational configurations rather than physical locations.

Coordinates may include

- institutional variables,
- organizational states,
- information distributions,
- economic indicators,
- technological configurations.

The manifold therefore represents the state space of relational evolution.

Definition 36.1 (Relational Manifold)

A **Relational Manifold** is a smooth differentiable manifold whose points correspond to admissible relational configurations of the modeled system.

Its dimensionality depends upon the number of independent structural variables rather than physical spatial dimensions.

36.3 Tangent Spaces

Associated with every point

$$p \in M$$

is a tangent space

$$T_pM.$$

Vectors in

$$T_pM$$

describe infinitesimal changes of the relational configuration.

Examples include

- incremental policy adjustments,
- marginal changes in communication flow,
- small organizational perturbations,
- infinitesimal economic fluctuations.

36.4 Fiber Bundles


A fiber bundle consists of four components,

$$(E, M, \pi, F), \qquad (36.1)$$

where

- E is the total space,
- M is the relational manifold,
- $\pi: E \rightarrow M$ is the projection,
- F is the fiber attached to every point.

The structure is illustrated schematically.



Each point on the relational manifold possesses its own internal relational degrees of freedom.

Definition 36.2 (Relational Fiber)

A **Relational Fiber**

$$F_p$$

contains the internal relational variables associated with the configuration

$$p \in M.$$

These internal variables may represent latent organizational characteristics, hidden interaction modes, or internal state variables depending on the application.

36.5 Sections

A relational field corresponds to a section

$$s: M \rightarrow E, \quad (36.2)$$

satisfying

$$\pi \circ s = \text{id}_M. \quad (36.3)$$

Thus,

every relational configuration possesses an associated internal state.

Interpretation

A section assigns,

to every observable relational situation,

an internal structural description.

Examples include

- latent organizational culture,
 - institutional trust,
 - hidden communication modes,
 - internal technological capability.
-

36.6 Connections

To compare fibers located at different points,
one requires a connection.

The connection is represented locally by

$$A_\mu,$$

introduced previously as the relational gauge field.

The connection specifies how internal relational states are transported across the manifold.

Definition 36.3 (Relational Connection)

A **Relational Connection** defines the rule for transporting internal relational information between neighboring configurations while preserving local gauge consistency.

Connections therefore describe structural continuity rather than physical motion.

36.7 Parallel Transport

Suppose a relational state

$$\Phi$$

moves along a curve

$$\gamma.$$

Parallel transport satisfies

$$D_\gamma \Phi = 0, \quad (36.4)$$

where

$$D$$

denotes the covariant derivative.

Parallel transport provides the mathematical description of preserving internal consistency during relational evolution.

Examples

Parallel transport may model

- policy implementation across institutions,
- knowledge transfer between organizations,
- standardization across multinational enterprises,
- technological diffusion through industrial ecosystems.

36.8 Curvature

Transport around a closed loop generally fails to return a relational state unchanged.

The curvature tensor is

$$R(X, Y) = D_X D_Y - D_Y D_X - D_{[X, Y]}. \quad (36.5)$$

Curvature measures intrinsic geometric incompatibility.

Within gauge theory,

this coincides with the field strength tensor.

Interpretation

Large relational curvature may correspond to

- conflicting institutional rules,
- incompatible governance structures,
- fragmented communication pathways,
- organizational inconsistency.

Small curvature indicates relatively coherent relational geometry.

36.9 Geodesics

The shortest (or more generally, stationary) paths on the manifold satisfy

$$\frac{D\dot{\gamma}}{dt} = 0. \quad (36.6)$$

These paths are called geodesics.

Within SQFT,

geodesics represent dynamically preferred trajectories under the chosen relational metric.

They do not necessarily correspond to optimal human decisions, but rather to stationary trajectories determined by the geometry of the model.

36.10 Metric Geometry

Let

$$g_{ij}$$

denote the metric tensor on

$$M.$$

Distances satisfy

$$ds^2 = g_{ij}dx^i dx^j. \quad (36.7)$$

The metric determines

- relational distance,
- structural similarity,
- local geometry,
- information propagation.

Depending on the application, the metric may be derived from Fisher information (Chapter 25), empirical covariance structures, or other geometrically meaningful constructions.

36.11 Holonomy

Transport around a closed loop produces a transformation

$$U(\gamma).$$

The collection of all such transformations forms the **holonomy group**.

Holonomy characterizes global geometric properties that cannot be inferred from purely local observations.

Within relational systems,

holonomy may represent accumulated structural effects resulting from cyclic organizational processes or repeated institutional interactions.

Theorem 36.1 (Equivalence Between Gauge Fields and Connections)

Suppose

1. the relational field is represented as a section of a principal fiber bundle;
2. the bundle admits a smooth connection;
3. local trivializations exist over the manifold.

Then the local gauge field

$$A_\mu$$

is precisely the local representation of the bundle connection, and covariant differentiation corresponds to differentiation with respect to that connection.

Proof

A connection on a principal fiber bundle determines a connection one-form. Under local trivialization, this one-form is represented by the gauge potential

$$A_\mu.$$

The induced covariant derivative on associated bundles is obtained by combining the ordinary derivative with the connection term, reproducing the gauge-covariant derivative introduced in Chapter 35. Conversely, local gauge transformations correspond exactly to changes of local trivialization, demonstrating that gauge fields are local coordinate representations of a global geometric connection.

Corollary 36.1

Gauge interactions in SQFT admit a natural geometric interpretation as connection structures on relational fiber bundles, allowing local relational dynamics to be analyzed using the tools of differential geometry.

36.12 Discussion

The fiber-bundle formulation elevates Social Quantum Field Theory from an operator-based formalism to a fully geometric field theory. Relational fields become sections of fiber bundles, gauge fields become geometric connections, and interactions arise from the curvature of those connections. This unification clarifies the relationship between gauge symmetry, information geometry, topology, and variational dynamics developed throughout the preceding chapters.

The geometric language also provides the foundation for advanced mathematical developments, including characteristic classes, index theorems, and geometric quantization, while offering a coherent framework for studying multiscale relational systems with both local and global structure.

The next chapter introduces **Geometric Quantization of Relational Fields**, demonstrating how classical relational geometry can be systematically transformed into a quantum relational theory through symplectic geometry, polarization, and operator quantization.

Chapter 37

Geometric Quantization of Relational Fields

37.1 Introduction

The preceding chapter established that relational dynamics admit a geometric formulation through manifolds, fiber bundles, gauge connections, and curvature. This geometric framework provides a classical description of relational evolution.

The next fundamental question is:

How can a classical relational geometry be systematically transformed into a quantum relational theory?

In modern mathematical physics, this question is addressed through **Geometric Quantization**, a procedure that constructs a Hilbert space and quantum operators from an underlying symplectic manifold. Rather than replacing geometry with quantum mechanics, geometric quantization demonstrates that quantum theory emerges naturally from geometric structure.

Within Social Quantum Field Theory (SQFT), geometric quantization provides a mathematically rigorous bridge between classical relational geometry and the operator formalism developed in earlier chapters.

37.2 Classical Phase Space

Let

$$P$$

denote the relational phase space.

Each point is represented by

$$(q^i, p_i), \quad (37.1)$$

where

- q^i denotes generalized relational coordinates,
- p_i denotes their conjugate relational momenta.

Unlike physical mechanics, these variables need not correspond to spatial position or mechanical momentum. Instead, they parameterize the dynamical state of the modeled relational system.

Definition 37.1 (Relational Phase Space)

A **Relational Phase Space** is a smooth manifold whose points specify complete instantaneous relational states together with their conjugate dynamical variables.

37.3 Symplectic Structure

The essential geometric structure of phase space is the **symplectic two-form**

$$\omega = dq^i \wedge dp_i. \quad (37.2)$$

This differential form satisfies

$$d\omega = 0, \quad (37.3)$$

and is non-degenerate.

The symplectic form defines the canonical geometry of relational dynamics.

Interpretation

Within SQFT,

the symplectic structure determines

- admissible relational trajectories,
 - conservation structure,
 - canonical transformations,
 - Hamiltonian evolution.
-

37.4 Poisson Brackets

Given smooth functions

$$f(q, p), \quad g(q, p),$$

their Poisson bracket is

$$\{f, g\} = \sum_i \left(\frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i} \right). \quad (37.4)$$

Poisson brackets generate classical relational dynamics.

The evolution of any observable satisfies

$$\frac{df}{dt} = \{f, H\}. \quad (37.5)$$

37.5 Canonical Quantization

Quantization replaces classical observables by operators,

$$f \rightarrow \hat{f}, \quad (37.6)$$

and replaces Poisson brackets by commutators,

$$\{f, g\} \rightarrow \frac{1}{i\hbar} [\hat{f}, \hat{g}]. \quad (37.7)$$

Consequently,

canonical coordinates satisfy

$$[\hat{q}^i, \hat{p}_j] = i\hbar \delta_j^i. \quad (37.8)$$

These relations define the algebra of relational observables.

37.6 Prequantization

The first stage of geometric quantization constructs a complex line bundle

$$L \rightarrow P,$$

whose curvature satisfies

$$F = -\frac{i}{\hbar} \omega. \quad (37.9)$$

This condition guarantees compatibility between geometry and operator theory.

Definition 37.2 (Prequantum Line Bundle)

A **Prequantum Line Bundle** is a complex line bundle over the relational phase space whose curvature reproduces the symplectic form.

37.7 Polarization

Prequantization produces an excessively large Hilbert space.

A **polarization** selects a physically meaningful subspace.

For example,

choosing the coordinate polarization yields wavefunctions

$$\psi(q),$$

instead of functions depending on both

$$(q, p).$$

Polarization therefore eliminates redundant degrees of freedom while preserving the quantum structure.

37.8 Relational Hilbert Space

After polarization,

the quantum states form a Hilbert space

$$H = L^2(P). \quad (37.10)$$

Each vector

$$|\Psi\rangle$$

represents a relational quantum state.

Observables become self-adjoint operators acting on

$$H.$$

37.9 Quantization of Relational Observables

Classical functions

$$f(q, p)$$

become operators

$$\hat{f}.$$

Examples include

$$q \rightarrow \hat{q},$$

$$q \rightarrow \hat{q},$$

$$p \rightarrow -i\hbar \frac{\partial}{\partial q}, \quad (37.11)$$

and

$$H(q, p) \rightarrow \hat{H}.$$

The Schrödinger equation then governs relational evolution,

$$i\hbar \frac{\partial}{\partial t} |\Psi\rangle = \hat{H} |\Psi\rangle. \quad (37.12)$$

37.10 Coherent States

Between purely classical and fully quantum descriptions lie **coherent states**,

$$|\alpha\rangle.$$

These satisfy

$$\hat{a}|\alpha\rangle = \alpha|\alpha\rangle, \quad (37.13)$$

where

$$\hat{a}$$

is the annihilation operator.

Coherent states exhibit minimal uncertainty and often provide useful semiclassical approximations.

Within SQFT,

they may represent highly coordinated relational configurations whose fluctuations remain comparatively small.

37.11 Semiclassical Limit

As

$$\hbar \rightarrow 0,$$

the quantum equations converge toward the classical Hamiltonian dynamics,

$$\frac{df}{dt} = \{f, H\}. \quad (37.14)$$

This correspondence ensures mathematical consistency between the classical geometric formulation of Chapter 36 and the quantum operator formulation developed throughout earlier chapters.

The limit should be understood as a formal mathematical correspondence rather than a literal physical limit for social systems.

Theorem 37.1 (Correspondence Principle for Relational Quantization)

Suppose

1. the relational phase space possesses a symplectic structure;
2. geometric quantization is performed through a compatible prequantization and polarization;
3. the resulting quantum operators satisfy the canonical commutation relations.

Then, in the formal semiclassical limit,

$$\hbar \rightarrow 0,$$

the commutator algebra converges to the classical Poisson algebra,

$$\frac{1}{i\hbar}[\hat{f}, \hat{g}] \rightarrow \{f, g\}.$$

Proof

The proof follows the standard correspondence principle of geometric quantization. The operator algebra constructed from the symplectic manifold reproduces the Poisson algebra to leading order in $\hbar$. Consequently, expectation values of suitably regular observables satisfy the classical equations of motion in the formal semiclassical limit.

Corollary 37.1

The geometric and operator formulations of SQFT constitute mutually consistent descriptions connected through geometric quantization, allowing classical relational geometry and quantum relational dynamics to be analyzed within a unified mathematical framework.

37.12 Discussion

Geometric quantization provides the final mathematical bridge between classical differential geometry and the quantum operator formalism of Social Quantum Field Theory. Beginning with a symplectic manifold, one systematically constructs Hilbert spaces, operators, and quantum evolution while preserving the underlying geometric structure. This unifies the Hamiltonian, variational, gauge-theoretic, and geometric perspectives developed throughout the preceding chapters.

Beyond its formal elegance, this framework establishes a common language through which classical approximations, semiclassical methods, and fully quantum relational models can be studied as different regimes of the same mathematical theory.

The next chapter develops **Path Integrals and Functional Methods for Relational Fields**, introducing Feynman's sum-over-histories formulation and functional integration, thereby providing an alternative but equivalent formulation of SQFT based on probability amplitudes over entire relational trajectories rather than individual dynamical equations.

Chapter 38

Path Integrals and Functional Methods for Relational Fields

38.1 Introduction

The previous chapter established the operator formulation of Social Quantum Field Theory (SQFT) through geometric quantization. States evolve according to Schrödinger dynamics, and observables are represented by operators acting on a Hilbert space.

An alternative—but mathematically equivalent—formulation was introduced by Richard Feynman through the **path integral**.

Rather than describing evolution by a single trajectory or wavefunction, the path integral assigns a probability amplitude to **every admissible history** connecting an initial and a final configuration.

Within SQFT, this formulation provides a powerful framework for analyzing uncertainty, collective behavior, emergent phenomena, and statistical ensembles of relational evolution.

Importantly, the "sum over histories" should be interpreted here as a mathematical integration over possible trajectories within the model, rather than as a claim that real social systems physically realize all possible histories.

38.2 Relational Histories

Let

$$\Phi(t)$$

represent a relational field configuration.

Suppose the system evolves from

$$\Phi_i$$

to

$$\Phi_f.$$

Instead of assuming one deterministic evolution,

SQFT considers the collection

$$\Omega(\Phi_i, \Phi_f)$$

of all admissible relational trajectories satisfying the prescribed boundary conditions.

Definition 38.1 (Relational History)

A **Relational History** is a continuous trajectory in the configuration space of relational fields connecting specified initial and final configurations.

Each history contributes to the transition amplitude according to the action functional.

38.3 Transition Amplitude

The transition amplitude between two relational states is formally written as

$$K(\Phi_f, \Phi_i) = \int D\Phi e^{\frac{i}{\hbar} S[\Phi]}. \quad (38.1)$$

where

- $D\Phi$ denotes integration over admissible relational histories,
- $S[\Phi]$ is the relational action introduced in Chapter 33.

The integral is understood as a functional integral over field configurations rather than an ordinary finite-dimensional integral.

38.4 Functional Integration

Unlike ordinary integration,

functional integration sums over functions.

Symbolically,

$$\int D\Phi = \prod_x \int d\Phi(x). \quad (38.2)$$

Although formal,

this notation becomes mathematically precise through lattice regularization, discretization, or other rigorous constructions.

Interpretation

Functional integration accounts simultaneously for

- local fluctuations,
 - alternative relational trajectories,
 - uncertainty,
 - collective dynamics.
-

38.5 Classical Limit

When

$$\hbar$$

is formally small,

the rapidly oscillating phases cancel except near stationary points of the action.

Consequently,

$$\delta S = 0$$

dominates the functional integral.

The classical equations of Chapter 33 therefore emerge naturally as the stationary-phase approximation of the path integral.

38.6 Correlation Functions

Observable quantities are computed through expectation values.

For an operator

$$O[\Phi],$$

the expectation value is

$$\langle O \rangle = \frac{\int D\Phi O[\Phi] e^{\frac{i}{\hbar} S[\Phi]}}{\int D\Phi e^{\frac{i}{\hbar} S[\Phi]}}. \quad (38.3)$$

Correlation functions characterize statistical dependencies among different regions of the relational field.

Examples

Correlation functions may quantify

- coordination among organizational units,
- long-range communication dependence,
- institutional synchronization,
- propagation of structural influence.

Their interpretation depends on the observable selected.

38.7 Generating Functional

Introduce an external source

$$J(x).$$

The generating functional is

$$Z[J] = \int D\Phi \exp \left[\frac{i}{\hbar} \left(S[\Phi] + \int J\Phi d^4x \right) \right]. \quad (38.4)$$

Functional differentiation with respect to

$$J$$

generates all correlation functions.

For example,

$$\left. \frac{\delta Z[J]}{\delta J(x)} \right|_{J=0} = \langle \Phi(x) \rangle. \quad (38.5)$$

38.8 Effective Action

The connected generating functional is

$$W[J] = -i\hbar \ln Z[J]. \quad (38.6)$$

Its Legendre transform defines the effective action

$$\Gamma[\Phi] = W[J] - \int J\Phi d^4x. \quad (38.7)$$

The effective action incorporates the influence of fluctuations while preserving a variational structure.

It therefore complements the effective action introduced in Chapter 33 from a functional perspective.

38.9 Perturbation Expansion

Suppose

$$S = S_0 + S_{\text{int}}, \quad (38.8)$$

where

- S_0 is the free action,
- S_{int} represents interactions.

The exponential may be expanded as

$$e^{\frac{i}{\hbar}S} = e^{\frac{i}{\hbar}S_0} \left(1 + \frac{i}{\hbar}S_{\text{int}} + \cdots \right). \quad (38.9)$$

This perturbative expansion forms the basis for diagrammatic methods introduced in the following chapter.

38.10 Euclidean Formulation

Performing the Wick rotation

$$t \rightarrow -i\tau \quad (38.10)$$

transforms the oscillatory integral into

$$Z = \int D\Phi e^{-S_E[\Phi]/\hbar}, \quad (38.11)$$

where

$$S_E$$

is the Euclidean action.

The Euclidean formulation often improves numerical stability and is widely used in lattice field theory and Monte Carlo simulations.

Within SQFT, it provides a practical computational representation rather than a statement about physical time.

38.11 Statistical Interpretation

The Euclidean path integral closely resembles the partition function of statistical mechanics,

$$Z = \sum_i e^{-\beta E_i}. \quad (38.12)$$

This analogy establishes a mathematical bridge between

- quantum field theory,
- statistical mechanics,
- information theory,
- relational dynamics.

Consequently,

many computational methods developed in statistical physics become applicable to SQFT.

Theorem 38.1 (Stationary-Phase Approximation)

Suppose

1. the relational action is twice continuously differentiable;
2. the functional integral is dominated by neighborhoods of isolated stationary points;

3. higher-order fluctuations remain sufficiently small.

Then the leading contribution to the transition amplitude is determined by the stationary solutions satisfying

$$\delta S = 0.$$

Proof

Applying the functional version of the stationary-phase method, the action is expanded about each stationary configuration. The linear term vanishes by definition, while the quadratic term determines the leading Gaussian contribution. Higher-order terms contribute corrections of increasing order. Hence, the dominant contribution arises from stationary histories satisfying the Euler–Lagrange equations.

Corollary 38.1

The path-integral and variational formulations of SQFT are mathematically equivalent at the level of stationary trajectories, while the path-integral formalism additionally provides a systematic framework for incorporating fluctuations around those trajectories.

38.12 Discussion

The path-integral formulation completes the three principal mathematical descriptions of Social Quantum Field Theory:

- the **Hamiltonian formulation**, emphasizing operators and time evolution;
- the **Lagrangian formulation**, emphasizing variational principles and symmetry;
- the **Functional formulation**, emphasizing ensembles of admissible relational histories.

These formulations are mathematically equivalent under appropriate regularity conditions, yet each offers distinct conceptual and computational advantages. Functional methods are particularly valuable for perturbation theory, effective actions, statistical inference, and numerical simulation.

The next chapter introduces **Diagrammatic Expansions and Relational Feynman Diagrams**, where perturbative interactions are represented graphically, providing an intuitive and computationally efficient language for calculating interaction amplitudes, correlation functions, and higher-order corrections within Social Quantum Field Theory.

Chapter 39

Diagrammatic Expansions and Relational Feynman Diagrams

39.1 Introduction

The previous chapter introduced the functional formulation of Social Quantum Field Theory (SQFT), in which relational dynamics are represented through path integrals over all admissible histories. Although the path-integral formalism provides a compact mathematical description, direct evaluation of functional integrals is rarely possible except for the simplest systems.

Perturbation theory provides a practical alternative. Rather than solving the complete theory exactly, one expands the interaction terms around a solvable free theory. Each term in this expansion can be represented graphically by a **Feynman diagram**, transforming lengthy algebraic expressions into intuitive combinatorial objects.

Within SQFT, these diagrams should be interpreted as graphical representations of terms in a perturbative expansion. They do **not** depict literal physical trajectories or causal flows between social actors.

39.2 Free and Interaction Actions

Suppose the relational action is decomposed as

$$S[\Phi] = S_0[\Phi] + S_{\text{int}}[\Phi], \quad (39.1)$$

where

- S_0 describes the free relational field,
- S_{int} contains interaction terms.

The generating functional becomes

$$Z = \int D\Phi e^{\frac{i}{\hbar}(S_0 + S_{\text{int}})}. \quad (39.2)$$

Expanding the exponential yields

$$e^{\frac{i}{\hbar}S_{\text{int}}} = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{iS_{\text{int}}}{\hbar} \right)^n. \quad (39.3)$$

Each order of the expansion corresponds to a collection of interaction diagrams.

Definition 39.1 (Perturbative Order)

The **perturbative order** of a diagram equals the power of the interaction action appearing in the perturbation expansion.

Higher-order diagrams generally describe increasingly indirect interaction processes.

39.3 Propagators

The fundamental building block is the propagator,

$$G(x, y) = \langle 0 | T \{ \Phi(x) \Phi(y) \} | 0 \rangle, \quad (39.4)$$

where

$$T$$

denotes time ordering.

The propagator describes the mathematical correlation between two relational events within the perturbative framework.

It should be interpreted as a Green's function of the governing field equations rather than a literal transmission of information.

39.4 Interaction Vertices

Suppose the interaction Lagrangian contains

$$L_{\text{int}} = -\frac{\lambda}{4!} \Phi^4. \quad (39.5)$$

Each interaction contributes a four-leg vertex.

Graphically,



represents a local interaction event in the perturbative expansion.

The number of connecting lines depends on the structure of the interaction term.

39.5 Diagrammatic Rules

Each Feynman diagram corresponds to an algebraic contribution obtained through a fixed set of rules.

A typical perturbative calculation proceeds by

1. assigning propagators to internal lines,


2. assigning interaction factors to vertices,
3. integrating over internal variables,
4. multiplying symmetry factors,
5. summing all diagrams of the required order.

These rules provide a systematic algorithm for computing perturbative corrections.

39.6 Tree Diagrams

Diagrams without closed loops are called **tree diagrams**.

Example:



Tree diagrams describe the leading-order approximation of relational interactions.

Because they contain no internal loops,

their evaluation is comparatively straightforward.


Interpretation

Tree-level approximations are useful when interaction strengths remain sufficiently weak that higher-order corrections are negligible.

39.7 Loop Diagrams

Closed interaction paths generate **loop diagrams**.

Example:



Loops represent higher-order corrections arising from repeated or indirect interactions within the perturbative expansion.

Mathematically,

they modify

- propagators,
- coupling constants,
- effective interaction strengths.

39.8 Connected and Disconnected Diagrams

A diagram is **connected** if every component is joined through interaction lines.


Disconnected diagrams factorize into independent contributions.

Only connected diagrams contribute directly to connected correlation functions, which characterize genuine relational dependence beyond independent subsystems.

39.9 Vacuum Diagrams

Diagrams without external lines are called **vacuum diagrams**.

Example:



Although vacuum diagrams do not describe observable transitions directly, they contribute to the normalization of the generating functional and the effective action.

Within SQFT,

they characterize background structural fluctuations rather than observable relational events.

39.10 Renormalized Diagrams

Loop corrections often produce divergent integrals in continuum field theories.

Renormalization introduces counterterms,

$$L \rightarrow L + \delta L, \quad (39.6)$$

which absorb these divergences into redefined parameters.

The renormalization procedures developed in Chapter 23 therefore reappear naturally within the diagrammatic formulation.

39.11 Effective Interactions

Summing infinitely many perturbative diagrams produces effective quantities,

including

- effective propagators,
- effective coupling constants,
- effective masses,
- effective actions.

These renormalized objects describe large-scale relational behavior after incorporating higher-order interaction effects.

This illustrates how microscopic interaction structure can influence emergent macroscopic dynamics.

Theorem 39.1 (Diagrammatic Expansion of Correlation Functions)

Suppose

1. the interaction action admits a convergent or asymptotic perturbative expansion within the regime of interest;
2. the free theory is exactly solvable;
3. the generating functional is differentiable with respect to external sources.

Then every perturbative contribution to an n -point correlation function is represented by a finite collection of Feynman diagrams at each perturbative order.

Proof

Expanding the interaction exponential in powers of the interaction action generates products of field operators. Applying Wick's theorem rewrites these products as sums over pairwise contractions. Each contraction corresponds uniquely to a propagator, while interaction terms correspond to vertices. Consequently, every perturbative contribution admits a one-to-one graphical representation by a Feynman diagram.

Corollary 39.1

Diagrammatic methods provide a compact computational language for evaluating perturbative corrections to correlation functions, transition amplitudes, and effective interactions within the SQFT framework.

39.12 Discussion

Relational Feynman diagrams transform perturbative calculations from lengthy algebraic manipulations into an intuitive graphical language. Propagators encode pairwise correlations, interaction vertices encode local coupling terms, and loops describe higher-order corrections arising from indirect relational processes. Although these diagrams are mathematical bookkeeping devices rather than literal pictures of organizational behavior, they greatly simplify both analytical derivations and computational implementations.

Together with the path-integral formalism, diagrammatic methods provide powerful tools for studying interacting relational fields, effective theories, and multiscale dynamics.

The next chapter introduces **Non-Perturbative Methods and Topological Solitons in Relational Fields**, addressing phenomena that cannot be captured by perturbative expansions alone, including stable structural configurations, collective excitations, instanton-like transitions, and topologically protected relational states.

Chapter 40

Non-Perturbative Methods and Topological Solitons in Relational Fields

40.1 Introduction

The perturbative methods developed in the previous chapter provide systematic approximations when interaction strengths remain sufficiently small. Many complex systems, however, exhibit phenomena that lie beyond the reach of perturbation theory.

Examples include

- abrupt institutional transformations,
- persistent organizational structures,
- collective behavioral synchronization,
- long-lived network configurations,
- topologically protected relational states.

Such phenomena often arise from the global geometry of the relational field rather than from small local fluctuations.

This chapter introduces the **non-perturbative formulation** of Social Quantum Field Theory (SQFT), emphasizing stable field configurations, topological invariants, instanton-like transitions, and soliton solutions.

Throughout this chapter, terms such as *soliton* and *instanton* are employed as mathematical analogies for stable or transition-like solutions of the governing field equations, not as literal physical objects within social systems.

40.2 Perturbative and Non-Perturbative Regimes

Suppose the interaction strength is characterized by

$$g.$$

Perturbation theory expands observables as

$$F(g) = \sum_{n=0}^{\infty} a_n g^n. \quad (40.1)$$

This expansion is useful only when

$$|g| \ll 1.$$

When interactions become strong,

the perturbative series may converge slowly or merely provide an asymptotic approximation.

Alternative mathematical techniques are then required.

Definition 40.1 (Non-Perturbative Phenomenon)

A **non-perturbative phenomenon** is a property of the relational field that cannot be adequately described by any finite-order perturbative expansion.

40.3 Stable Field Configurations

Consider the relational action

$$S[\Phi].$$

Stationary configurations satisfy

$$\delta S = 0. \quad (40.2)$$

Among these solutions,

certain configurations remain stable under sufficiently small perturbations,

$$\Phi \rightarrow \Phi + \delta\Phi.$$

Such stability depends upon the structure of the action and the associated energy landscape.

Interpretation

Within SQFT,

stable field configurations may model

- persistent institutions,
- long-lived organizational arrangements,
- resilient technological ecosystems,
- enduring scientific communities.

40.4 Soliton Solutions

Suppose the nonlinear field equation admits localized solutions

$$\Phi_s(x, t).$$

These satisfy

$$\Phi_s(x, t) = \Phi_s(x - vt), \quad (40.3)$$

preserving their overall form during propagation.

Such solutions are called **solitons**.

Definition 40.2 (Relational Soliton)

A **Relational Soliton** is a stable localized solution of the nonlinear relational field equations whose structural identity persists throughout evolution within the assumptions of the model.

Interpretation

Illustrative analogies include

- highly stable institutional norms,
- persistent communication protocols,
- long-lived organizational cultures,
- robust technological standards.

These examples are intended as mathematical interpretations rather than direct physical equivalences.

40.5 Topological Stability

Some configurations remain stable not because they minimize local energy, but because they belong to distinct topological classes.

Suppose

$$\Phi: M \rightarrow N.$$

The homotopy class

$$[\Phi]$$

cannot be continuously deformed into another class.

The associated topological charge is

$$Q = \int q(x) d^3x, \quad (40.4)$$

where

$$q(x)$$

is the topological charge density.

Because

$$Q$$

is invariant under continuous deformations,

the corresponding configuration exhibits topological protection.

40.6 Domain Walls

Suppose the relational potential possesses two distinct stable minima,

$$\Phi_1, \quad \Phi_2.$$

The interface separating these phases is called a **domain wall**.

Graphically,



Domain walls represent boundaries between alternative stable relational structures.

Examples include

- institutional jurisdiction boundaries,
- organizational divisions,

- technological standard interfaces,
- administrative transitions.

40.7 Vortices

When the relational phase winds around a closed loop,

$$\oint \nabla \theta \cdot dl = 2\pi n, \quad (40.5)$$

the integer

$$n$$

defines a winding number.

Configurations possessing non-zero winding numbers cannot be removed by smooth deformations.

Within SQFT,

vortex-like structures represent mathematically persistent circulation patterns in relational variables.

40.8 Instanton-Like Transitions

Certain transitions occur through non-perturbative pathways connecting distinct stationary configurations.

Such solutions satisfy the Euclidean field equations,

$$\delta S_E = 0. \quad (40.6)$$

The corresponding transition amplitude behaves approximately as

$$P \propto e^{-S_E/\hbar}. \quad (40.7)$$

Within SQFT,

instanton-like solutions provide a mathematical representation of rare structural transitions between otherwise stable relational configurations.

40.9 Collective Excitations

Stable backgrounds may support small excitations,

$$\Phi = \Phi_s + \eta. \quad (40.8)$$

Linearization around

produces effective equations governing collective modes.

These excitations characterize local fluctuations without destroying the underlying stable configuration.

Examples include

- temporary organizational disturbances,
- short-lived communication anomalies,
- localized institutional adaptations.

40.10 Numerical Non-Perturbative Methods

Analytical solutions are rarely available for strongly interacting systems.

Numerical approaches include

- lattice discretization,
- tensor-network methods,
- variational optimization,
- Monte Carlo sampling,
- gradient-flow algorithms.

These methods enable the investigation of stable relational structures beyond perturbative approximations.

40.11 Emergent Structural Robustness

Topologically protected configurations exhibit remarkable robustness against local perturbations.

This robustness arises because changing the global topological class generally requires discontinuous transformations or sufficiently large perturbations rather than small local modifications.

Within SQFT,

such protection provides a mathematical explanation for why certain organizational structures remain stable despite continual local fluctuations.

Whether a particular empirical system exhibits topological protection must be established through application-specific modeling and validation.

Theorem 40.1 (Topological Stability of Relational Solitons)

Suppose

1. the relational field possesses multiple homotopy classes;
2. the topological charge

$$Q$$

is conserved under continuous deformations;

3. the field evolution remains continuous.

Then a configuration with non-trivial topological charge cannot be continuously transformed into the trivial configuration.

Proof

Continuous evolution preserves homotopy class. Since the topological charge uniquely labels distinct classes, any continuous deformation must preserve

$$Q.$$

Therefore, a configuration possessing non-zero topological charge cannot evolve continuously into a configuration with zero charge. Such transformations require discontinuities, singular configurations, or modifications of the underlying topology.

Corollary 40.1

Within the SQFT framework, topologically distinct relational configurations exhibit structural robustness against sufficiently small continuous perturbations, providing a mathematical mechanism for persistent large-scale organizational patterns.

40.12 Discussion

Non-perturbative methods extend Social Quantum Field Theory beyond the domain of weak interactions, enabling the study of stable structures, abrupt transitions, and topologically protected configurations that cannot be captured through perturbative expansions alone. Solitons, domain walls, vortices, and instanton-like solutions illustrate how global geometric and topological properties can govern long-term relational behavior.

Together with perturbation theory, renormalization, gauge symmetry, and differential geometry, these methods complete the principal analytical toolkit of SQFT, allowing both weakly and strongly interacting relational systems to be studied within a unified mathematical framework.

The next chapter develops **Quantum Anomalies and Symmetry Breaking Beyond the Classical Theory**, examining situations in which symmetries preserved at the classical level fail after quantization, thereby extending SQFT to include anomaly structures, effective symmetry breaking, and their implications for relational dynamics.



輸入留言

這個網誌中的熱門文章

Foundational Charter of the Social Quantum Field Theory Research Community


7月 19, 2026

The SQFT Charter Foundational Charter of the Social Quantum Field Theory Research Community Preamble The Social Quantum Field Theory (SQFT) Research Community is founded upon the conviction that mathematics advances through open inquiry ...

[閱讀完整內容](#)

社會量子場論（SQFT）超越布迪厄的場域理論，建構關係社會學的形式化 1~10

7月 15, 2026



預印本 社會量子場論與場論的形式化 周易先 獨立研究者 · 版本 2.0（數學版） · 2026年7月 摘要 本文提出社會量子場論（SQFT）作為用以表示複雜社會系統中關係動態的形式數學框架。SQFT並不主張社 ...

[閱讀完整內容](#)

量子之影：台灣QNF-3量子導航系統的崛起與其地緣政治影響




量子之影：台灣QNF-3量子導航系統的崛起與其地緣政治影響 摘要 在 2035年，全球局勢緊張之際，台灣的量子導航技術成為重塑國際權力格局的關鍵力量。本篇科幻論文式敘事，記錄了國家中山科學研 ...

[閱讀完整內容](#)

 技術提供：Blogger

主題圖片來源：[Michael Elkan](#)



IH

[瀏覽簡介](#)

封存



[檢舉濫用情形](#)