# Model-based optimal inference of spike times and calcium dynamics given noisy and intermittent calcium-fluorescence imaging


## Joshua T Vogelstein<sup>1</sup>, Baktash Babadi<sup>2</sup>, Liam Paninski<sup>2,3</sup>

<sup>1</sup>Department of Neuroscience, Johns Hopkins School of Medicine, <sup>2</sup>Department of Neuroscience, Columbia University, <sup>3</sup>Department of Statistics, Columbia University

joshuav@jhu.edu, bb2280@columbia.edu, liam@stat.columbia.edu

#### **Abstract**

The goal of this work is to infer spike trains, having only the fluorescence time-series available to us. To do so, we propose a parametric model of the experimental system. Then, we develop a number of algorithms, each iteratively estimating the parameters and then using those parameters to infer the hidden spike trains. Our algorithms make significant advances over previous approaches, in that they naturally facilitate: (i) intermittent observations, (ii) stimulus information, (iii) saturation effects, (iv) error bars, and (v) refractoriness.




### References


- 1) Smetters D, Majewska A, Yuste R. *Detecting action potentials in neu-ronal populations with calcium imaging*. Methods. 1999. Jun; 18(2):215-21
- 2) Kerr JN, Greenberg D, Helmchen F. *Imaging input and output of neo-cortical networks in vivo.* PNAS 2005 Sep 27;102(39):14063-8
- 3) Yaksi E, Friedrich RW. Reconstruction of firing rate changes across neuronal populations by temporally deconvolved Ca2+ imaging. Nat Methods. 2006 May;3(5):344-6.
- 3) Sato T, Gray NW, Mainen ZF, Svoboda K. *The Functional Microarchitecture of the Mouse Barrel Cortex*. PLOS Biol 2007 Jul 10; 5(7):e189.


  4) Vogelstein JT and Paninski L *Model-based optimal inference of*
- spike times and calcium dynamics given noisy and intermittent calcium-fluorescence imaging. Under review at Biophysical Journal.


## Acknowldgments


Support for JTV was provided by NIDCD DC00109. LP is supported by an NSF CA-REER award and by an Alfred P. Sloan Research Fellowship.

















9. comparison				
	other	barrier	PPR	SMC
parameters	variable	$\{a,b,c,d,k,A\}$	$\{a,b,c,d,k,A\}$	$+\{\alpha,\beta,\gamma,\delta,\omega,\sigma_h,\vec{a}_h\}$
speed	O(T)	O(T)	O(T)	$O(N^2T)$
intermittency	X	$\checkmark$	X	$\checkmark$
stimulus	X	$\checkmark$	<b>✓</b>	$\checkmark$
saturation	X	X	X	$\checkmark$
errorbars	X	X	X	$\checkmark$
refractoriness	X	X	X	$\checkmark$
noise assumptions	?	gaussian	gaussian	arbitrary

