

Hogfish

Species: *Lachnolaimus maximus*, Stock code: Hogfish - Florida Keys - East Florida


Region: Southeast Shelf

Marine Ecoregion: Florida Keys - East Florida


Reconstructed catch data used from years 1986 - 2012

For figure captions and method see <http://www.seaaroundus.org/cmsy-method>



Catch Hogfish - Florida Keys - East Floi



Stock size



Exploitation



extext

Results for management (based on BSM analysis)

$F_{msy} = 0.0873$, 95% CL = 0.0536 - 0.13 (if $B > 1/2 B_{msy}$ then $F_{msy} = 0.5 r$)

$F_{msy} = 0.0692$, 95% CL = 0.0425 - 0.103 (r and F_{msy} are linearly reduced if $B < 1/2 B_{msy}$)

$MSY = 26.6$, 95% CL = 19.7 - 37.1; $B_{msy} = 302$, 95% CL = 207 - 518 (1000 tonnes)







Biomass in last year = 123, 95% CL = 76.8 - 202 (1000 tonnes)

B/B_{msy} in last year = 0.397, 95% CL = 0.283 - 0.575

Fishing mortality in last year = 0.0269, 95% CL = 0.0154 - 0.0478

$F/F_{msy} = 0.392$, 95% CL = 0.202 - 0.821

Comment: NOAA Assessment model SS3. RF OK

A: Hogfish - Florida Keys - East Flor**B: Finding viable r-k****C: Analysis of viable r-k****D: Stock size****E: Exploitation rate****F: Equilibrium curve**

extext

Results of CMSY analysis conducted in JAGS

$r = 0.322$, 95% CL = 0.201 - 0.434; $k = 325$, 95% CL = 246 - 525 (1000 tonnes)

MSY = 26.2, 95% CL = 20.2 - 33.9 (1000 tonnes/year)

Relative biomass last year = 0.125 k , 95% CL = 0.0501 - 0.271

Exploitation $F/(r/2)$ in last year = 1.16

Results from Bayesian Schaefer model using catch and CPUE

$r = 0.175$, 95% CL = 0.107 - 0.261; $k = 604$, 95% CL = 415 - 1035

r - k log correlation = -0.772

MSY = 26.6, 95% CL = 19.7 - 37.1 (1000 tonnes/year)

Relative biomass in last year = 0.125 k , 95% CL = 0.0501 - 0.271

Exploitation $F/(r/2)$ in last year = 0.6

$q = 3.27$, 95% CL = 2.06 - 4.81

Prior range of q = 1.12 - 28.2

Relative abundance data type = CPUE

Prior initial relative biomass = 0.15 - 0.3 expert

Prior intermediate relative biomass = 0.0678 - 0.323 in year 2005 default

Prior final relative biomass = 0.02 - 0.25 expert

Prior range for r = 0.2 - 0.8 default, prior range for k = 70.2 - 443 (1000 tonnes) default

Source for relative biomass:

NOAA Fisheries. (2020). Assessment Summary Data. Retrieved from www.st.nmfs.noaa.gov/stocksmart. 12/13/2020