


**a** Three data sources and transport directions


Scenario-specific endpoints: 1-year mortality for NHANES -> MIMIC-IV; hospital mortality for MIMIC-IV ICU <=> eICU.

**b**

## Cohort sizes and endpoints by direction

| Direction               | Endpoint              | Source                                           | Target evaluation                             |
|-------------------------|-----------------------|--------------------------------------------------|-----------------------------------------------|
| NHANES<br>-> MIMIC-IV   | 1-year<br>mortality   | NHANES<br>N=6,564<br>Deaths=166 (2.5%)           | MIMIC-IV<br>N=33,201<br>Deaths=6,297 (19.0%)  |
| MIMIC-IV ICU<br>-> eICU | Hospital<br>mortality | MIMIC-IV ICU<br>N=25,906<br>Deaths=2,988 (11.5%) | eICU<br>N=13,971<br>Deaths=1,263 (9.0%)       |
| eICU<br>-> MIMIC-IV ICU | Hospital<br>mortality | eICU<br>N=55,486<br>Deaths=5,242 (9.4%)          | MIMIC-IV ICU<br>N=6,395<br>Deaths=712 (11.1%) |

Target evaluation sizes correspond to primary transport-performance analyses.

**c** NHANES-MIMIC separability increased with labs**d** ICU separability increased with labs and vitals