



# Echelon Forms and Row Reduction

Department of Computer Engineering  
Sharif University of Technology

Hamid R. Rabiee [rabiee@sharif.edu](mailto:rabiee@sharif.edu)

Maryam Ramezani [maryam.ramezani@sharif.edu](mailto:maryam.ramezani@sharif.edu)



# Table of contents

**01**

Row-Reduced Matrix

**02**

Echelon Form

**03**

Row-Reduced Echelon Form

**04**

Solutions of  
a Linear System

**05**

Geometric  
Interpretation



01

# Row-Reduced Matrix

# Row-Reduced Matrix

## Definition

□ A  $m \times n$  matrix  $R$  is called **row-reduced** if:

1. **Leading entries=1**: The first non-zero entry in each non-zero row of  $R$  is equal to 1.
2. All other entries in the columns corresponding to leading entries must be zero: Any column of  $R$  that contains a leading entry (the first non-zero entry) of a row must have all its other entries equal to zero.

# Row-Reduced Matrix

## Example

□ Are following matrices Row-Reduced Matrix?

a.  $n \times n$  identity matrix

b. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

c. 
$$\begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

d. 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -6 \end{bmatrix}$$

# Row-Reduced matrix for Every Matrix

## Theorem (1)

Every  $m \times n$  matrix is row-equivalent to a row-reduced matrix.

**Theorem 4.** Every  $m \times n$  matrix over the field  $F$  is row-equivalent to a row-reduced matrix.

*Proof.* Let  $A$  be an  $m \times n$  matrix over  $F$ . If every entry in the first row of  $A$  is 0, then condition (a) is satisfied in so far as row 1 is concerned. If row 1 has a non-zero entry, let  $k$  be the smallest positive integer  $j$  for which  $A_{1j} \neq 0$ . Multiply row 1 by  $A_{1k}^{-1}$ , and then condition (a) is satisfied with regard to row 1. Now for each  $i \geq 2$ , add  $(-A_{ik})$  times row 1 to row  $i$ . Now the leading non-zero entry of row 1 occurs in column  $k$ , that entry is 1, and every other entry in column  $k$  is 0.

Now consider the matrix which has resulted from above. If every entry in row 2 is 0, we do nothing to row 2. If some entry in row 2 is different from 0, we multiply row 2 by a scalar so that the leading non-zero entry is 1. In the event that row 1 had a leading non-zero entry in column  $k$ , this leading non-zero entry of row 2 cannot occur in column  $k$ ; say it occurs in column  $k' \neq k$ . By adding suitable multiples of row 2 to the various rows, we can arrange that all entries in column  $k'$  are 0, except the 1 in row 2. The important thing to notice is this: In carrying out these last operations, we will not change the entries of row 1 in columns 1,  $\dots$ ,  $k$ ; nor will we change any entry of column  $k$ . Of course, if row 1 was identically 0, the operations with row 2 will not affect row 1.

Working with one row at a time in the above manner, it is clear that in a finite number of steps we will arrive at a row-reduced matrix. ■

02

# Echelon Form



# Echelon form

## Definition

- A rectangular matrix is **in echelon form** (or **row echelon form**) if it has the following three properties:
1. All nonzero rows are above any rows of all zeros.
  2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
  3. All entries in a column below a leading entry are zeros.

$$\begin{bmatrix} 2 & -3 & 2 & 1 \\ 0 & 1 & -4 & 8 \\ 0 & 0 & 0 & \frac{5}{2} \end{bmatrix}$$

Echelon form



03

# Row-Reduced Echelon Form



# Row-Reduced Echelon form

## Definition

If a matrix in echelon form satisfies the following additional conditions, then it is in **reduced echelon form** (or **reduced row echelon form**):

1. The leading entry in each non-zero row is 1.
2. Each leading 1 is the only non-zero entry in its columns.
3. The leading 1 in the second row or beyond is to the right of the leading 1 in the row just above.
4. Any row containing only 0's is at the bottom.


$$\begin{array}{ccc|c} e_1 & e_2 & e_3 & \\ \hline 1 & 0 & 0 & 29 \\ 0 & 1 & 0 & 16 \\ 0 & 0 & 1 & 3 \end{array}$$

Reduced Echelon form

# Row-Reduced echelon matrix for Every Matrix

## Theorem (2)

Every  $m \times n$  matrix is row-equivalent to a row-reduced echelon matrix.



Proof is same as the Theorem (1).



# Reduced Echelon Form (RREF)

## Example

Are following matrices RREF?

*a.*  $0_{m \times n}$

*b.* 
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

*c.* 
$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

*d.* 
$$\begin{bmatrix} 0 & 1 & -3 & 0 & 0.5 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

# Existence and Uniqueness Questions

Two fundamental questions about a linear system:

1. Is the system consistent? That is, does at least one solution exist?
2. If a solution exists, is it the only one? That is, is the solution unique?



04

# Solutions of a Linear System



# Elementary Row Operations

## Example

□ Augmented matrix for a linear system:

$$\begin{bmatrix} 1 & 0 & -5 & 1 \\ 0 & 1 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{aligned} x - 5z &= 1 \\ y + z &= 4 \\ 0 &= 0 \end{aligned}$$

$$\begin{cases} x = 1 + 5z \\ y = 4 - z \\ z \text{ is free variable} \end{cases}$$

□  $x, y$ : basic variable       $z$ : free variable

□ This system is consistent, because the solution set can be described explicitly by solving the reduced system of equations for the basic variables in terms of the free variables.

# Existence and Uniqueness Questions


## Theorem

A linear system is **consistent** if and only if the rightmost column of the augmented matrix is not a pivot column – that is, if and only if an echelon form of the augmented matrix has no row of the form  $[0 \quad \cdots \quad 0 \quad b]$  with nonzero  $b$ .

- If a linear system is consistent, then the solution set contains either:
  - A unique solution, when there are no free variables
  - Infinitely many solutions, when there is at least one free variable



# Find all solutions of a linear system

- 
1. Write the augmented matrix of the system.
  2. Use the row reduction algorithm to obtain an equivalent augmented matrix in echelon form. Decide whether the system is consistent. If there is no solution, stop; otherwise, go to the next step.
  3. Continue row reduction to obtain the reduced echelon form.
  4. Write the system of equations corresponding to the matrix obtained in step 3.
  5. Rewrite each nonzero equation from step 4 so that its one basic variable is expressed in terms of any free variables appearing in the equation.



# Existence of Solutions

## Example

Let  $A = \begin{bmatrix} 1 & 3 & 4 \\ -4 & 2 & -6 \\ -3 & -2 & -7 \end{bmatrix}$  and  $b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ . Is the equation  $Ax = b$  consistent for all possible  $b_1, b_2, b_3$ ?

## Solution

Row reduce the augmented matrix for  $Ax = b$ :

$$\left[ \begin{array}{ccc|c} 1 & 3 & 4 & b_1 \\ -4 & 2 & -6 & b_2 \\ -3 & -2 & -7 & b_3 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & 4 & b_1 \\ 0 & 14 & 10 & b_2 + 4b_1 \\ 0 & 7 & 5 & b_3 + 3b_1 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 3 & 4 & b_1 \\ 0 & 14 & 10 & b_2 + 4b_1 \\ 0 & 0 & 0 & b_3 + 3b_1 - \frac{1}{2}(b_2 + 4b_1) \end{array} \right]$$

The third entry in column 4 equals  $b_3 + 3b_1 - \frac{1}{2}(b_2 + 4b_1)$ . The equation  $Ax = b$  is not consistent for every  $b$  because some choices of  $b$  can make  $b_1 - \frac{1}{2}b_2 + b_3$  nonzero.

# Existence of solutions

## Example

True or False?

Equation  $Ax = b$  is consistent, if its augmented matrix  $[A \ b]$  has one pivot column in each rows? (Having one leading entry in each rows)

# Homogeneous Linear Systems

## Definition

A system of linear equations is said to be **homogeneous** if it can be written in the form  $Ax = 0$ , where  $A$  is a matrix and  $0$  is the zero vector.

- **Trivial solution:**  $Ax = 0$  always has at least one solution, namely,  $x = 0$  (the zero vector)
- **Nontrivial solution:** The non-zero solution for  $Ax = 0$ .

## Fact

The homogenous equation  $Ax = 0$  has a nontrivial solution if and only if the equation has at least one free variable.

$$\left( \begin{array}{ccc|c} 1 & 3 & 4 & 0 \\ 2 & -1 & 2 & 0 \\ 1 & 0 & 1 & 0 \end{array} \right) \xrightarrow{RREF} \left( \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right)$$

# Existence Of Solutions

## Fact

The equation  $Ax = b$  has a solution if and only if  $b$  is a linear combination of the columns of  $A$ .



05


# Geometric Interpretation

# Line ( $\mathbb{R}^2$ )


The line  $\ell$  with equation  $2x + y = 0$

$\mathbf{n} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  and  $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$ , then the equation becomes  $\mathbf{n} \cdot \mathbf{x} = 0$ .


$\ell$  as  $\mathbf{x} = t\mathbf{d}$ .




The line  $2x + y = 0$



A normal vector  $\mathbf{n}$



A direction vector  $\mathbf{d}$



$\mathbf{n} \cdot (\mathbf{x} - \mathbf{p}) = 0$

**Definition** The normal form of the equation of a line  $\ell$  in  $\mathbb{R}^2$  is

$$\mathbf{n} \cdot (\mathbf{x} - \mathbf{p}) = 0 \quad \text{or} \quad \mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{p}$$

where  $\mathbf{p}$  is a specific point on  $\ell$  and  $\mathbf{n} \neq \mathbf{0}$  is a normal vector for  $\ell$ .

The general form of the equation of  $\ell$  is  $ax + by = c$ , where  $\mathbf{n} = \begin{bmatrix} a \\ b \end{bmatrix}$  is a normal vector for  $\ell$ .



## Lines in $\mathbb{R}^2$

Normal Form	General Form	Vector Form	Parametric Form
$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{p}$	$ax + by = c$	$\mathbf{x} = \mathbf{p} + t\mathbf{d}$	$\begin{cases} x = p_1 + td_1 \\ y = p_2 + td_2 \end{cases}$

# Plane ( $\mathbb{R}^3$ )

$$\mathbf{n} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \text{ and } \mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix},$$

$$ax + by + cz = d \text{ (where } d = \mathbf{n} \cdot \mathbf{p} \text{)}$$



## Lines and Planes in $\mathbb{R}^3$

	Normal Form	General Form	Vector Form	Parametric Form
Lines	$\begin{cases} \mathbf{n}_1 \cdot \mathbf{x} = \mathbf{n}_1 \cdot \mathbf{p}_1 \\ \mathbf{n}_2 \cdot \mathbf{x} = \mathbf{n}_2 \cdot \mathbf{p}_2 \end{cases}$	$\begin{cases} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2 \end{cases}$	$\mathbf{x} = \mathbf{p} + t\mathbf{d}$	$\begin{cases} x = p_1 + td_1 \\ y = p_2 + td_2 \\ z = p_3 + td_3 \end{cases}$
Planes	$\mathbf{n} \cdot \mathbf{x} = \mathbf{n} \cdot \mathbf{p}$	$ax + by + cz = d$	$\mathbf{x} = \mathbf{p} + s\mathbf{u} + t\mathbf{v}$	$\begin{cases} x = p_1 + su_1 + tv_1 \\ y = p_2 + su_2 + tv_2 \\ z = p_3 + su_3 + tv_3 \end{cases}$



# Nonhomogeneous Systems & General Solution

## Example

Describe all solutions of  $A\mathbf{x} = \mathbf{b}$ , where:  $A = \begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & -8 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 7 \\ -1 \\ -4 \end{bmatrix}$

$$\left[ \begin{array}{ccc|c} 3 & 5 & -4 & 7 \\ -3 & -2 & 4 & -1 \\ 6 & 1 & -8 & -4 \end{array} \right] \sim \left[ \begin{array}{ccc|c} 1 & 0 & -\frac{4}{3} & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

## Example

Describe all solutions of  $A\mathbf{x} = \mathbf{0}$ , where:  $A = \begin{bmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ .

## Example

Describe all solutions of  $A\mathbf{x} = \mathbf{b}$ , where:  $[A \ \mathbf{b}] = \begin{bmatrix} 1 & 2 & -1 & 1 & 3 \\ 1 & 1 & -1 & 1 & 1 \\ 1 & 3 & -1 & 1 & 5 \end{bmatrix}$ .

$$\left[ \begin{array}{ccccc|c} 1 & 2 & -1 & 1 & 3 & 3 \\ 0 & 1 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

# Nonhomogeneous Systems & General Solution

## Question

Can we change the order of columns in an augmented matrix???

$$\begin{cases} ax + by + cz = d \\ a'x + b'y + c'z = d' \\ a''x + b''y + c''z = d'' \end{cases}$$

Is equivalent to

$$\begin{cases} ax + cz + by = d \\ a'x + c'z + b'y = d' \\ a''x + c''z + b''y = d'' \end{cases}$$

# Resources

- ❑ Chapter 1: Kenneth Hoffman and Ray A. Kunze. Linear Algebra. PHI Learning, 2004.
- ❑ Chapter 1: David C. Lay, Steven R. Lay, and Judi J. McDonald. Linear Algebra and Its Applications. Pearson, 2016.
- ❑ Chapter 2: David Poole, Linear Algebra: A Modern Introduction. Cengage Learning, 2014.
- ❑ Chapter 1: Gilbert Strang. Introduction to Linear Algebra. Wellesley-Cambridge Press, 2016.

