

**PART - A**  
(Compulsory Question)

\*\*\*\*\*

- 1 Answer the following: (10 X 02 = 20 Marks)
- Construct the truth table  $\neg(7P \vee 7Q)$ .
  - Prove that  $(\exists x)(P(x) \wedge Q(x)) \Rightarrow (\exists x)P(x) \wedge (\exists x)Q(x)$ .
  - Let  $X = \{1, 2, \dots, 7\}$  and  $R = \{ \langle x, y \rangle \mid x - y \text{ is divisible by } 3 \}$  show that R is an equivalence relation. Draw the graph of R.
  - Let  $\lfloor \sqrt{x} \rfloor$  be the greatest integer  $\leq \sqrt{x}$ . Show that  $\lfloor \sqrt{x} \rfloor$  is primitive recursive.
  - Let  $\langle G, * \rangle$  be a finite cyclic group generated by an element  $a \in G$ . Prove that if G is of order n, i.e.  $|G| = n$  then  $a^n = e$  so that  $G = \{a, a^2, a^3, \dots, a^n = e\}$ . Furthermore n is the least +ve integer for which  $a^n = e$ .
  - Prove that the minimum weight of the nonzero code words in a group code is equal to its minimum distance.
  - Let  $\langle L, \leq \rangle$  be a lattice. For any  $a, b, c \in L$  then prove that  $b \leq c \Rightarrow a * b \leq a * c$ .
  - Prove the Boolean identity  $(a \wedge b) \vee (a \wedge b') = a$ .
  - Prove that a tree with n vertices has precisely n-1 edges.
  - A label identifier for a computer program consists of one letter followed by three digits. If repetitions are allowed, how many distinct label identifiers are possible.

**PART - B**

(Answer all five units, 5 X 10 = 50 Marks)

**UNIT - I**

- Show that the formula  $Q \vee (P \wedge \neg Q) \vee (\neg P \wedge \neg Q)$  is a tautology.
  - Obtain the principal conjunctive normal form of the formula given by  $(7P \rightarrow R) \wedge (Q \Rightarrow P)$ .
- OR**

  - Show that  $R \wedge (P \vee Q)$  is a valid conclusion from the premises  $P \vee Q$ ,  $Q \rightarrow R$ ,  $P \rightarrow M$  and  $\neg M$ .
  - Show that  $\neg P(a, b)$  follows logically from  $(x)(y) (P(x, y)) \rightarrow W(x, y)$  and  $\neg W(a, b)$ .

**UNIT - II**

- Let Z be the set of integers and let R be the relation called congruence modulo 3 defined by;  
 $R = \{ \langle x, y \rangle \mid x \in \mathbb{Z} \wedge y \in \mathbb{Z} \wedge (x - y) \text{ is divisible by } 3 \}$ . Determine the equivalence classes generated by the elements of  $\mathbb{Z}$ .
  - Let  $X = \{2, 3, 6, 12, 24, 36\}$  and the relation  $\leq$  be such that  $x \leq y$  if x divides y. Draw the Hasse diagram of  $\langle X, \leq \rangle$ .

**OR**

- Let  $F_x$  be the set of all one to one, onto mappings from X onto X where  $X = \{1, 2, 3\}$ . Find all the elements of  $F_x$  and find the inverse of each element.

**UNIT - III**

- Show that every cyclic group of order n is isomorphic to the group  $\langle z_n, t_n \rangle$ .
  - Prove that a subset  $S \neq \emptyset$  of G is a subgroup of  $\langle G, * \rangle$ . If for any pair of elements  $a, b \in S$ ,  $a * b^{-1} \in S$ .

**OR**

- Prove that a code can correct all combinations of K or fewer errors if and only if the minimum distance between any two code words is at least  $2K+1$ .

**UNIT - IV**

- Prove that a connected graph G is Euler if and only if all the vertices of G are even degree.

**OR**

- Explain travelling sales man's problem.

**UNIT - V**

- In how many ways can the 26 letters of the alphabet be permuted so that none of the patterns car, dog, pun or byte occurs?

**OR**

- Use generating functions to determine how many four element subsets of  $S = \{1, 2, 3, \dots, 15\}$  contain no consecutive integers.

\*\*\*\*\*