
# Pappus chain

From Wikipedia, the free encyclopedia

In geometry, the **Pappus chain** is a ring of circles between two tangent circles investigated by Pappus of Alexandria in the 3rd century AD.

## **Contents**

- 1 Construction
- 2 Properties
  - 2.1 Centers of the circles
    - 2.1.1 Ellipse
    - 2.1.2 Coordinates
  - 2.2 Radii of the circles
  - 2.3 Circle inversion
  - 2.4 Steiner chain
- 3 References
- 4 Bibliography
- 5 External links



A Pappus chain

## Construction

The arbelos is defined by two circles,  $C_{\rm U}$  and  $C_{\rm V}$ , which are tangent at the point  ${\bf A}$  and where  $C_{\rm U}$  is enclosed by  $C_{\rm V}$ . Let the radii of these two circles be denoted as  $r_{\rm U}$  and  $r_{\rm V}$ , respectively, and let their respective centers be the points  ${\bf U}$  and  ${\bf V}$ . The Pappus chain consists of the circles in the shaded grey region, which are externally tangent to  $C_{\rm U}$  (the inner circle) and internally tangent to  $C_{\rm V}$  (the outer circle). Let the radius, diameter and center point of the  $n^{\rm th}$  circle of the Pappus chain be denoted as  $r_n$ ,  $d_n$  and  ${\bf P}_n$ , respectively.

## **Properties**

#### Centers of the circles

### **Ellipse**

All the centers of the circles in the Pappus chain are located on a common ellipse, for the following reason. The sum of the distances from the  $n^{\rm th}$  circle of the Pappus chain to the two centers  ${\bf U}$  and  ${\bf V}$  of the arbelos circles equals a constant

$$\overline{\mathbf{P}_n\mathbf{U}}+\overline{\mathbf{P}_n\mathbf{V}}=(r_U+r_n)+(r_V-r_n)=r_U+r_V$$

Thus, the foci of this ellipse are **U** and **V**, the centers of the two circles that define the arbelos; these points correspond to the midpoints of the line segments **AB** and **AC**, respectively.

#### **Coordinates**

If r = AC/AB, then the center of the *n*th circle in the chain is:

$$(x_n,y_n)=\left(rac{r(1+r)}{2[n^2(1-r)^2+r]}\ ,\ rac{nr(1-r)}{n^2(1-r)^2+r}
ight)$$


#### Radii of the circles

If r = AC/AB, then the radius of the *n*th circle in the chain is:

$$r_n = rac{(1-r)r}{2[n^2(1-r)^2 + r]}$$

#### Circle inversion

The height  $h_n$  of the center of the  $n^{th}$  circle above the base diameter ACB equals n times  $d_n$ .[1] This may be shown by inverting in a circle centered on the tangent point A. The circle of inversion is chosen to intersect the  $n^{\text{th}}$ circle perpendicularly, so that the  $n^{\text{th}}$  circle is transformed into itself. The two arbelos circles,  $C_{\rm II}$  and  $C_{\rm V}$ , are transformed into parallel lines tangent to and sandwiching the  $n^{th}$  circle; hence, the other circles of the Pappus chain are transformed into similarly sandwiched circles of the same diameter. The initial circle  $C_0$  and the final circle  $C_n$  each contribute  $\frac{1}{2}d_n$ to the height  $h_n$ , whereas the circles  $C_1$ - $C_{n-1}$ each contribute  $d_n$ . Adding these contributions together yields the equation  $h_n = n d_n$ .



Under a particular inversion centered on  $\mathbf{A}$ , the four initial circles of the Pappus chain are transformed into a stack of four equally sized circles, sandwiched between two parallel lines. This accounts for the height formula  $h_n = n \ d_n$  and the fact that the original points of tangency lie on a common circle.

The same inversion can be used to show that the points where the circles of the Pappus chain are tangent to one another lie on a common circle. As noted above, the inversion centered at point  $\bf A$  transforms the

arbelos circles  $C_{\rm U}$  and  $C_{\rm V}$  into two parallel lines, and the circles of the Pappus chain into a stack of equally sized circles sandwiched between the two parallel lines. Hence, the points of tangency between the transformed circles lie on a line midway between the two parallel lines. Undoing the inversion in the circle, this line of tangent points is transformed back into a circle.

#### Steiner chain

In these properties of having centers on an ellipse and tangencies on a circle, the Pappus chain is analogous to the Steiner chain, in which finitely many circles are tangent to two circles.

## References

1. Ogilvy, pp. 54-55.

## **Bibliography**

- Ogilvy, C. S. (1990). Excursions in Geometry. Dover. pp. 54–55.
   ISBN 0-486-26530-7.
- Bankoff, L. (1981). "How did Pappus do it?". In Klarner, D. A. *The Mathematical Gardner*. Boston: Prindle, Weber, & Schmidt. pp. 112–118.
- Johnson, R. A. (1960). Advanced Euclidean Geometry: An elementary treatise on the geometry of the triangle and the circle (reprint of 1929 edition by Houghton Miflin ed.). New York: Dover Publications. pp. 116–117. ISBN 978-0-486-46237-0.
- Wells, D. (1991). *The Penguin Dictionary of Curious and Interesting Geometry*. New York: Penguin Books. pp. 5–6. ISBN 0-14-011813-6.

### **External links**

- Floer van Lamoen and Eric W. Weisstein. "Pappus Chain" (http://mathworld.wolfram.com/PappusChain.html). *MathWorld*.
- Tan, Stephen. "Arbelos" (http://www.math.ubc.ca/~cass/courses/m308/projects/tan/html/home.html).

Retrieved from "https://en.wikipedia.org/w/index.php?title=Pappus\_chain&oldid=749693691"

- This page was last edited on 15 November 2016, at 18:28.
- Text is available under the Creative Commons Attribution-ShareAlike

License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.