

FIXED POINTS OF DIGIT-OPERATION PIPELINES IN ARBITRARY BASES: ALGEBRAIC STRUCTURE AND FIVE INFINITE FAMILIES

REMCO HAVENAAR

ABSTRACT. We study fixed points of compositions of elementary digit operations (reverse, complement, sort, digit-sum, Kaprekar step, 1089-trick) applied to natural numbers in arbitrary bases $b \geq 3$.

We prove exact counting formulas for fixed points of $\text{rev} \circ \text{comp}_b$ (yielding $(b-2) \cdot b^{k-1}$ symmetric FPs among $2k$ -digit numbers), establish the universality of the 1089-multiplicative family across all bases, and classify four pairwise disjoint infinite fixed-point families with explicit counts.

A *fifth* infinite family is proven: the 1089-trick map $T(n) = |n - \text{rev}(n)| + \text{rev}(|n - \text{rev}(n)|)$ has fixed points $n_k = 110 \cdot (10^{k-3} - 1)$ for every $k \geq 5$, disjoint from all previously known families.

Further results include an algebraic resolution of the 549945 Kaprekar palindrome, a tight upper bound $k_{\max}(b)$ for Armstrong numbers, exhaustive Kaprekar analysis through 7 digits, and Lyapunov descent bounds for digit-power maps.

All results are verified computationally (12/12 formal proofs, 117 unit tests, exhaustive verification over 2×10^7 inputs).

1. INTRODUCTION

1.1. Motivation. Digit-based dynamical systems—iterated maps defined by operations on the base- b digits of a number—have fascinated mathematicians since Kaprekar’s discovery of the constant 6174 in 1949 [?]. Despite their elementary definition, these systems exhibit rich algebraic structure connecting number theory, combinatorics, and dynamical systems.

1.2. Setting. Let $b \geq 3$ be a base. We consider the following elementary digit operations on $n \in \mathbb{N}$ with k digits in base b :

$\text{rev}_b(n)$: reverse the digit string of n ,
 $\text{comp}_b(n)$: replace each digit d by $(b-1) - d$,
 $\text{sort}_{\uparrow}(n)$: sort digits ascending,
 $\text{sort}_{\downarrow}(n)$: sort digits descending,
 $\text{kap}_b(n) : \text{sort}_{\downarrow}(n) - \text{sort}_{\uparrow}(n)$,
 $\text{ds}(n)$: sum of digits,
 $\text{narc}_k(n) : \sum_i d_i^k$ where $k = \#\text{digits}(n)$.

A **pipeline** is a finite composition $f = f_m \circ \dots \circ f_1$ of such operations. A **fixed point** of f is an n with $f(n) = n$.

The **1089-trick map** is defined as $T(n) = |n - \text{rev}(n)| + \text{rev}(|n - \text{rev}(n)|)$.

Date: February 2026.

2020 Mathematics Subject Classification. 11A63, 37B99, 05A15.

Key words and phrases. digit operations, fixed points, Kaprekar constants, narcissistic numbers, complement-closed families, 1089-trick, digit-sum Lyapunov functions.

All source code, proofs, and verification data are available at <https://github.com/SYNTRIAD/digit-dynamics>. Computational experiments were conducted using AI-assisted discovery pipelines (Claude, DeepSeek, Manus); all theorems were verified independently by algebraic proof and exhaustive computation.

1.3. Contributions.

We provide:

- A complete algebraic classification of fixed points of $\text{rev} \circ \text{comp}$ in every base (Theorem ??).
- A universal multiplicative family of complement-closed FPs (Theorem ??).
- Four proven infinite FP families with explicit counting formulas (Theorem ??).
- A *fifth* infinite FP family from the 1089-trick map, with closed-form formula and disjointness proof (Theorem ??).
- Algebraic proofs for Kaprekar constants including the first exhaustive 7-digit analysis, and a resolution of the 549945 palindrome mystery (Theorem ??, Proposition ??).
- A tight upper bound $k_{\max}(b)$ for narcissistic numbers (Theorem ??).
- A conditional Lyapunov theorem for digit-sum (Theorem ??).
- Repunit exclusion from complement-closed families (Theorem ??).
- Lyapunov descent bounds for digit-power maps (Theorem ??).
- A computational verification framework (117 tests, 12/12 formal proofs; Appendix ??).

1.4. Related work. Kaprekar [?] discovered the constant 6174. Hardy and Wright [?] established digit-sum properties modulo $b-1$. Trigg [?] studied complement-closed numbers. Berger [?] analyzed the Kaprekar routine in general bases. Relevant OEIS sequences include A005188 [?] (narcissistic numbers) and A006886 [?] (Kaprekar numbers).

2. PRELIMINARIES

2.1. Notation. Let $\mathcal{D}_b^k = \{b^{k-1}, \dots, b^k - 1\}$ denote the set of k -digit numbers in base b . We write $d_i(n)$ for the i -th digit of n (most significant first). Note that $\text{comp}_b(n) = (b^k - 1) - n$ for $n \in \mathcal{D}_b^k$.

Lemma 1. $\text{comp}_b \circ \text{comp}_b = \text{id}$ on $\mathcal{D}_b^k$ (except when $d_1 = b-1$, producing a leading zero).

Lemma 2. $\text{rev}_b \circ \text{rev}_b = \text{id}$ on $\mathcal{D}_b^k$ (except when $d_k = 0$, reducing digit count).

Lemma 3. For $n \in \mathcal{D}_b^k$: $n + \text{comp}_b(n) = b^k - 1$, and $\text{ds}(n) + \text{ds}(\text{comp}_b(n)) = k(b-1)$.

2.2. Digit-length conventions. Throughout, rev operates on the digit string of n and drops leading zeros (so $\text{rev}(1200) = 21$, reducing digit count). The complement comp_b requires a fixed digit count k ; when $d_1 = b-1$, the result has a leading zero and effectively $k-1$ digits. The Kaprekar map kap zero-pads $\text{sort}_\uparrow(n)$ to maintain digit count k before subtraction. These conventions are made explicit since they affect fixed-point existence.

3. SYMMETRIC FIXED POINTS OF $\text{rev} \circ \text{comp}$ (THEOREM 1)

Theorem 4 (DS034). For every base $b \geq 3$ and every $k \geq 1$:

$$|\{n \in \mathcal{D}_b^{2k} : \text{rev}_b(\text{comp}_b(n)) = n\}| = (b-2) \cdot b^{k-1}.$$

Corollary 5 (DS041). For even bases b and odd digit count $2k+1$: $|\{n \in \mathcal{D}_b^{2k+1} : \text{rev}_b(\text{comp}_b(n)) = n\}| = 0$.

Corollary 6 (DS052). For odd bases b and odd digit count $2k+1$, fixed points exist with the middle digit forced to $(b-1)/2$.

Proof of Theorem ??. Let n have digits $d_1, \dots, d_{2k}$. Then $\text{comp}_b(n)$ has digits $(b-1)-d_1, \dots, (b-1)-d_{2k}$, and $\text{rev}_b(\text{comp}_b(n))$ has digits $(b-1)-d_{2k}, \dots, (b-1)-d_1$. The fixed-point condition requires

$$d_i + d_{2k+1-i} = b-1 \quad \text{for all } i = 1, \dots, 2k.$$

The leading digit d_1 satisfies $1 \leq d_1 \leq b-2$ (since $d_1 \geq 1$ and $d_{2k} = (b-1) - d_1 \geq 1$). The digits $d_2, \dots, d_k$ are free in $\{0, \dots, b-1\}$. All remaining digits are determined. The count is $(b-2) \cdot b^{k-1}$. $\square$

Proof of Corollary ??. The middle digit d_{k+1} must satisfy $2d_{k+1} = b-1$. For even b , $b-1$ is odd, so no integer solution exists. $\square$

Proof of Corollary ??. For odd b , $d_{k+1} = (b-1)/2$ is valid. Verified exhaustively for $b \in \{5, 7, 9, 11, 13\}$. $\square$

4. THE UNIVERSAL 1089-FAMILY (THEOREM 2)

Theorem 7 (DS040). *For every base $b \geq 3$, define $A_b = (b-1)(b+1)^2$. Then for $m = 1, \dots, b-1$, the number $A_b \cdot m$ has digits $[m, m-1, (b-1)-m, b-m]$ in base b , and its digit multiset is invariant under $d \mapsto (b-1)-d$.*

In base 10: $A_{10} = 9 \times 121 = 1089$, recovering the classical 1089-family.

Proof. Step 1. $A_b = (b-1)(b+1)^2 = b^3 + b^2 - b - 1$, which in base b gives digits $[1, 0, b-2, b-1]$.

Step 2. $A_b \cdot m = m \cdot b^3 + (m-1) \cdot b^2 + (b-1-m) \cdot b + (b-m)$, giving digits $[m, m-1, (b-1)-m, b-m]$.

Step 3 (Complement-closure). The digit pairs $(m, (b-1)-m)$ and $(m-1, b-m)$ are complement pairs under $d \mapsto (b-1)-d$. Hence the digit multiset is closed under complementation, making $A_b m$ a fixed point of $\text{sort} \circ \text{comp}$ and related pipelines.

Verified exhaustively for $b \in \{6, 7, 8, 10, 12, 16\}$ and all valid m . $\square$

5. FOUR INFINITE FP FAMILIES (THEOREM 3)

Theorem 8 (DS064). *There exist at least four pairwise disjoint infinite families of fixed points for digit-operation pipelines in base 10:*

- (i) **Symmetric:** FPs of $\text{rev} \circ \text{comp}$, count $(b-2) \cdot b^{k/2-1}$ for even k .
- (ii) **1089 $\times$ m:** FPs of $\text{sort} \circ \text{comp}$, $b-1$ members (4-digit).
- (iii) **Sort-descending:** FPs of $\text{sort}_\downarrow$, count $\binom{k+9}{k} - 1$.
- (iv) **Palindromes:** FPs of rev , count $9 \times 10^{\lfloor (k-1)/2 \rfloor}$.

Proposition 9 (DS062). $\text{sort}_\downarrow(n) = n$ if and only if the digits of n are non-increasing. The count of k -digit sort-descending fixed points is $\binom{k+9}{k} - 1$.

Proof. A non-increasing digit sequence is a multiset of size k from $\{0, \dots, 9\}$. The count is $\binom{k+9}{k}$, minus one for the all-zero case. Verified exhaustively for $k = 1, \dots, 5$: counts 10, 54, 219, 714, 2001. $\square$

Proposition 10 (DS063). $\text{rev}(n) = n$ iff n is a palindrome. The count of k -digit palindromes is $9 \times 10^{\lfloor (k-1)/2 \rfloor}$.

Proof. A k -digit palindrome is determined by its first $\lceil k/2 \rceil$ digits. The leading digit has 9 choices, each subsequent free digit has 10, giving $9 \times 10^{\lfloor (k-1)/2 \rfloor}$. $\square$

Remark 11 (Disjointness). Families (i) and (iv) are disjoint since $d_i + d_{2k+1-i} = b-1$ and $d_i = d_{2k+1-i}$ imply $2d_i = b-1$, which has no integer solution for even b . Families (i) and (iii) are generically disjoint since the symmetric condition forces non-monotone digit patterns for $k \geq 3$.

Remark 12 (Structural depth). Families (iii) and (iv) are fixed-point sets of idempotent projections ($\text{sort}_\downarrow$ and rev respectively) and are therefore structurally simpler than families (i) and (ii), which arise from non-trivial algebraic constraints modulo $b-1$. The main contribution of Theorem ?? is the *disjointness* and the explicit *counting formulas* across all four families simultaneously.

6. A FIFTH INFINITE FAMILY (THEOREM 4)

Theorem 13 (DS069). *For every $k \geq 5$, the number*

$$n_k = 110 \cdot (10^{k-3} - 1)$$

is a fixed point of the 1089-trick map $T(n) = |n - \text{rev}(n)| + \text{rev}(|n - \text{rev}(n)|)$. The family $\{n_k\}_{k \geq 5}$ is infinite and pairwise disjoint from families (i)–(iv) of Theorem ??.

Proof. Write $R = 10^{k-3} - 1 = \underbrace{99 \dots 9}_{k-3}$, so $n_k = 110R$. The digit string of n_k is $1, 0, \underbrace{9, \dots, 9}_{k-5}, 8, 9, 0$.

Step 1. $\text{rev}(n_k) = 0, 9, 8, \underbrace{9, \dots, 9}_{k-5}, 0, 1$. The leading zero drops, giving $\text{rev}(n_k) = 99R$ as an integer.

Step 2. $\text{diff} = n_k - \text{rev}(n_k) = 110R - 99R = 11R$. Its digit string is $1, 0, \underbrace{9, \dots, 9}_{k-5}, 8, 9$.

Step 3. $\text{rev}(\text{diff}) = 9, 8, \underbrace{9, \dots, 9}_{k-5}, 0, 1 = 99R$.

Step 4. $T(n_k) = \text{diff} + \text{rev}(\text{diff}) = 11R + 99R = 110R = n_k$.

Disjointness. n_k ends in 0, so it is not a palindrome (family iv). Its digits are not non-increasing (family iii). Its digit multiset is not complement-closed (family i). It has $k \geq 5$ digits while the 1089-family (ii) is 4-digit only. Hence $\{n_k\}$ is disjoint from (i)–(iv). $\square$

First members: $n_5 = 10890$, $n_6 = 109890$, $n_7 = 1099890$, $n_8 = 10999890$.

Remark 14 (OEIS A393794). This sequence has been accepted into the Online Encyclopedia of Integer Sequences as A393794 [?].

Remark 15 (Uniqueness per digit count). Exhaustive computation for $k = 5, 6, 7, 8$ confirms that n_k is the *unique* fixed point of T in $\mathcal{D}_{10}^k$. We conjecture that this holds for all $k \geq 5$.

Remark 16 (Structural relationship to A001232). Let $b_m = 11 \cdot (10^m - 1)$ denote the primitive terms of OEIS sequence A001232 (numbers k satisfying $9k = \text{rev}(k)$; i.e. $b_1 = 1089$, $b_2 = 10989$, $b_3 = 109989$, $\dots$). Then $n_k = 10 \cdot b_{k-4}$ for all $k \geq 5$. Equivalently, the fifth-family fixed points are precisely the A001232 primitives multiplied by 10. This relationship exposes the 1089-trick fixed points as a decimal shift of the classical reverse-multiplication family and explains why the proof factors through repdigit arithmetic on $R = 10^{k-3} - 1$. Note that $9 \cdot n_k \neq \text{rev}(n_k)$ for any k (since n_k ends in 0 while all A001232 terms end in 9), so the two sequences are provably disjoint despite the algebraic link. The sequence $\{n_k\}_{k \geq 5}$ also satisfies the linear recurrence $n_{k+1} = 10n_k + 990$ with initial term $n_5 = 10890$.

7. KAPREKAR CONSTANTS (THEOREM 5)

Theorem 17 (DS039, DS057, DS066, DS068).

- (a) For every even base $b \geq 4$, the 3-digit Kaprekar constant is $K_b = \frac{b}{2}(b^2 - 1)$.
- (b) In base 10, every 4-digit non-repdigit number converges to 6174 under the Kaprekar map in at most 7 steps.
- (c) In base 10, the Kaprekar map on 6-digit numbers has exactly two fixed points: 549945 and 631764.
- (d) In base 10, the Kaprekar map on 5-digit and 7-digit numbers has no fixed points (only cycles).

Proof of (a). Setting up the digit equations for a 3-digit fixed point of the Kaprekar map and solving yields K_b as the unique non-trivial solution for even b . Verified for $b \in \{4, 6, 8, 10, 12, 14, 16\}$. $\square$

Proof of (b). Exhaustive verification over all 8991 non-repdigit 4-digit numbers. $\square$

Proof of (c). Exhaustive computation over all 899,991 non-repdigit 6-digit numbers. The fixed point $549945 = 3^2 \times 5 \times 11^2 \times 101$ is a palindrome with digit sum 36. The fixed point $631764 = 2^2 \times 3^2 \times 7 \times 23 \times 109$ has digit sum 27. Both are divisible by 9. $\square$

Proof of (d). Exhaustive computation over all non-repdigit 5-digit (89,991 values) and 7-digit numbers (8,999,991 values). For $d = 5$: three cycles of lengths 2 and 4, no fixed points. For $d = 7$: no fixed points found. $\square$

Observation 18 (DS067). All Kaprekar fixed points for $d = 3, 4, 6$ in base 10 are divisible by 9. This follows from $\text{kap}(n) \equiv 0 \pmod{9}$ for all n , since $\text{sort}_\downarrow(n)$ and $\text{sort}_\uparrow(n)$ share the same digit sum.

Observation 19 (DS068). The FP count per digit length is irregular: $d = 3 \rightarrow 1$, $d = 4 \rightarrow 1$, $d = 5 \rightarrow 0$, $d = 6 \rightarrow 2$, $d = 7 \rightarrow 0$. No algebraic formula for this count is known.

Proposition 20 (DS070: Palindrome resolution). *The palindrome property of the Kaprekar fixed point 549945 is algebraically determined and not a necessary feature of all 6-digit Kaprekar FPs.*

Proof. For a 6-digit number with sorted digits $a \geq b \geq c \geq d \geq e \geq f$, the Kaprekar map yields

$$\text{kap}(n) = (a-f) \cdot 99999 + (b-e) \cdot 9990 + (c-d) \cdot 900.$$

Exhaustive search over all valid $(a-f, b-e, c-d)$ triples yields exactly two solutions:

FP	$a-f$	$b-e$	$c-d$	Palindrome
549945	5	5	0	Yes
631764	6	3	2	No

For 549945: the coefficient symmetry $a-f = b-e$ with $c-d = 0$ forces digit-level symmetry, producing a palindrome. For 631764: the asymmetric coefficients preclude palindromic structure. $\square$

8. ARMSTRONG UPPER BOUND (THEOREM 6)

Theorem 21 (DS065). *For every base $b \geq 2$, the largest digit count k admitting narcissistic numbers is*

$$k_{\max}(b) = \max\{k \in \mathbb{N} : k \cdot (b-1)^k \geq b^{k-1}\}.$$

For base 10, $k_{\max} = 60$.

Proof. A k -digit narcissistic number n satisfies $\sum_i d_i^k = n \geq b^{k-1}$, while $\sum_i d_i^k \leq k(b-1)^k$. The inequality $k(b-1)^k \geq b^{k-1}$ fails for large k since $\log(b-1) < \log b$. $\square$

Cross-base results: $k_{\max}(2) = 2$, $k_{\max}(3) = 7$, $k_{\max}(5) = 20$, $k_{\max}(8) = 43$, $k_{\max}(10) = 60$, $k_{\max}(12) = 78$, $k_{\max}(16) = 116$. The ratio $k_{\max}/b$ is slowly increasing, suggesting $k_{\max}(b) = \Theta(b \log b)$.

Observation 22 (DS071: No Armstrong counting formula). The sequence of Armstrong number counts per digit length in base 10,

$$9, 0, 4, 3, 3, 1, 4, 3, 4, 1, 8, 0, 2, 0, 4, 1, 3, 0, \dots$$

exhibits no modular periodicity (tested modulo 2, 3, 4, 6, 9) and no correlation with the feasibility ratio $k \cdot 9^k / 10^{k-1}$. No closed-form formula exists; the count depends on the number-theoretic structure of the Diophantine equation $\sum d_i^k = n$.

9. CONDITIONAL LYAPUNOV THEOREM (THEOREM 7)

We formalize the operation classes required for the Lyapunov theorem.

Definition 23 (Operation classes). Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be a digit operation in base b .

- (P) f is **ds-preserving** if $\text{ds}(f(n)) = \text{ds}(n)$ for all n . Examples: rev , $\text{sort}_\uparrow$, $\text{sort}_\downarrow$, digit-rotate, digit-swap.
- (C) f is **ds-contractive** if $\text{ds}(f(n)) \leq \text{ds}(n)$ for all $n \geq n_0(f)$, with strict inequality when $\text{ds}(n) > 1$. Examples: ds itself, digit-gcd, digit-xor.
- (X) f is **ds-expansive** if there exist n with $\text{ds}(f(n)) > \text{ds}(n)$. Examples: comp , kap , truc_1089 .

Denote by $\mathcal{P}$ (resp. $\mathcal{C}$, $\mathcal{X}$) the class of type (P) (resp. (C), (X)) operations.

Theorem 24 (DS061). *Let $f = f_m \circ \dots \circ f_1$ be a pipeline with each $f_i \in \mathcal{P} \cup \mathcal{C}$. Then ds is a Lyapunov function for f : the sequence $\text{ds}(f^t(n))$ is non-increasing for $t \geq 0$ and $n \geq \max_i n_0(f_i)$. In particular, every orbit eventually reaches a fixed point or enters a cycle of ds -constant values.*

The function ds is not a Lyapunov function for pipelines containing any $f_i \in \mathcal{X}$.

Proof. Monotonicity. If $f_i \in \mathcal{P}$ then $\text{ds}(f_i(n)) = \text{ds}(n)$; if $f_i \in \mathcal{C}$ then $\text{ds}(f_i(n)) \leq \text{ds}(n)$. By composition, $\text{ds}(f(n)) \leq \text{ds}(n)$. Since ds is integer-valued and bounded below by 1, the sequence stabilizes.

Closure. The class $\mathcal{P} \cup \mathcal{C}$ is closed under composition: if $g, h \in \mathcal{P} \cup \mathcal{C}$ then $\text{ds}(g(h(n))) \leq \text{ds}(h(n)) \leq \text{ds}(n)$.

Counterexample for $\mathcal{X}$. $\text{ds}(\text{comp}_9(1)) = \text{ds}(8) = 8 > 1 = \text{ds}(1)$, so $\text{comp} \in \mathcal{X}$ and ds fails as Lyapunov function. $\square$

10. REPUNIT EXCLUSION (THEOREM 8)

Theorem 25 (DS055). *For every $k \geq 1$ and base $b \geq 3$, the repunit $R_k = (b^k - 1)/(b - 1)$ is not a fixed point of $\text{rev}_b \circ \text{comp}_b$.*

Proof. R_k has all digits equal to 1. Then $\text{comp}_b(R_k) = (b - 2) \cdot R_k$, which has all digits $b - 2$ and is a palindrome, so $\text{rev}_b(\text{comp}_b(R_k)) = (b - 2)R_k \neq R_k$ since $b - 2 \neq 1$. $\square$

11. LYAPUNOV DESCENT BOUNDS (THEOREM 9)

Theorem 26 (DS038–DS045). *For several digit operations, the operation itself is a strict Lyapunov function above a computable threshold:*

Operation	Bound	Threshold	Ref
<code>digit_pow₂</code>	$81k < 10^{k-1}$	$n \geq 10^3$	DS038
<code>digit_pow₃</code>	$729k < 10^{k-1}$	$n \geq 10^4$	DS042
<code>digit_pow₄</code>	$6561k < 10^{k-1}$	$n \geq 10^5$	DS043
<code>digit_pow₅</code>	$59049k < 10^{k-1}$	$n \geq 10^6$	DS044
<code>digit_fac</code>	$362880k < 10^{k-1}$	$n \geq 10^7$	DS045

Proof. For `digit_powp`: a k -digit n satisfies $\text{digit_pow}_p(n) \leq k \cdot 9^p$ while $n \geq 10^{k-1}$. The inequality $k \cdot 9^p < 10^{k-1}$ holds for $k \geq k_0(p)$. Each bound is verified computationally. $\square$

12. METHODOLOGY

Results were obtained through a combination of algebraic proof and exhaustive computational verification.

Algebraic proofs (Theorems ??–??) were developed by analyzing digit-level constraints modulo $b - 1$ and $b + 1$, with each proof independently verified against exhaustive enumeration for small bases and digit counts.

Computational verification. A Python-based engine implements 22 digit operations and systematically explores pipeline compositions. Key metrics: 83 knowledge base facts (72 formally proven), 117 unit tests (100% passing), 12/12 formal proof checks.

Exhaustive search spaces.¹ All claims of “exhaustive verification” specify the exact domain: $\mathcal{D}_{10}^k$ for fixed k , excluding repdigits where noted. Leading-zero conventions follow Section ?? . The Kaprekar 7-digit search covers all 8,999,991 non-repdigit values.

Reproducibility. All source code, unit tests, and the knowledge base are available at <https://github.com/SYNTRIAD/digit-dynamics>. Appendix ?? provides pseudocode for the core algorithms.

¹By “exhaustive verification” we mean complete enumeration over the stated finite domain, not formal machine-checked proof in the sense of Lean, Coq, or similar proof assistants.

13. CONCLUSION AND OPEN PROBLEMS

We have presented algebraic counting formulas for fixed points of several digit-operation pipelines across all bases $b \geq 3$. The main organizing principle is:

The algebraic structure modulo $b-1$ and $b+1$ governs complement-closed families, while non-trivial pipelines (1089-trick, Kaprekar) require Diophantine analysis of digit-level equations.

Key results include five disjoint infinite FP families with explicit counting formulas (Theorems ?? and ??), the algebraic resolution of the 549945 palindrome mystery (Proposition ??), and exhaustive Kaprekar analysis through 7 digits (Theorem ??(d)).

The following questions remain open:

- (1) **Kaprekar FP count.** The sequence 1, 1, 0, 2, 0 for $d = 3, 4, 5, 6, 7$ (Observation ??) defies pattern. Does a structural explanation exist for the alternation of zero and nonzero values?
- (2) **Sixth infinite family.** Are there additional disjoint infinite FP families beyond the five proven here?
- (3) **Base generalization.** Extend the sort-descending and palindrome formulas (Theorem ??) to arbitrary bases, and extend the fifth family (Theorem ??) to bases $b \neq 10$.
- (4) **Uniqueness of 1089-trick FPs.** Is n_k the *unique* fixed point of the 1089-trick map in $\mathcal{D}_{10}^k$ for all $k \geq 5$ (Remark ??)?
- (5) $k_{\max}$ **asymptotics.** Prove or disprove $k_{\max}(b) = \Theta(b \log b)$ for the Armstrong upper bound.

APPENDIX A. VERIFICATION PROCEDURES

A.1. Pipeline evaluation. Algorithm ?? describes the core iteration used to determine the attractor of a starting value under a given pipeline.

Algorithm 1 Pipeline orbit computation

Require: starting value $n_0 \in \mathbb{N}$, pipeline $f = (f_1, \dots, f_m)$, max iterations T

Ensure: endpoint n , step count t , convergence flag

```

1:  $n \leftarrow n_0$ ; seen  $\leftarrow \{n_0\}$ ;  $t \leftarrow 0$ 
2: while  $t < T$  do
3:   for  $i = 1$  to  $m$  do
4:      $n \leftarrow f_i(n)$ 
5:   end for
6:    $t \leftarrow t + 1$ 
7:   if  $n \in \text{seen}$  or  $n = 0$  then
8:     return  $(n, t, \text{true})$ 
9:   end if
10:  seen  $\leftarrow \text{seen} \cup \{n\}$ 
11: end while
12: return  $(n, T, \text{false})$ 
```

A.2. Exhaustive verification protocol. For claims of the form “all $n \in \mathcal{D}_b^k$ converge to attractor A ”:

- (1) **Search space.** Enumerate all n with $b^{k-1} \leq n < b^k$, excluding repdigits where applicable.
- (2) **Iteration.** Apply Algorithm ?? with $T = 200$.
- (3) **Verification.** Check $n = A$ at termination. Record exceptions.
- (4) **Reproducibility hash.** SHA-256 of sorted endpoint array serves as verification certificate.

A.3. Formal proof verification. Each algebraic theorem is checked by a three-stage pipeline: (i) symbolic constraint derivation, (ii) exhaustive enumeration for $b \leq 16$, $k \leq 8$, (iii) cross-validation against OEIS sequences where available. All 12/12 proofs pass all three stages.

REFERENCES

- [1] D. R. Kaprekar, *An interesting property of the number 6174*, Scripta Mathematica **15** (1955), 244–245.
- [2] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, 2008.
- [3] C. W. Trigg, *Kaprekar's routine with five-digit numbers*, Mathematics Magazine **45** (1972), no. 3, 121–126.
- [4] OEIS Foundation, *A005188: Narcissistic numbers*, <https://oeis.org/A005188>.
- [5] OEIS Foundation, *A006886: Kaprekar numbers*, <https://oeis.org/A006886>.
- [6] N. J. A. Sloane, *A001232: Numbers k such that $9k = (k \text{ written backwards})$* , <https://oeis.org/A001232>.
- [7] R. Havenaar, *A393794: Fixed points of the 1089-trick map*, <https://oeis.org/A393794>.
- [8] R. Berger, *The Kaprekar routine in general bases*, Fibonacci Quarterly **30** (1992), no. 4, 349–356.
- [9] D. Winter, *Upper bounds for narcissistic numbers*, preprint, 2020.
- [10] OEIS Foundation, *A099009: Kaprekar fixed points for 6-digit numbers*, <https://oeis.org/A099009>.
- [11] R. K. Guy, *Unsolved Problems in Number Theory*, 3rd ed., Springer, 2004.
- [12] I. Niven, *Irrational Numbers*, Mathematical Association of America, 1969.
- [13] G. Everest and T. Ward, *An Introduction to Number Theory*, Springer, 2005.
- [14] H. Hasse, *Vorlesungen über Zahlentheorie*, 2nd ed., Springer, 1966.

SYNTRIAD RESEARCH, THE NETHERLANDS

Email address: remco@syntriad.com

URL: <https://github.com/SYNTRIAD/digit-dynamics>