

# Problem Set 3

Shirley Yuan

In collaboration with Zeshun, Shirley and Zunda

$$\mathbf{2} \quad D = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}.$$

Let  $\det(\lambda I - D) = 0$

$$\begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -2 \\ 0 & 0 & \lambda \end{vmatrix} = \lambda^3 = 0$$

$$\Rightarrow \lambda = 0$$

$\therefore$  Algebraic multiplicity is 3. The corresponding eigenvector is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

$\therefore$  Geometric multiplicity is 1.

**4**

**i)** Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , then  $A^H = A$  implies that  $b = \bar{c}$ .

Recall that if  $z \in \mathbb{C}$ , then  $z\bar{z} = |z|^2 \geq 0$

Now,  $p(\lambda) = \lambda^2 - (a+d)\lambda + (ad-bc)$

$$\Delta = (a+d)^2 - 4(ad-bc) = (a-d)^2 + 4b\bar{b} = (a-d)^2 + 4|b|^2 \geq 0$$

$\Rightarrow$  Real roots.

**2** Suppose  $\lambda$  is an eigenvalue of  $A$  where  $A^H = -A$ .  $x$  is the corresponding eigenvector.

Then,  $\langle Ax, x \rangle = \langle \lambda x, x \rangle = \lambda \langle x, x \rangle$ .

Also,  $\langle Ax, x \rangle = \langle x, A^H x \rangle = \langle x, -Ax \rangle = -\langle x, \lambda x \rangle = -\lambda \langle x, x \rangle$ .

So we have  $\bar{\lambda} = -\lambda$

$\Rightarrow \lambda$  is pure imaginary.

**6** Suppose  $A$  is an upper triangular matrix.

$$A = \begin{bmatrix} a_1 & & & * \\ & a_2 & & \\ & & \ddots & \\ 0 & & & a_n \end{bmatrix}$$

Then the characteristic polynomial is

$$\det(zI - A) = \begin{vmatrix} z - a_1 & & & * \\ & z - a_2 & & \\ & & \ddots & \\ 0 & & & z - a_n \end{vmatrix} = \prod_{i=1}^n (z - a_i) = 0$$

Note that this polynomial has n zeros, which are  $a_1, a_2, \dots, a_n$  respectively.

The case of upper triangular matrix is the same.

8

1 Since  $V = \text{span}(s)$ , it suffices to show that the four vectors are linearly independent.

Let  $a \sin(x) + b \cos(x) + c \sin(2x) + d \cos(2x) = 0, \quad \forall x \in \mathbb{R}$ .

Let  $x = 0$ :  $b + d = 0$

Let  $x = \frac{\pi}{2}$ :  $a - d = 0$

Let  $x = \pi$ :  $-b + d = 0$

Let  $x = \frac{\pi}{4}$ :  $a \sin(\frac{\pi}{4}) + b \cos(\frac{\pi}{4}) + c \sin(\frac{\pi}{2}) + d \cos(\frac{\pi}{2}) = 0$

From the above four conditions we can get  $a = 0, b = 0, c = 0, d = 0$ .

Since the only case that can let  $a \sin(x) + b \cos(x) + c \sin(2x) + d \cos(2x) = 0, \quad \forall x \in \mathbb{R}$ , is when  $a = b = c = d = 0$ ,

$\Rightarrow$  They are linearly independent.

$$2 \quad D = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 2 & 0 \end{bmatrix}$$

3  $V_1 = \{\sin x, \cos x\}, V_2 = \{\sin 2x, \cos 2x\}$

13 To diagonalize A, we first need to find eigenvalues and eigenvectors.

$$p(\lambda) = \lambda^2 - 1.4\lambda + 0.4 = 0$$

$$\Rightarrow \lambda_1 = 1, \lambda_2 = \frac{2}{5}$$

And  $\Sigma_1 = \text{span}([2, 1]^T), \Sigma_2 = \text{span}([1, -1]^T)$

So A is semisimple.

$$\text{Let } p = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}, \text{ then } p^{-1} = \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{2}{3} \end{bmatrix}.$$

$$D = \begin{bmatrix} 1 & 0 \\ 0 & 0.4 \end{bmatrix}$$

Then  $D = p^{-1}Ap$ .

15 Since  $A \in M_n(\mathbb{F})$  is semisimple, we can diagonalize  $A = pDp^{-1}$ , where D is diagonalized, and  $\{\lambda_i\}_1^n$  are the diagonal entries of D.

$$\begin{aligned}
\text{Now, } f(A) &= f(pDp^{-1}) \\
&= a_0I + a_1pDp^{-1} + \dots + a_n pD^n p^{-1} \\
&= p[a_0I + a_1D + \dots + a_n D^n]p^{-1} \\
&= pf(D)p^{-1}
\end{aligned}$$

Observe that  $f(A)$  and  $f(D)$  are similar, so they have the same eigenvalues.

Also note that  $f(D)$  is also diagonal, so each entry along the diagonal is  $f(D)_{ii} = a_0 + a_1 d_{ii} + \dots + a_n d_{ii}^n = f(d_{ii})$ , where  $D = [d_{ij}]_{ij}$

Hence, the eigenvalues of  $f(D)$  are just its diagonals, which are  $\{f(\lambda_1), f(\lambda_2), \dots, f(\lambda_n)\}$

**16**

$$\begin{aligned}
& \mathbf{1} \quad A = pD^n p^{-1} \\
&= \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0.4^n \end{bmatrix} \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{2}{3} \end{bmatrix} \\
&\therefore \lim_{n \rightarrow \infty} A^n = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{2}{3} \end{bmatrix} \\
&= \begin{bmatrix} \frac{2}{3} & \frac{2}{3} \\ \frac{1}{3} & \frac{1}{3} \end{bmatrix} \\
&\text{Let } B = \begin{bmatrix} \frac{2}{3} & \frac{2}{3} \\ \frac{1}{3} & \frac{1}{3} \end{bmatrix}, \text{ then it follows immediately from the definition of limit.}
\end{aligned}$$

**2** The choice of norm does not affect the answer.

**3** Let  $f(x) = 3 + 5x + x^3$ , then the eigenvalues of  $f(A)$  are  $f(\lambda_1) = f(1) = 9, f(\lambda_2) = f(0.4) = 5.064$ .

**18** Take  $\vec{y}$  an eigenvector corresponding to  $\lambda$ . Then,

$$\begin{aligned}
x^T A y &= x^T \lambda y = (\lambda x^T y) \\
&\Rightarrow x^T A = \lambda x^T
\end{aligned}$$

**20** Let  $B = U^H A U$ , then,

$$B^H = U^H A^H U = U^H A U = B, \text{ since } A^H = A$$

**24**

$$\mathbf{1} \quad p(\vec{x}) = \frac{\langle x, Ax \rangle}{||x||^2}$$

Observe that the denominator is always a real number. Hence to show that  $p(\vec{x}) \in \mathbb{R}$ , it suffices to show that  $\langle x, Ax \rangle \in \mathbb{R}$ .

$$\text{Now } \langle x, Ax \rangle = \langle A^H x, x \rangle = -\langle Ax, x \rangle$$

$$\text{Since by definition, } \langle x, Ax \rangle = \langle A\bar{x}, x \rangle, \text{ we have } \langle Ax, x \rangle = \langle A\bar{x}, x \rangle \in \mathbb{R}$$

This implies

$$\langle x, Ax \rangle \in \mathbb{R}$$

$$\Rightarrow p(x) \in \mathbb{R}$$

**2** If  $A^H = -A$ , then

$$\langle x, Ax \rangle = \langle A^H, x \rangle = -\langle Ax, x \rangle$$

$$\text{Also, } \langle x, Ax \rangle = \langle A\bar{x}, x \rangle$$

$$\therefore \langle A\bar{x}, x \rangle = -\langle Ax, x \rangle$$

This implies  $\langle x, Ax \rangle = \langle A\bar{x}, x \rangle \in \mathbb{C} \setminus \mathbb{R} \cup \{0\}$

Hence  $p(\bar{x}) = \frac{\langle x, Ax \rangle}{\|x\|^2}$  is pure imaginary number.

**25**

**1** Since  $A \in M_n(\mathbb{C})$  is a normal matrix, its eigenspace  $\{x_1, x_2, \dots, x_n\}$  spans  $\mathbb{C}^n$ .

Observe that  $\forall j = 1, 2, \dots, n$ ,

$$(x_1x_1^H + x_2x_2^H + \dots + x_nx_n^H)x_j = x_1x_1^Hx_j + x_2x_2^Hx_j + \dots + x_nx_n^Hx_j = x_j$$

This holds for any  $j$ .

Since  $\{x_1, x_2, \dots, x_n\}$  spans  $\mathbb{C}^n$ ,  $\forall \vec{v} = \mathbb{C}^n$ ,  $\vec{v} = \sum a_i \vec{x}_i$

Let  $B = x_1x_1^H + x_2x_2^H + \dots + x_nx_n^H$ , then  $B\vec{v} = \sum a_i B\vec{x}_i = \sum a_i \vec{x}_i = \vec{v}$ .

Let  $\vec{v} = \vec{e}_1, \vec{e}_2, \dots, \vec{e}_n$  respective, then we get

$$Be_1 = e_1, Be_2 = e_2, \dots, Be_n = e_n$$

Hence  $B = I$

**2** Since  $A$  is a normal matrix and  $\{x_1, x_2, \dots, x_n\}$  forms an orthonormal eigenbasis,  $A$  admits a diagonalization.

$A = pDp^{-1} = pDp^H$ , where

$$p = [x_1, x_2, \dots, x_n] \quad D = \begin{bmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_n \end{bmatrix}$$

$$p^{-1} = p^H = \begin{bmatrix} x_1^H \\ x_2^H \\ \vdots \\ x_n^H \end{bmatrix}, \text{ since } p \text{ is an orthonormal matrix.}$$

$$\text{Hence, } A = [x_1, x_2, \dots, x_n] \begin{bmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_n \end{bmatrix} \begin{bmatrix} x_1^H \\ x_2^H \\ \vdots \\ x_n^H \end{bmatrix} = \sum \lambda_i x_i x_i^H$$

**27** Suppose 
$$\begin{bmatrix} a_{11} & \dots & a_{1n} \\ a_{21} & \lambda_2 & a_{2n} \\ \vdots & & \vdots \\ a_{n1} & & a_{nn} \end{bmatrix}$$

By definition,  $\forall x, x^H A x > 0$

Now, let  $x = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} = e_i^{-1}$

then  $e_1^H A e_1 = \begin{bmatrix} 1 & 0 & \dots & 0 \end{bmatrix} \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{bmatrix} = a_{11} > 0$

Similarly, let  $x = e_2, e_3, \dots, e_n$

we have  $a_{22} > 0, a_{33} > 0, \dots, a_{nn} > 0$

Here all diagonal elements are positive and real.

## 28 Proof:

First we introduce the following lemmas used in the proof.

- Lemma 1: The diagonals of a positive semi-definite matrix are greater than or equal to zero. (Proof similar to exercise 4.27)
- Lemma 2:  $tr(AB) = tr(BA)$  (Proof can be found in Problem Set 2)
- Lemma 3: If  $A \in M_n(\mathbb{F})$  is a positive semi-definite matrix,  $D \in M_n(\mathbb{F})$  is a diagonal matrix with non-negative diagonals, then  $0 \leq tr(AD) \leq tr(A)tr(D)$ .

*Proof.* Suppose

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & & & \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

,

$$D = \begin{bmatrix} d_{11} & & & \\ & d_{22} & & \\ & & \ddots & \\ & & & d_{nn} \end{bmatrix}$$

,

then  $tr(AD) = \sum_{i=1}^n a_{ii}d_i \geq 0$ , since  $a_{ii} \geq 0$  and  $d_i \geq 0$  for  $\forall i$

$$tr(A)tr(D) = (\sum_{i=1}^n a_{ii})(\sum_{i=1}^n d_i) = \sum_{i=1}^n a_{ii}d_i + \sum_{i \neq j} a_{ii}d_j \geq \sum_{i=1}^n a_{ii}d_i$$

$$\Rightarrow tr(A)tr(D) \geq tr(AD) \geq 0$$

□

Now since  $B$  is a positive semi-definite matrix, it admits a diagonalization s.t.  $B = PDP^{-1} = PDP^H$ , where

$$P = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}$$

is an orthonormal eigenbasis,

$$D = \begin{bmatrix} d_{11} & & & \\ & d_{22} & & \\ & & \ddots & \\ & & & d_{nn} \end{bmatrix}$$

is diagonal matrix with  $d_i \geq 0 \quad \forall i$ .

$$\begin{aligned} \text{Then } tr(AB) &= tr(APDP^H) = tr(P^H APD) \leq tr(P^H AP)tr(D) \\ &= tr(APP^H)tr(D) = tr(A)tr(D) = tr(A)tr(B). \end{aligned}$$

Meanwhile,  $\|AB\|_F^2 = tr(AA^H BB^H) \leq tr(AA^H) tr(BB^H) = \|A\|_F \|B\|_F^2$ , which makes  $\|\cdot\|_F$  a matrix norm.

### 31

1 Suppose  $A$  has rank  $r$ , then  $A^H A$  is positive definite and has  $r$  distinct eigenvalues.

Let  $s = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$  be an orthonormal eigenspace of  $A^H A$ , and  $\{\sigma_1^2, \sigma_2^2, \dots, \sigma_n^2\}$  be the corresponding eigenvalues, where  $\sigma_1^2 \geq \sigma_2^2 \geq \dots \geq \sigma_n^2$ .

Since  $s$  spans  $\mathbb{F}^n$ ,  $\forall \vec{x} \in \mathbb{F}^n$ , we have

$$\vec{x} = \sum_{i=1}^n c_i \vec{v}_i, c_i \in \mathbb{F}, \forall i, \text{ and}$$

$$\|\vec{x}\|_2 = \sqrt{(\sum c_i v_i^T)(\sum c_i v_i)} = \sqrt{(\sum c_i^2)}$$

Hence if  $\|\vec{x}\|_2 = 1$ , then  $\sum_{i=1}^n c_i^2 = 1$

Now, observe that  $\|Ax\|_2^2 = \langle Ax, Ax \rangle = (Ax)^H Ax = x^H A^H Ax$

$$= (\sum_{i=1}^n c_i \vec{v}_i^H)(A^H A)(\sum_{i=1}^n c_i \vec{v}_i)$$

$$= (\sum_{i=1}^n c_i \vec{v}_i^H)(\sum_{i=1}^n c_i A^H A \vec{v}_i)$$

$$= (\sum_{i=1}^n c_i \vec{v}_i^H)(\sum_{i=1}^n c_i \sigma_i^2 \vec{v}_i) = \sum c_i^2 \sigma_i^2, \text{ where } s = \{v_1, v_2, \dots, v_n\}$$

Note that when  $\sigma_i^2 = 1$ , and  $\sigma_1^2 \geq \sigma_2^2 \geq \dots \geq \sigma_n^2$ ,

$$\sum c_i^2 \sigma_i^2 \leq \sigma_1^2$$

$$\text{Hence, } \|A\|_2^2 = \sup_{\|\vec{x}\|_2=1} \|Ax\|_2^2 = \sigma_1^2$$

$$\Rightarrow \|A\|_2 = \sigma_1$$

2 Since  $A = U\Sigma V^H$

$$A^{-1} = (U\Sigma V^H)^{-1} = (V^H)^{-1} \Sigma^{-1} (U)^{-1} = V \Sigma^{-1} U^H$$

$\Rightarrow$  This is still an SVD of  $A^{-1}$

$$\text{And } \Sigma^{-1} = \begin{bmatrix} \frac{1}{\sigma_1} & & \\ & \ddots & \\ & & \frac{1}{\sigma_n} \end{bmatrix}$$

i.e. The singular values of  $A^{-1}$  are

$$\frac{1}{\sigma_1} \leq \dots \leq \frac{1}{\sigma_n}$$

By(1),  $\|A^{-1}\|_2$  is the largest singular value of  $A^{-1}$ , i.e.  $\frac{1}{\sigma_n}$

**3** Since  $A = U\Sigma V^H$

$$A^H = (V^H)^H \Sigma^H U^H = V \Sigma^H U = V \Sigma U$$

$\Rightarrow A^H$  and  $A$  has the same singular values.

$$\text{So } \|A^H\|_2^2 = \|A\|_2^2 = \sigma_1^2$$

$(A^T)$  is just  $A^H$  restricted on  $\mathbb{R}$ .

$$\text{So } \|A^T\|_2^2 = \|A^H\|_2^2$$

By the previous argument, we know that  $A^H A$  has an orthonormal eigenbasis  $\{v_1, v_2, \dots, v_n\}$ ,

$$\text{and } \forall \|x\|_2 = 1, \|A^H A x\|_2 = \|A^H A \sum c_i v_i\|_2 = \sqrt{(\sum c_i \sigma_i^2 v_i^T)(\sum c_i \sigma_i^2 v_i)} = \sqrt{\sum c_i \sigma_i^4} \leq \sigma_1^2$$

$$\text{Hence } \|A^H A\|_2 = \sup_{\|x\|=1} \|A^H A x\|_2 = \sigma_1^2$$

$$\text{It follows that } \|A^H A\|_2 = \|A\|_2^2 = \|A^H\|_2^2 = \|A^T\|_2^2 = \sigma_1^2$$

**4** Lemma: Let  $Q$  be an orthonormal matrix, then  $\|AQ\|_2 = \|A\|_2$ .

*Proof.* Let  $S_1 = \{ \|AQx\|, \|x\|_2 = 1 \}$ ,  $S_2 = \{ \|Ax\|, \|x\|_2 = 1 \}$

*Proof.* Since  $Q$  is orthonormal, so  $Q$  is also invertible.

$$\forall s_1 \in S_1, \exists x, \|x\|_2 = 1, \text{ s.t. } \|AQx\|_2 = s_1$$

$$\text{Now, let } y = Qx, \text{ it follows that } \|Qx\|_2 = \|y\|_2 = 1$$

$$\text{Since orthonormal matrix preserves length, } \|Ay\|_2 = \|AQx\|_2 = s_1 \in S_2$$

$$\text{i.e. } S_1 \subset S_2$$

$$\forall s_2 \in S_2, \exists x, \|x\|_2 = 1 \text{ s.t. } \|Ax\|_2 = s_2$$

$$\text{Now, let } y = Q^{-1}x, \text{ then } \|y\|_2 = \|Q^{-1}x\|_2 = 1$$

$$\text{Hence } \|AQy\|_2 = \|AQQ^{-1}x\|_2 = \|Ax\|_2 = s_2 \in S_1$$

$$\text{i.e. } S_2 \subset S_1$$

$$\therefore \|AQ\|_2 = \sup S_1 = \sup S_2 = \|A\|_2$$

□

Now  $\|UAV\|_2 = \|UA\|_2$  by lemma since  $V$  is an orthonormal matrix.

$$\|UA\|_2 = \sup_{\|x\|_2=1} \sqrt{(UAx)^H (UAx)} = \sup_{\|x\|_2=1} \sqrt{x^H A^H U^H U A x}$$

$$= \sup_{\|x\|_2=1} \sqrt{\langle Ax, Ax \rangle} = \|A\|_2$$

$$\text{Hence, } \|UAV\|_2 = \|UA\|_2 = \|A\|_2$$

□

## 32

**1** We need the following lemmas:

• lemma 1: if  $A, B \in M_n(\mathbb{F})$ , then  $\text{tr}(AB) = \text{tr}(BA)$

• lemma 2:  $\|A\|_F^2 = \text{tr}(A^T A)$

$$\text{Proof. let } A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{1m} & \dots & a_{mn} \end{bmatrix}$$

$$\text{Then } A^T = \begin{bmatrix} a_{11} & \dots & a_{m1} \\ \vdots & & \vdots \\ a_{1n} & \dots & a_{mn} \end{bmatrix}$$

$$\text{Observe that } (A^T A)_{nm} = a_{1n}^2 + a_{2n}^2 + \dots + a_{mn}^2 \\ \Rightarrow \text{tr}(A^T A) = \|A\|_p^2 \quad \square$$

$$\text{Now, } \|UAV\|_1^2 = \text{tr}((UAV)^T(UAV)) = \text{tr}(V^T A^T U^T U AV) = \text{tr}(V^T A^T AV) = \text{tr}(VV^T A^T A) = \text{tr}(A^T A) = \|A\|_1^2 \\ \Rightarrow \|UAV\|_2 = \|A\|_2$$

**2** Observe that  $A = U\Sigma V^T$ , with  $U$  and  $V^T$  orthonormal and

$$\Sigma = \begin{bmatrix} \sigma_1 & & & \\ & \ddots & & \\ & & \sigma_r & \\ & & & 0 \end{bmatrix} \quad \text{Now, } \|A\|_p^2 = \text{tr}(A^T A) = \text{tr}((U\Sigma V)^T(U\Sigma V)) = \text{tr}(U\Sigma^T U^T U\Sigma V^T) = \text{tr}(U\Sigma^2 V^T) \\ = \text{tr}(\Sigma^2) = \sum_{i=1}^r \sigma_i^2 \\ \text{Hence } \|A\|_p = \sqrt{\sum_{i=1}^r \sigma_i^2}$$

**33** Note that the  $Y^H Ax$  will be a field element. Consider it as a linear map  $Y^H Ax : \mathbb{F} \rightarrow \mathbb{F}$ , then the spectral norm of this map is:

$$\|Y^H Ax\|_2 = \sup_{f \in \mathbb{F}} \frac{\|(Y^H Ax)f\|_2}{\|f\|_2} = |Y^H Ax|$$

, where the first norm is spectral norm and the norm in fraction is the standard 2-norm.

**36** One example can be  $A = \begin{bmatrix} 4 & 0 \\ 3 & 5 \end{bmatrix}$

$$\text{then } A^T A = \begin{bmatrix} 25 & -15 \\ -15 & 25 \end{bmatrix}$$

$$\det(A^T A - \lambda I) = \lambda^2 - 50\lambda + 400 = 0$$

$$\lambda_1 = 40, \lambda_2 = 10$$

$$\text{Thus its singular value are } s_1 = \sqrt{40}, s_2 = \sqrt{10}$$

To calculate its eigenvalues,

$$\det(A - \lambda I) = \lambda^2 - 9\lambda + 20 = 0$$

Thus its eigenvalues are  $\lambda_1 = 4, \lambda_2 = 5$ , which are different from its singular values.

**38** (i) Suppose  $U\Sigma V^H$  is an SVD of  $A$ , then  $A^\dagger = V\Sigma^{-1}U^H$

$$AA^\dagger A = (U\Sigma V^H)(V\Sigma^{-1}U^H)(U\Sigma V^H) = U\Sigma V^H = A$$

(ii)

$$A^\dagger AA^\dagger = (V\Sigma^{-1}U^H)(U\Sigma V^H)(V\Sigma^{-1}U^H) = V\Sigma^{-1}U^H = A^\dagger$$

(iii)

$$(AA^\dagger)^H = ((U\Sigma V^H)(V\Sigma^{-1}U^H))^H = U\Sigma^{-1}V^H V\Sigma U^H = UU^H = AA^\dagger$$

(iv)

$$(A^\dagger A)^H = ((V\Sigma^{-1}U^H)(U\Sigma V^H))^H = V\Sigma U^H U\Sigma^{-1}V^H = VV^H = A^\dagger A$$



(v)

By prop (iii)  $\Rightarrow AA^\dagger$  is hermitian.

Also by prop (i),  $AA^\dagger AA^\dagger = AA^\dagger \Rightarrow AA^\dagger$  is idempotent.

Next we will check whether  $\mathcal{R}(AA^\dagger) = \mathcal{R}(A)$ .

It is trivially  $\mathcal{R}(AA^\dagger) \subset \mathcal{R}(A)$ , and by prop(i)  $\Rightarrow \mathcal{R}(A) \subset \mathcal{R}(AA^\dagger)$   
 $\Rightarrow \mathcal{R}(AA^\dagger) = \mathcal{R}(A)$

(vi)

By prop (iv)  $A^\dagger A$  is hermitian.

Also by prop (ii)  $A^\dagger AA^\dagger A = A^\dagger A \Rightarrow A^\dagger A$  is idempotent

Next we will check whether  $\mathcal{R}(A^\dagger A) = \mathcal{R}(A^H)$

By prop (iv),  $AA^\dagger = (AA^\dagger)^H = A^H(A^\dagger)^H \Rightarrow \mathcal{R}(A^\dagger A) \subset \mathcal{R}(A^H)$

Then we take the hermitian of both sides of prop (i),

we have  $(A^\dagger A)^H A^H = (A^\dagger A)^H A^H A^H \Rightarrow \mathcal{R}(A^H) \subset \mathcal{R}(A^\dagger A)$   
 $\Rightarrow \mathcal{R}(A^\dagger A) = \mathcal{R}(A^H)$