

A Gini coefficient
0·6
0·5
0·4
0·3
0·2
Gini coefficient
0·537
0·246 → 0·266
Gini of incidence rate
Gini of case counts
2004
2009
2014
2019
2024
Year
B Coefficient of variation
140
120
100
80
60
40
CV (%)
126·4%
no further convergence
47·6%
43·7%
2004
2009
2014
2019
2024
Year
C Between-province ratio
300×
200×
100×
50×
20×
10×
5×
Highest ÷ lowest
285×
7·1×
6·0×
2004
2009
2014
2019
2024
Year
D Total (2024) versus early syphilis (2020)
high total, high early: transmission-driven
high total, low early: detection-driven
Tibet
Hainan
Shanghai
Liaoning
Jiangsu
Xinjiang
Zhejiang
Guangdong
Guizhou
Hebei
Yunnan
Qinghai
15
12·5
10
7·5
5
2·5
0
2020 primary + secondary rate (per 100,000)
20
40
60
80
100
2024 notified incidence rate (per 100,000)
80%
60%
40%
20%
0%
-20%
E Rank shift between the two indicators
red: shift ≥15 places (n=5); nine provinces ≥12
Qinghai
Hunan
Yunnan
Liaoning
Jilin
1
5
10
15
20
25
30
Liaoning
Jilin
Hunan
Qinghai
Yunnan
2020 total incidence (31 provinces)
2020 primary + secondary (31 provinces)