

Foundations of Certified Programming Language and Compiler Design

Dr.-Ing. Sebastian Ertel

Composable Operating Systems Group, Barkhausen Institute



- **There is an internship (“Komplexpraktikum”) about LEAN at TUD!**
- Please have a look at [https://iccl.inf.tu-dresden.de/web/Theorem_Proving_with_LEAN_\(WS2023\)](https://iccl.inf.tu-dresden.de/web/Theorem_Proving_with_LEAN_(WS2023))
- Meet Stephan and Lukas and join their research!

Outline



Lecture	Logic Coq/Lean	Formalisms	PL Haskell
1	Propositional and first-order logic		
2			Functional programming
3		Syntax and Semantics	
4			The untyped lambda calculus
5		Types	
6			The typed lambda calculus
7			Polymorphism
8		Curry-Howard	
9			Higher-order types
10			Dependent types

Goals



Let's enter a more powerful logic system that allows us to express

- (in-)equalities and
- arithmetics.



Motivating examples

- Natural numbers with addition:

$$0 + 1 = 1$$

- Generalization:

$$0 + x = x$$

- And even tougher ones:

$$x + 0 = x, x + y = y + x$$

- Verification of the correctness of a compiler transformation `trans` for a program p :

$$\text{exec}(\text{trans } p) = \text{exec } p$$

From Propositional to First-Order Logic



<i>Syntax</i>	$A ::=$	formulas/propositions:
	X	propositional variables
	$A \Rightarrow A$	implication
	$A \wedge A$	conjunction
	\top	truth
	$A \vee A$	disjunction
	\perp	falsity
	$\neg A$	negation

Examples:

Propositional Logic

$$X_1 \vee X_2 \Rightarrow X_3$$



From Propositional to First-Order Logic

Syntax $A ::=$

- | $P(t_1, \dots, t_n)$
- | $A \Rightarrow A$
- | $A \wedge A$
- | \top
- | $A \vee A$
- | \perp
- | $\neg A$
- | $\forall x.A$
- | $\exists x.A$

formulas/propositions:

predicates on terms

implication

conjunction

truth

disjunction

falsity

negation

universally quantified term variables

existentially quantified term variables

Examples:

Propositional Logic

$$X_1 \vee X_2 \Rightarrow X_3$$

First-Order Logic

$$\forall x. \exists y. (x \times y = 1 \wedge y \times x = 1)$$



Syntax $P ::=$
| $P \in \mathcal{P}$ predicates:
 predicate symbol

- ... with $a :: \mathcal{P} \rightarrow \mathbb{N}$ defining the arity of predicates.
- Let $\mathcal{P} = \mathcal{X}$ where \mathcal{X} is the set of propositional variables such that $\forall X \in \mathcal{X}. a(X) = 0$,
- then

Propositional formula

$X \vee \neg Y$

First-order formula

$X() \vee \neg Y()$



Syntax $t ::=$

$\quad | \quad x$
 $\quad | \quad f(t_1, \dots, t_n)$
 $f ::= f \in \Sigma$

terms:

variables

function application

function symbols

- ... with an associated arity: $a :: \Sigma \rightarrow \mathbb{N}$
- and \mathcal{T} as the smallest set of terms.

Example:

Predicates

$\mathcal{P} = \{= : 2\}$

Terms

$\Sigma = \{\times : 2, 1 : 0\}$

Formula

$\forall x. \exists y. (x \times y = 1 \wedge y \times x = 1)$

Bound and Free Variables



Syntax $A ::= \dots$ formulas/propositions:

- | $\forall x.A$ universally quantified term variables
- | $\exists x.A$ existentially quantified term variables

- We say: “Term variable x is *bound* in formula A .”
- A variable that is not bound is *free*.
- A formula A is *closed* when it does not contain free variables, i.e., $FV(A) = \emptyset$.

Free Variables

Definitions and Notation

- ... of a formula A :

$$\begin{aligned}
 FV_A &:: A \rightarrow \{x | x \in X\} \\
 FV_A(P(t_1, \dots, t_n)) &= FV_t(t_1) \cup \dots \cup FV_t(t_n) \\
 FV_A(A_1 \Rightarrow A_2) = FV_A(A_1 \vee A_2) = FV_A(A_1 \wedge A_2) &= FV_A(A_1) \cup FV_A(A_2) \\
 FV_A(\top) = FV_A(\perp) &= \emptyset \\
 FV_A(\neg A_1) &= FV_A(A_1) \\
 FV_A(\forall x. A_1) = FV_A(\exists x. A_1) &= FV_A(A_1) \setminus \{x\}
 \end{aligned}$$

- ... of a term t :

$$\begin{aligned}
 FV_t &:: t \rightarrow \{x | x \in X\} \\
 FV_t(x) &= \{x\} \\
 FV_t(f(t_1, \dots, t_n)) &= FV_t(t_1) \cup \dots \cup FV_t(t_n)
 \end{aligned}$$

- ... of a context Γ :

$$\begin{aligned}
 FV_\Gamma &:: \Gamma \rightarrow \{x | x \in X\} \\
 FV_\Gamma(x : A_1, \dots, x : A_n) &= FV_A(A_1) \cup \dots \cup FV_A(A_n)
 \end{aligned}$$

- Notation: $A(x_1, \dots, x_n)$ where x_1, \dots, x_n are free variable of A



Definition (Substitution)

A *substitution* is a function $\sigma :: \mathcal{X} \rightarrow \mathcal{T}$ such that the set $\{x \in \mathcal{X} \mid \sigma(x) \neq x\}$ is finite.

- We write $t[\sigma(x)]$ for the term where every occurrence of x in t is replaced by $\sigma(x)$.
- The replacement function is trivial:

$$\begin{aligned}x[\sigma] &= \sigma(x) \\ f(t_1, \dots, t_n)[\sigma] &= f(t_1[\sigma], \dots, t_n[\sigma])\end{aligned}$$

- Notation:

$A(t_1, \dots, t_n)$ instead of $A[t_1/x_1, \dots, t_n/x_n]$



- We extend the natural deduction rules of propositional logic with the following rules:

$$\frac{\Gamma \vdash \forall x.A}{\Gamma \vdash A[t/x]} (\forall_E)$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash \forall x.A} (\forall_I) \text{ if } x \notin FV_\Gamma(\Gamma)$$

$$\frac{\Gamma \vdash \exists x.A_1 \quad \Gamma, A_1 \vdash A_2}{\Gamma \vdash A_2} (\exists_E) \text{ if } x \notin FV_\Gamma(\Gamma) \cup FV_A(A_2)$$

$$\frac{\Gamma \vdash A[t/x]}{\Gamma \vdash \exists x.A} (\exists_I)$$



Definition (First-order Theory)

A first-order theory $\Theta(\mathcal{P}, \Sigma)$ on a set of predicates \mathcal{P} and a signature of terms Σ is a (possibly infinite) set of closed formulas also called *axioms*.

Definition (Provability)

A formula A is *provable* if there is $\Gamma \subseteq \Theta$ such that $\Gamma \vdash A$ is provable.

- Properties:
 - consistent** when \perp is not provable in the theory
 - complete** when for every formula A , **either** A **or** $\neg A$ is provable in the theory
 - decidable** when there is an algorithm that decides whether a given formula is provable or not

The decidable theory of equality with uninterpreted functions



$\Theta := (\mathcal{P}, \Sigma)$ where $\mathcal{P} = \{=: 2\}$

$\Sigma = \{f : n \mid f \in \mathbf{Strings}, n \in \mathbb{N}\}$

- Axioms:

$$\boxed{t = t'}$$

$\frac{}{t = t}$ Reflexivity

$\frac{t_2 = t_1}{t_1 = t_2}$ Symmetry

$\frac{t_1 = t_{12} \quad t_{12} = t_2}{t_1 = t_2}$ Transitivity

$\frac{f = f' \quad t_1 = t'_1 \quad \dots \quad t_n = t'_n}{f(t_1, \dots, t_n) = f'(t'_1, \dots, t'_n)}$ Function Congruence

$\frac{P(A_1, \dots, A_m) \quad A_1 = A'_1 \quad \dots \quad A_m = A'_m}{P(A'_1, \dots, A'_m)}$ Predicate Congruence



- Axiomatizes the addition over natural numbers.
- Theory of equality over the signature $\Sigma = \{0 : 0, S : 1, + : 2\}$
- with the following axioms:

$$\frac{0 = S(x)}{\perp} \text{ (ax}_{=\perp}\text{)} \qquad \frac{S(x) = S(y)}{x = y} \text{ (S}_{\text{inj}}\text{)} \qquad \frac{}{0 + x = x} \text{ (ax}_{+1}\text{)} \qquad \frac{}{S(x) + y = S(x + y)} \text{ (ax}_{+2}\text{)}$$

Recurrence Principle



$$\frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \text{ (cases)}$$

$$\frac{x = 0 \vdash x + 0 = x \quad x = S(y) \vdash S(y) + 0 = S(y)}{x + 0 = x} \text{ (cases)}$$

Recurrence Principle



$\pi_S :=$

$\pi_0 :=$

$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \text{ (cases)}$

`fn pi_left_identity(x:Nat) { }`

Recurrence Principle



$$\pi_S :=$$

$$\pi_0 := 0 + 0 = 0 \quad =: (x + 0 = x)[0/x]$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \text{ (cases)}$$

Recurrence Principle



$$\pi_S :=$$

$$\pi_0 := \overline{0 + 0 = 0} \text{ (ax}_{+1}\text{)}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \text{ (cases)}$$



$$\pi_S := \frac{\frac{\forall x. S(x) + 0 = S(x+0) \quad \forall x. S(x+0) = S(x)}{\forall x. S(x) + 0 = S(x)} \quad (=_{\text{transitivity}})}{\quad} \quad (\forall_E)$$

$$\pi_0 := \overline{0 + 0 = 0} \quad (\text{ax}_{+1})$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \quad (\text{cases})$$



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x + 0)}}{\forall x.S(x) + 0 = S(x)} \text{ (ax}_{+2}\text{)} \quad \forall x.S(x + 0) = S(x)}{\frac{\forall x.S(x) + 0 = S(x)}{S(x) + 0 = S(x)} \text{ (}\forall_E\text{)}} \text{ (=transitivity)}$$

$$\pi_0 := \frac{}{0 + 0 = 0} \text{ (ax}_{+1}\text{)}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x + 0 = x} \text{ (cases)}$$



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x + 0)}}{\quad} (\text{ax}_{+2}) \quad \frac{\frac{S = S}{\quad} \quad \frac{\forall x.x + 0 = x}{\quad} (\text{fun cong})}{\forall x.S(x + 0) = S(x)} (\text{=transitivity})}{\frac{\forall x.S(x) + 0 = S(x)}{S(x) + 0 = S(x)}} (\forall_E)$$

$$\pi_0 := \overline{0 + 0 = 0} (\text{ax}_{+1})$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x + 0 = x} (\text{cases})$$

Recurrence Principle



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x + 0)}}{\quad} (\text{ax}_{+2}) \quad \frac{\frac{\overline{S = S}}{\quad} (\text{ax}_{S-\text{cong}}) \quad \forall x.x + 0 = x}{\forall x.S(x + 0) = S(x)} (\text{fun cong})}{\frac{\forall x.S(x) + 0 = S(x)}{S(x) + 0 = S(x)} (\forall_E)} (= \text{transitivity})$$

$$\pi_0 := \overline{0 + 0 = 0} (\text{ax}_{+1})$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x + 0 = x} (\text{cases})$$

Recurrence Principle



$$\begin{array}{c}
 \frac{\overline{\forall x.S(x) + 0 = S(x+0)} \text{ (ax}_{+2}\text{)}}{\overline{\forall x.S(x) + 0 = S(x)}} \text{ (}\forall_E\text{)} \\
 \frac{\overline{S = S} \text{ (ax}_{S\text{-cong}\text{)}} \quad \forall x.x + 0 = x \text{ (fun cong)}}{\forall x.S(x+0) = S(x)} \text{ (=transitivity)} \\
 \pi_S := \frac{\overline{\forall x.S(x) + 0 = S(x)}}{S(x) + 0 = S(x)}
 \end{array}$$

$$\pi_0 := \overline{0 + 0 = 0} \text{ (ax}_{+1}\text{)}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x + 0 = x} \text{ (cases)}$$

$$f(n) := \left\{ \begin{array}{l} \end{array} \right.$$

Recurrence Principle



$$\begin{array}{c}
 \frac{\overline{\forall x.S(x) + 0 = S(x+0)} \text{ (ax}_{+2}\text{)}}{\overline{\forall x.S(x) + 0 = S(x)}} \text{ (}\forall_E\text{)} \\
 \frac{\overline{S=S} \text{ (ax}_{S\text{-cong}}\text{)} \quad \forall x.x+0=x \text{ (fun cong)}}{\forall x.S(x+0)=S(x)} \text{ (=transitivity)} \\
 \pi_S := \frac{\overline{\forall x.S(x) + 0 = S(x)}}{S(x) + 0 = S(x)}
 \end{array}$$

$$\pi_0 := \overline{0+0=0} \text{ (ax}_{+1}\text{)}$$

$$f(n) := \begin{cases} \pi_0 & \text{if } n \text{ is } 0 \end{cases}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x+0=x} \text{ (cases)}$$



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x+0)}}{\quad} (\text{ax}_{+2}) \quad \frac{\frac{\overline{S=S}}{\quad} (\text{ax}_{S-\text{cong}}) \quad \frac{\overline{\forall x.f(x)}}{\forall x.x+0=x} (\text{fun cong})}{\forall x.S(x+0)=S(x)} (\text{fun cong})}{\frac{\forall x.S(x)+0=S(x)}{S(x)+0=S(x)} (\forall_E)} (=transitivity)$$

$$\pi_0 := \overline{0+0=0} (\text{ax}_{+1})$$

$$f(n) := \begin{cases} \pi_0 & \text{if } n \text{ is } 0 \\ \pi_S[y/x] & \text{if } n \text{ is } S(y) \end{cases}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x+0=x} (\text{cases})$$

Recurrence Principle



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x + 0)}}{(\text{ax}_{+2})} \quad \frac{\frac{\overline{S = S}}{(\text{ax}_S\text{-cong})} \quad \frac{\overline{\forall x.f(x)}}{\forall x.x + 0 = x} \quad (\text{fun cong})}{\forall x.S(x + 0) = S(x)} \quad (\text{=transitivity})}{\frac{\forall x.S(x) + 0 = S(x)}{S(x) + 0 = S(x)}} \quad (\forall_E)$$

π_S :=

$$\pi_0 := \overline{0 + 0 = 0} \quad (\text{ax}_{+1})$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x + 0 = x} \quad (\text{cases})$$

$$f(n) := \begin{cases} \pi_0 & \text{if } n \text{ is } 0 \\ \pi_S[y/x] & \text{if } n \text{ is } S(y) \end{cases}$$

```
fn f(x:Nat) {
  match x {
    0 => pi_0,
    S(y) => pi_S(y)
  }
}
```

Recurrence Principle



$$\pi_S := \frac{\frac{\overline{\forall x.S(x) + 0 = S(x+0)}}{\quad} (\text{ax}_{+2}) \quad \frac{\frac{\overline{S=S}}{\quad} (\text{ax}_{S-\text{cong}}) \quad \frac{\forall x.f(x)}{\forall x.x+0=x}}{\forall x.S(x+0)=S(x)} (\text{fun cong})}{\frac{\forall x.S(x)+0=S(x)}{S(x)+0=S(x)}} (= \text{transitivity}) (\forall_E)$$

$$\pi_0 := \overline{0+0=0} (\text{ax}_{+1})$$

$$f(n) := \begin{cases} \pi_0 & \text{if } n \text{ is } 0 \\ \pi_S[y/x] & \text{if } n \text{ is } S(y) \end{cases}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x.x+0=x} (\text{cases})$$

axiom scheme:

$\frac{A(0) \quad (\forall x.A(x) \rightarrow A(S(x)))}{\forall x.A x} (\text{Rec})$

Induction Principle



$$\pi_S := \frac{\frac{\overline{S(x) + 0 = S(x + 0)}}{S(x) + 0 = S(x + 0)} \text{ (ax}_{+2}\text{)} \quad \frac{\frac{\overline{S = S} \text{ (ax}_{S\text{-cong}}) \quad x + 0 = x \vdash x + 0 = x}{x + 0 = x \vdash S(x + 0) = S(x)} \text{ (fun cong)}}{x + 0 = x \vdash S(x) + 0 = S(x)} \text{ (=transitivity)}}{x + 0 = x \Rightarrow S(x) + 0 = S(x)} \text{ (}\Rightarrow_I\text{)}$$

$$\pi_0 := \overline{0 + 0 = 0} \text{ (ax}_{+0}\text{)}$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \text{ (Ind)}$$

$\frac{A(0) \quad (\forall x. A(x) \Rightarrow A(S(x)))}{\forall x. A(x)} \text{ (Ind)}$

Induction Principle



$$\pi_S := \frac{\frac{\frac{}{S(x) + 0 = S(x+0)}{(\text{ax}_{+2})} \quad \frac{\frac{}{S=S}(\text{ax}_{S-\text{cong}}) \quad \frac{}{x+0=x \vdash x+0=x}(\text{ax})}{x+0=x \vdash S(x+0)=S(x)}(\text{fun cong})}{x+0=x \vdash S(x)+0=S(x)}(=\text{transitivity})}{x+0=x \Rightarrow S(x)+0=S(x)}(\Rightarrow_I)$$

$$\pi_0 := \frac{}{0+0=0} (\text{ax}_{+0})$$

$$\pi_{\text{Left Identity}} := \frac{\pi_0 \quad \pi_S}{\forall x. x+0=x} (\text{Ind})$$

$\frac{A(0) \quad (\forall x. A(x) \Rightarrow A(S(x)))}{\forall x. A(x)} (\text{Ind})$

Induction Principle



Induction Hypothesis

$$\pi_S := \frac{\frac{\overline{S(x) + 0 = S(x+0)}}{(ax_{+2})} \quad \frac{\frac{\overline{S = S}}{(ax_{S-cong})} \quad \frac{\overline{x + 0 = x \vdash x + 0 = x}}{(ax)}}{x + 0 = x \vdash S(x+0) = S(x)} \quad (fun\ cong)}{\frac{x + 0 = x \vdash S(x) + 0 = S(x)}{x + 0 = x \Rightarrow S(x) + 0 = S(x)}} \quad (\Rightarrow_I)$$

$$\pi_0 := \overline{0 + 0 = 0} \quad (ax_{+0})$$

$$\pi_{Left\ Identity} := \frac{\pi_0 \quad \pi_S}{\forall x. x + 0 = x} \quad (Ind)$$

$\frac{A(0) \quad (\forall x. A(x) \Rightarrow A(S(x)))}{\forall x. A\ x} \quad (Ind)$



$$\frac{x = 0 \vdash A(x) \quad (\forall x. A(x) \Rightarrow A(S(x)))}{\forall x. A(x)} \text{ (Ind)} \quad \frac{x = 0 \vdash A(x) \quad x = S(y) \vdash A(S(y))}{\forall x. A(x)} \text{ (cases)}$$

- Important point to remember:
 - Cases does not provide a **induction hypothesis**, i.e.,
 - it does not cover recursion!



$$\frac{A(0) \quad (\forall x. A(x) \Rightarrow A(S(x)))}{\forall x. A x} \text{ (Ind)}$$

$$\frac{A(0) \quad (\forall x. A(x) \rightarrow A(S(x)))}{\forall x. A x} \text{ (Rec)}$$

- Look what just happened:
 - We used a proof just like a function!
 - There seems to be a deeper connection between implication and a function (type)?
- Let's explore this connection in this lecture!