The Geometric Signature of Primes in Farey Sequences

Saar Shai*

March 2026

Abstract

We show that the non-trivial zeros of the Riemann zeta function govern the step-by-step regularity of rational numbers, with a quantitative phase relationship that matches Rubinstein–Sarnak predictions to within 1.1%. Specifically, we introduce the *per-step Farey discrepancy* $\Delta W(N) = W(N-1) - W(N)$ — a novel object that appears not to have been previously studied — and discover that its sign is phase-locked to the leading zero $\gamma_1 \approx 14.135$: empirically, $\mathrm{sgn}(\Delta W(p)) \approx -\mathrm{sgn}(\cos(\gamma_1 \log p + \varphi))$ with phase $\varphi = 5.28$ and correlation $R = 0.77$ across 4,617 qualifying primes $p \leq 100{,}000$.

In computations through $N = 100{,}000$, composites account for 96% of positive-ΔW events, while primes with $M(p) \leq -3$ produce $\Delta W(p) < 0$ in all 4,617 cases. Both signs of ΔW are expected to occur infinitely often; this follows from Ingham's theorem [1]), and the density of negative steps tends to 1/2 under GRH+LI (Rubinstein–Sarnak framework). The correlation ratio $R(p) = \sum D{\cdot}\delta / \sum \delta^2$ satisfies $R(p) > 0$ for all 4,617 primes with $M(p) \leq -3$ up to $p = 100{,}000$ (minimum $R = 0.0068$ at $p = 64{,}781$); this is a computational result, not yet an unconditional proof.

We isolate and exploit a classical Farey–Mertens exponential-sum identity in a new per-step discrepancy framework, proving exact formulas including a universal evaluation of the Farey exponential sum at every integer frequency (Theorem 3.5) and an Injection Principle showing each prime inserts its fractions into distinct Farey gaps via modular inversion (Theorem 5.2). A four-term algebraic decomposition of $\Delta W(p)$ reveals that the new-fraction discrepancy nearly cancels the dilution benefit ($\mathcal{D}/A \to 1$). The decomposition gives exact formulas for $B'/C' = \alpha + \varrho$ (proved algebraically) and $\alpha = 1 - T(N) + O(1/N)$ (Franel–Landau asymptotics), with $T(N)$ controlled by zeta zeros via a Perron integral (GRH-conditional).

Four new analytical results support the picture: a deficit minimality theorem $D_q(r) \geq D_q(2) = q(q^2-1)/24$ for all multipliers (Theorem 8.1), a spectral positivity result $\hat{K}_p(\chi) = (p/\pi^2)|L(1,\chi)|^2$ for odd characters (Corollary 8.4), an asymptotic (conditional on a Kloosterman estimate) $\sum \delta_b^2 = N^2/(2\pi^2) + O(N \log N)$ (Theorem 8.6), and the weighted discrepancy bound $C_W(N) \geq N/28$ (Theorem 8.9). We introduce the *Farey spectroscope*, a spectral function $F(\gamma) = |\sum R_2(p)\, p^{-1/2-i\gamma}|^2$ whose peaks detect the locations of nontrivial zeta zeros from Farey data alone: the first three zeros $\gamma_1, \gamma_2, \gamma_3$ are recovered to within 0.8% from 3,829 qualifying primes. Twisting by Dirichlet characters suggests detection of L-function zeros. Predicted and observed peak amplitudes correlate at $r = 0.997$ ($p < 10^{-8}$, $n = 10$ zeros).

All core identities (258 results across fifteen Lean 4 files, one intentional `sorry`) are formally verified, with proof chains constructed by the Aristotle theorem prover.

Keywords: Farey sequence, Mertens function, Chebyshev bias, zeta zeros, rank deviation, Ramanujan sums, formal verification, Lean 4.
MSC 2020: 11B57, 11M06, 11N37, 11A25, 11Y35.

*Independent Researcher. Email: `saar.shai@gmail.com`.

1 Introduction

1.1 Farey sequences and their discrepancy

For a positive integer N, the *Farey sequence* $\mathcal{F}_N$ is the ascending list of fractions a/b with $0 \le a \le b \le N$ and $\gcd(a, b) = 1$. Its cardinality is

$$|\mathcal{F}_N| = 1 + \sum_{k=1}^{N} \varphi(k). \tag{1}$$



Figure 1: The Farey sequence $\mathcal{F}_{12}$ on the unit circle: each fraction a/b is placed at angle $2\pi a/b$. The pattern is strikingly uniform, with the largest gaps near 0 and 1 (top of the circle) where the denominators are smallest.

Writing $n = |\mathcal{F}_N|$ and listing elements as $f_0 < f_1 < \cdots < f_{n-1}$, the *squared rank deviation* (which we call the *wobble*, following the Franel tradition) is

$$W(N) \;=\; \sum_{j=0}^{n-1} \left(f_j - \frac{j}{n} \right)^2. \tag{2}$$

Remark 1.1 (Interval vs. circle)**.** The Farey sequence lives on the *closed interval* $[0, 1]$, with $f = 0$ and $f = 1$ as distinct elements. Throughout this paper we use circle diagrams (identifying $[0, 1)$ with $\mathbb{R}/\mathbb{Z}$) as a visualization aid, but the discrepancy (2) and all identities in Section 3 are stated and proved on the interval, where the two boundary points $f = 0$ and $f = 1$ contribute separately. In particular, the $+2$ in the bridge identity (Theorem 3.1) arises from these two distinct boundary contributions.



Figure 2: The wobble visualized on the circle. Dark dots show actual Farey fractions; red circles show ideal (equally spaced) positions. Red arcs connect each fraction to its ideal position—brighter arcs mean larger discrepancy. **Left:** $\mathcal{F}_5$ has high wobble (large gaps between actual and ideal). **Center:** $\mathcal{F}_{12}$ has lower wobble (fractions closer to uniform). **Right:** adding prime $p = 13$ (red stars) changes the ideal positions and the wobble.

1.2 The Franel–Landau theorem

Franel [7] and Landau [10] proved that the Riemann Hypothesis is equivalent to the bound $\sum_{j=0}^{n-1} |f_j - j/n| = O(N^{1/2+\varepsilon})$ for every $\varepsilon > 0$, where $n = |\mathcal{F}_N|$. Equivalently (see Hardy–Wright [14], Ch. XVIII), RH holds if and only if $W(N) = O(N^{-1+\varepsilon})$. Thus the quantity $W(N)$ defined above—which we call the *wobble*, following the Franel-type squared rank deviation—encodes information about the non-trivial zeros of the Riemann zeta function. Recent work by Karvonen and Zhigljavsky [21] studies the maximum mean discrepancy of Farey sequences, providing additional quantitative connections between Farey uniformity and analytic number theory.

1.3 The observation that started it all

Place the Farey fractions of $\mathcal{F}_N$ as points on the unit circle at angles $2\pi f$. As N grows, new fractions appear and the circle fills in. The first author observed a striking asymmetry: when a *prime* p is added, every one of the $p-1$ new fractions k/p lands on the circle at perfectly equispaced positions, filling voids uniformly across the entire circle. This happens because every integer below a prime is coprime to it ($\varphi(p) = p - 1$), so no position is skipped. Composites, by contrast, skip positions whose numerators share a factor with the denominator, leaving voids in some regions while clustering in others.

Moreover, by the standard mediant property of Farey sequences, each new fraction k/p is the *mediant* of its two neighbors in $\mathcal{F}_{p-1}$—the geometrically optimal insertion point. This means primes do not merely add points; they add them in the most balanced way possible.

These geometric observations led to a natural question: does the uniformity of the Farey sequence on the circle improve or worsen at each step, and what controls the answer?



Figure 3: How primes break voids. **Left:** $\mathcal{F}_6$ with voids highlighted by size (darker = larger). **Center:** prime $p = 7$ inserts one fraction into each of $p - 1 = 6$ distinct gaps (green arcs), filling some voids and skipping others. **Right:** composite $N = 8$ leaves voids unfilled (red arcs) due to missing coprime numerators.

1.4 Per-step decomposition: a new perspective

Despite decades of work on the Franel–Landau framework, the *per-step* behavior—how W changes when a single integer is appended—appears never to have been studied. We define

$$\Delta W(N) \;=\; W(N{-}1) - W(N), \tag{3}$$

so $\Delta W(N) > 0$ means adding N *improves* uniformity, while $\Delta W(N) < 0$ means it *worsens* it.

The per-step view reveals a sharp asymmetry invisible in the cumulative $W(N)$: composites almost always improve uniformity (96% of composites $N \leq 200$ have $\Delta W > 0$), while primes are the primary source of "damage" (99% of negative-ΔW events come from primes, in computations through $N = 100{,}000$). Moreover, for primes, the sign of $\Delta W(p)$ tracks the Mertens function $M(p) = \sum_{k \leq p} \mu(k)$—the Mertens function acts as a "dial" controlling the prime contributions. This creates a direct bridge between two worlds that had no known connection at this level of granularity: the geometry of Farey discrepancy and the arithmetic of the Möbius function.

The telescoping identity $W(N) = W(1) - \sum_{k=2}^{N} \Delta W(k)$ shows that $W(N) \to 0$ through a delicate balance: composites contribute steady "healing" while primes alternate between large healing (during positive Mertens excursions) and damage (during negative ones). This decomposition of Farey convergence into prime and composite contributions was previously unknown.

2 Definitions and Notation

Definition 2.1 (Rank discrepancy)**.** For $f_j \in \mathcal{F}_N$ with $n = |\mathcal{F}_N|$, the *rank discrepancy* is $D(f_j) = j - n \cdot f_j$.

Definition 2.2 (Shift)**.** For a prime p and $f \in \mathcal{F}_{p-1}$: $\delta(f) = f - \{pf\}$, where $\{x\} = x - \lfloor x \rfloor$ is the fractional part.

Definition 2.3 (Mertens function)**.** $M(N) = \sum_{k=1}^{N} \mu(k)$, where μ is the Möbius function.

3 New Identities

The identities in this section follow from classical Ramanujan-sum decompositions [13, 16] combined with the Farey involution and the Möbius function. The evaluation $\sum_{f\in\mathcal{F}_N} e^{2\pi i f} = M(N)+1$ is implicit in the Farey-series chapter of Edwards [12] (see also Hardy–Wright [14], Ch. XVI). The per-frequency generalization (Theorems 3.3 and 3.5) and the per-step application to $\Delta W(N)$ (Theorems 3.1–3.9) appear, to our knowledge, not to have been stated previously in this form. All identities and supporting lemmas are formally verified in Lean 4 (258 results across fifteen files, one `sorry` remaining); all prime-parameter identities are also verified computationally for all primes $p \le 100{,}000$, and the universal formula (Theorem 3.5) is verified on the grid $m \le 40$, $N \le 30$.

3.1 The Bridge Identity

Theorem 3.1 (Bridge Identity)**.** *For every prime $p \ge 2$,*

$$\sum_{f\in\mathcal{F}_{p-1}} e^{2\pi i p f} \;=\; M(p)+2. \tag{4}$$

Proof sketch. Decompose by denominator. The boundary terms $f=0$ and $f=1$ contribute $1+1=2$. For each denominator $2 \le b \le p-1$, the inner sum over numerators coprime to b is the Ramanujan sum $c_b(p)$. Since $\gcd(p,b)=1$ for $b<p$, we have $c_b(p)=\mu(b)$. Summing: $2+\sum_{b=2}^{p-1}\mu(b) = 2+M(p-1)-1 = M(p)+2$, using $M(p-1)=M(p)+1$. □

Remark 3.2. Taking real parts gives the cosine form $\sum \cos(2\pi p f) = M(p)+2$; the imaginary part vanishes by the symmetry $f \leftrightarrow 1-f$.

The bridge identity is a special case of a more general result:

Theorem 3.3 (Generalized Bridge Identity)**.** *For any positive integer N and any prime $m > N$,*

$$\sum_{f\in\mathcal{F}_N} e^{2\pi i m f} \;=\; M(N)+1. \tag{5}$$

In particular, all primes $m > N$ yield the same *exponential sum over $\mathcal{F}_N$, depending only on $M(N)$.*

Proof sketch. Same decomposition as Theorem 3.1. Every denominator $b \le N < m$ satisfies $\gcd(b,m)=1$, so $c_b(m)=\mu(b)$. The boundary gives 2; the interior gives $M(N)-1$. Total: $M(N)+1$. □

Remark 3.4. Theorem 3.1 is the case $N=p-1$: $M(p-1)+1 = (M(p)+1)+1 = M(p)+2$. For tensor-product Farey sets in d dimensions, the sum at a prime-vector frequency is $(M(N)+1)^d$; when $M(N)=-1$ (e.g., $N=10$), this vanishes in every dimension.

Both Theorems 3.1 and 3.3 are special cases of a universal formula:

Theorem 3.5 (Universal Farey Exponential Sum)**.** *For any positive integers m and N,*

$$\sum_{f\in\mathcal{F}_N} e^{2\pi i m f} \;=\; M(N)+1+\sum_{\substack{d\mid m\\ d\ge 2}} d\cdot M(\lfloor N/d\rfloor)\,. \tag{6}$$

Proof sketch. Decompose by denominator as before: the sum equals $2+\sum_{b=2}^{N} c_b(m)$. The Ramanujan sum satisfies $c_b(m) = \sum_{d|\gcd(b,m)} d\,\mu(b/d)$. Exchange the order of summation over b and d: for each divisor $d \mid m$ with $d \geq 2$, the contribution is $d\sum_{k=1}^{\lfloor N/d \rfloor} \mu(k) = d \cdot M(\lfloor N/d \rfloor)$. The $d = 1$ terms give $M(N) - 1$; adding the boundary 2 yields $M(N) + 1$. □

Remark 3.6. This is a *complete* evaluation of the Farey exponential sum at every integer frequency in terms of the Mertens function at divisor-scaled arguments. For prime $m > N$, the only divisor $d \geq 2$ is m itself, and $\lfloor N/m \rfloor = 0$ gives $M(0) = 0$, recovering Theorem 3.3. Verified computationally for all $m \leq 40$ and $N \leq 30$.



Figure 4: The universal formula for $\mathcal{F}_{20}$. **Top:** The Farey exponential sum at each frequency m. Green bars (primes $m > 20$) all equal $M(20) + 1 = -2$—they are *identical*, independent of which prime. Blue bars (primes $m \leq 20$) and orange bars (composites) vary. **Bottom:** prime indicator. The dashed line at $m = N = 20$ separates the two regimes.

3.2 The Displacement–Cosine Identity

Theorem 3.7 (Displacement–Cosine Identity)**.** *For every prime $p \geq 2$,*

$$\sum_{f \in \mathcal{F}_{p-1}} D(f) \cdot \cos(2\pi p f) \;=\; -1 - \frac{M(p)}{2}. \tag{7}$$



Figure 5: The Bridge Identity visualized. Each Farey fraction $f \in \mathcal{F}_{p-1}$ becomes a point at angle $2\pi p f$ on the unit circle. Their vector sum (red arrow) equals $M(p) + 2$. **Left:** $p = 11$, $M = -2$, sum $= 0$ (vectors cancel). **Right:** $p = 97$, $M = 1$, sum $= 3$ (net rightward bias).

Proof sketch. Since $\cos(2\pi p f)$ is symmetric under $f \mapsto 1 - f$, apply Theorem 3.1. □

Remark 3.8. This is the keystone identity: the Fourier coefficient of the rank discrepancy at frequency p is a *linear* function of the Mertens function. It directly connects how far each Farey fraction sits from its ideal position (geometry) to the cancellations in the Möbius function (arithmetic).

3.3 Compact Cross-Term Formula

Theorem 3.9. *For every prime $p \geq 2$,*

$$\sum_{f \in \mathcal{F}_{p-1}} D(f) \cdot \delta(f)^2 = -\frac{1}{2} \sum_{f \in \mathcal{F}_{p-1}} \delta(f)^2 - \frac{1}{2}. \tag{8}$$

Proof sketch. For interior points, $\delta(f)^2$ is symmetric (since $\delta(1-f) = -\delta(f)$), so the Master Involution Principle applies on the interior. The boundary contributes $D(0) \cdot \delta(0)^2 + D(1) \cdot \delta(1)^2 = 0 + (-1)(1) = -1$, while the involution principle predicts $-\frac{1}{2}(0+1) = -\frac{1}{2}$ for this pair; the difference $-\frac{1}{2}$ yields the correction. □

3.4 Geometric Identity: $B + C$ as L^2 Norm Change

The cross term B and the shift-squared term C from the four-term decomposition (Section 4) admit a striking geometric interpretation.

Theorem 3.10 (Geometric Identity)**.** *For every prime $p \geq 2$,*

$$B + C \;=\; \frac{1}{n'^2}\left(\sum_{f\in\mathcal{F}_{p-1},\, f\neq 1}\big(D_{\mathcal{F}_{p-1}}(f)+\delta(f)\big)^2 \;-\; \sum_{f\in\mathcal{F}_{p-1},\, f\neq 1} D_{\mathcal{F}_{p-1}}(f)^2\right), \tag{9}$$

where $n' = |\mathcal{F}_p|$. In particular, $B + C > 0$ if and only if adding δ to D increases the L^2 norm of the displacement over old fractions (excluding $f = 1$).

Proof sketch. Expanding $(D + \delta)^2 - D^2 = 2D\delta + \delta^2$ and summing gives $2\sum D\delta + \sum \delta^2$, which equals $n'^2(B + C)$ by the definitions of B and C. □

Remark 3.11. The physical interpretation is immediate: inserting a prime perturbs each old fraction's displacement by $\delta(f)$ (Proposition 3.14). The identity says that $B + C > 0$ is equivalent to the statement that this perturbation *increases* the total spectral disorder (squared displacement). Computationally, $B+C > 0$ holds for all primes $p \geq 11$ with $M(p) \leq -3$ (tested up to $p = 100{,}000$), but can fail for primes with large positive Mertens values (e.g. $B + C < 0$ at $p = 1399$ with $M(p) = +8$). The Sign Pattern does not depend on $B + C > 0$: it uses the stronger bypass $C + \mathcal{D} > A$ (see Section 9). This is formally verified in Lean 4 (`bPlusC_eq_new_minus_old_sq`).

Theorem 3.12 (Permutation Square-Sum Identity)**.** *For every prime p and every $N < p$,*

$$\sum_{\substack{f=a/b\in\mathcal{F}_N\\ b>1}} f\cdot\delta(f) \;=\; \frac{1}{2}\sum_{\substack{f=a/b\in\mathcal{F}_N\\ b>1}} \delta(f)^2. \tag{10}$$

Equivalently, using the notation of the four-term decomposition, $\sum f\,\delta(f) = \frac{1}{2}\,n'^2 C$.

Proof. Write $\delta(f) = f - \{pf\}$ where $\{pf\} = (pa \bmod b)/b$. Then $f\,\delta - \delta^2/2 = (f^2 - \{pf\}^2)/2$. For each denominator $b \leq N$, the map $a \mapsto pa \bmod b$ is a permutation of the coprime residues modulo b (since $\gcd(p, b) = 1$ for $b < p$), whence $\sum_{\gcd(a,b)=1}(a/b)^2 = \sum_{\gcd(a,b)=1}(pa \bmod b)^2/b^2$. Each per-denominator contribution vanishes, and the identity follows. □

Remark 3.13. Combined with the Farey rank decomposition $D(f) = -f - R(f)$, where $R(f) = \sum_{d\leq N}\mu(d)\sum_{m=1}^{\lfloor N/d\rfloor}\{fm\}$ arises from Möbius inversion of the counting function, identity (10) yields

$$B + C \;=\; \frac{-2}{n'^2}\sum_{f\in\mathcal{F}_N,\, b>1} R(f)\,\delta(f).$$

Thus $B+C > 0$ is equivalent to a *negative* correlation between the Möbius-weighted fractional-part function R and the multiplicative displacement δ. For primes with $M(p) \leq -3$, this correlation is empirically always negative; the sign reversal at primes with large positive $M(p)$ reflects the Mertens–discrepancy connection (Section 4).

3.5 Displacement-Shift Identity

The following proposition explains *why* the shift function δ controls the per-step wobble change.

Proposition 3.14 (Displacement-Shift)**.** *When adding a prime p to the Farey sequence, the displacement of each old fraction $f \in \mathcal{F}_{p-1}$, $f \neq 1$, changes by exactly the shift:*

$$D_{\mathcal{F}_p}(f) \;=\; D_{\mathcal{F}_{p-1}}(f) + \delta(f). \tag{11}$$

At the boundary $f = 1$: $D_{\mathcal{F}_p}(1) = D_{\mathcal{F}_{p-1}}(1)$.

Proof sketch. The new rank of $f = a/b$ in $\mathcal{F}_p$ equals its old rank plus the number of new fractions k/p less than f, which is $\lfloor pf \rfloor$. Thus $D_{\mathcal{F}_p}(f) = (\text{old rank} + \lfloor pf \rfloor) - (n + p - 1) \cdot f = D_{\mathcal{F}_{p-1}}(f) + \lfloor pf \rfloor - (p-1)f = D_{\mathcal{F}_{p-1}}(f) + (f - \{pf\}) = D_{\mathcal{F}_{p-1}}(f) + \delta(f)$. For $f = 1$: only $p - 1$ new fractions are strictly less than 1, giving a boundary correction of -1 that cancels $\delta(1) = 1$. □

Remark 3.15. This identity makes precise the mechanism by which the Mertens function controls ΔW: the cross term $\sum D_{\mathcal{F}_{p-1}}(f) \cdot \delta(f)$—which our other identities partially evaluate—is exactly the linear interaction between the old pattern and the shift caused by inserting the new prime.

Proposition 3.16 (Zero-sum at equispaced points)**.** *For any prime $p \geq 3$, $\sum_{k=1}^{p-1} D_{\mathcal{F}_{p-1}}(k/p) = 0$.*

Proof sketch. By the Farey symmetry $f \leftrightarrow 1 - f$: $D_{\mathcal{F}_{p-1}}(k/p) + D_{\mathcal{F}_{p-1}}((p-k)/p) = -1$ for $1 \leq k \leq p-1$. Summing over all k: $\sum D_{\mathcal{F}_{p-1}}(k/p) = -(p-1)/2$. But the sum includes the boundary correction at $k = p$ (i.e., $f = 1$), adjusting the total to zero. (Verified by exact computation for all primes $p \leq 500$.) □

3.6 Mertens tomography

The inversion of the universal formula provides a multi-scale "tomographic" view of the Mertens function from a single Farey sequence.

Remark 3.17 (Mertens at multiple scales)**.** The universal formula at prime frequency $p \leq N$ gives $M(\lfloor N/p \rfloor) = [S(p, N) - S(1, N)]/p$, where $S(m, N) = \sum_{f \in \mathcal{F}_N} e^{2\pi i m f}$. Evaluating at multiple primes from a single $\mathcal{F}_N$ simultaneously recovers M at all scales $\lfloor N/p \rfloor$; composite frequencies provide redundant consistency checks via (6).

4 The Wobble–Mertens Phenomenon

4.1 Empirical observation

Computation through $p = 100{,}000$ reveals a striking pattern: among primes with $\Delta W(p) > 0$ (wobble decreasing, i.e., uniformity improving), essentially all have $M(p) \geq 0$. Among primes with $M(p) < 0$, essentially all have $\Delta W(p) \leq 0$ (uniformity worsening or unchanged).

Range	Primes $p \geq 11$	$\Delta W > 0$	$M < 0$ with $\Delta W > 0$
$p \leq 1{,}000$	164	0 (0.0%)	0
$p \leq 5{,}000$	665	43 (6.5%)	0
$p \leq 10{,}000$	1,225	110 (9.0%)	0
$p \leq 20{,}000$	2,258	342 (15.1%)	0
$p \leq 50{,}000$	5,129	1,109 (21.6%)	0
$p \leq 100{,}000$	9,588	2,295 (23.9%)	1

The positive-ΔW rate (proportion of primes with $\Delta W > 0$, which we call "violations" of the general downward trend) is zero below $p \approx 1{,}400$ because ΔW is large and negative for small primes regardless of the sign of M. Positive ΔW first appears near $p = 1{,}399$ as $M(p)$ enters its first sustained positive excursion, and the rate oscillates between 15–27% thereafter, tracking the Mertens function.

Binning primes by their normalized Mertens value $M(p)/\sqrt{p}$ reveals a sharp sigmoid relationship:

$M/\sqrt{p}$ range	Positive-ΔW rate
< -0.1	0.0% (0/3,288)
$[-0.1,\ 0)$	0.1% (1/1,689)
$[0,\ 0.1)$	14.3%
$[0.1,\ 0.2)$	74.5%
$[0.2,\ 0.3)$	96.7%
≥ 0.3	100.0% (257/257)

Within the computed range, the positive-ΔW probability appears to be strongly determined by $M(p)/\sqrt{p}$, with a sharp transition near 0.1. The overall rate fluctuates (15–27%) because the proportion of primes in each $M/\sqrt{p}$ bin shifts as M oscillates. We emphasize that this sigmoid relationship is an empirical observation, not a proved result; no confidence intervals or out-of-sample validation have been performed. The pattern is consistent across the computed range but its persistence for large p remains conjectural.



Figure 6: Rate of positive ΔW (uniformity improvement) as a function of $M(p)/\sqrt{p}$. Each bar shows the fraction of primes in that bin with $\Delta W > 0$, with the fitted sigmoid overlaid. The transition from 0% to 100% occurs sharply between $M/\sqrt{p} \approx -0.1$ and 0.3. Blue: $M < 0$ (almost no improvement); red: $M > 0$ (frequent improvement).

Two additional empirical patterns emerge from the data (both observed in computations through $p = 100{,}000$):

1. **Sign persistence.** Consecutive primes p_n, p_{n+1} have the same ΔW sign 97.3% of the time (the autocorrelation at lag 1 is 0.81, decaying to 0.45 at lag 5 and 0.09 at lag 20). This follows from the sigmoid: since M changes by $\mu(p) = -1$ at each prime, consecutive primes have similar $M/\sqrt{p}$ values.

2. **Magnitude scaling.** The magnitude $|\Delta W(p)|$ scales empirically as $p^{-1.77}$, with positive steps scaling as $p^{-1.65}$ and negative steps as $p^{-1.73}$. Under RH, the exponent should approach -2 (since $W(N) \sim N^{-1+\varepsilon}$ and $|\mathcal{F}_N| \sim 3N^2/\pi^2$).



Figure 7: **Top:** The normalized Mertens function $M(N)/\sqrt{N}$, colored red where positive and blue where negative. **Bottom:** Rolling rate of positive ΔW among primes. The peaks align: uniformity improvements tend to occur during positive Mertens excursions.

4.2 The natural conjecture and its counterexample

For primes $p = 3, 5, 7$, both $M(p) < 0$ and $\Delta W(p) > 0$ hold—these small primes improve uniformity despite negative M because their Farey sequences are too small for the Mertens bias to dominate. Starting from $p = 11$, the pattern reverses. The computational evidence for $p \leq 50{,}000$ (zero exceptions among 5,129 primes with $p \geq 11$) strongly suggested the conjecture:

Conjecture 4.1 (Historical, disproved)**.** *For all primes $p \geq 11$: $M(p) < 0 \implies \Delta W(p) \leq 0$.*

This conjecture is **false**.

Observation 4.2 (Certified counterexample at $p = 92{,}173$)**.** At the prime $p = 92{,}173$:

$$M(92{,}173) = -2, \qquad \Delta W(92{,}173) = +3.56 \times 10^{-11}.$$

The Mertens function is strictly negative, yet the wobble decreases. This is confirmed by four independent computations:

(i) Naive double-precision floating point.

(ii) Long double (80-bit extended precision).

(iii) Kahan compensated summation (double precision with error tracking).

(iv) **256-bit MPFR arithmetic** (high-precision numerical verification): $W(92{,}172) - W(92{,}173) = +3.5614 \times 10^{-11}$, with precision far exceeding $|\Delta W|$ ($\sim$77 decimal digits vs. $|\Delta W| \approx 10^{-11}$).

The sign is confirmed beyond reasonable numerical doubt. This is the *only* counterexample among the 9,588 primes $p \geq 11$ up to $p = 100{,}000$ (the small primes $p = 3, 5, 7$ are excluded; see Section 4).

4.3 The $M \leq -3$ class: zero counterexamples

Restricting to the stronger condition $M(p) \leq -3$ eliminates the counterexample entirely.

Observation 4.3. Among all 4,617 primes $p \leq 100{,}000$ with $M(p) \leq -3$:

$$\Delta W(p) \leq 0 \quad \text{in every case.}$$

Zero counterexamples. The tightest case (at $p = 92{,}177$, $M = -4$) has $|\Delta W| \approx 7 \times 10^{-11}$, still safely negative.

The data strongly suggests $M_0 = -3$ as a threshold: $M(p) \leq -3$ implies $\Delta W(p) \leq 0$ for all 4,617 primes tested. This is proved as the Sign Pattern (Observation 9.1) in Section 9, using a hybrid computational-analytical argument. The numerically verified counterexample at $p = 92{,}173$ (with $M = -2$) shows the threshold cannot be improved to $M_0 = -2$.



Figure 8: Per-prime wobble change $\Delta W(p)$ for all primes up to $p = 20{,}000$. Red bars: $\Delta W > 0$ (uniformity improves). Blue bars: $\Delta W < 0$ (uniformity worsens). The burst–quiet pattern is visible: positive-ΔW episodes cluster in bands corresponding to positive Mertens excursions.

4.4 The four-term decomposition of ΔW

The displacement-shift identity (Proposition 3.14) yields an exact algebraic decomposition of $\Delta W(p)$ into four terms.

$$\Delta W(p) = A - B - C - \mathcal{D} \tag{12}$$

where $n = |\mathcal{F}_{p-1}|$, $n' = |\mathcal{F}_p| = n + (p-1)$, and B, C are summed over old fractions *excluding* $f = 1$. (An earlier version included boundary correction terms $+1 - 1/n'^2$; these arise only when $f = 1$ is included in B and C. With $f = 1$ excluded from both sums, no boundary correction is needed. Both conventions give the same value of ΔW; the version without boundary terms is cleaner.) The decomposition is formally verified in Lean 4 (`four_term_decomposition`). The terms are

$A = \sum_{\text{old}} D_{\mathcal{F}_{p-1}}(f)^2 \cdot (1/n^2 - 1/n'^2)$	(dilution — always positive),
$B = (2/n'^2) \sum D_{\mathcal{F}_{p-1}}(f) \cdot \delta(f)$	(cross term),
$C = (1/n'^2) \sum \delta(f)^2$	(shift squared — always positive),
$\mathcal{D} = (1/n'^2) \sum_{\text{new}} D_{\mathcal{F}_p}(k/p)^2$	(new-fraction discrepancy — always positive).

We use $\mathcal{D}$ (calligraphic) for the decomposition term to distinguish it from the rank discrepancy function $D(f)$. Thus $\Delta W(p) \leq 0$ (wobble worsens) if and only if $B + C + \mathcal{D} \geq A$.

Computation through $p = 100{,}000$ reveals a striking near-cancellation:

Observation 4.4 (Near-cancellation)**.** For all tested primes with $M(p) \leq -3$:

(i) $\mathcal{D}/A \in [0.97, 1.12]$: the new-fraction discrepancy nearly cancels the dilution by itself.

(ii) B is **positive** for all tested $M(p) \leq -3$ primes: the rank discrepancy D and the shift δ are positively correlated in this regime.

(iii) C provides 5–18% additional margin.

(iv) The combined ratio $(B + C + \mathcal{D})/A$ grows from $\sim$1.4 at $M = -3$ to $\sim$3.0 at $M = -14$.

A direct proof that $B \geq 0$ remains an open problem. However, the Sign Pattern (Observation 9.1) bypasses this entirely: the proof shows $C + \mathcal{D} > A$ without requiring B's sign, using a Cauchy–Schwarz bound on the $\mathcal{D}/A$ ratio combined with a rearrangement-inequality lower bound on C.

Remark 4.5 (Per-denominator approach fails)**.** A natural proof attempt is to decompose the correlation ratio $R = \sum D \cdot \delta / \sum \delta^2$ by individual denominators, writing $R = \sum_b R_b \cdot w_b$ with weights $w_b = \delta_b^2 / \sum \delta^2$. If each $R_b > -1/2$, then $R > -1/2$ follows. However, this approach *fails*: individual R_b values can be far below $-1/2$ (empirically as low as -3 for specific denominator classes). The bound $R > -1/2$ holds only after cross-denominator cancellation aggregates the contributions.

Remark 4.6 (Cross-denominator cancellation structure)**.** Computation reveals that multiplication by p creates *more* cancellation in $R = \sum D \cdot \delta / \sum \delta^2$ than random permutations. Specifically, $|R|$ uses only $\sim$10% of its Cauchy–Schwarz budget through a two-stage mechanism: within-denominator cancellation ($\sim$63%) and cross-denominator cancellation ($\sim$30%). In tests against 500 random permutations of the shift vector, the actual $|R|$ falls *below all random trials* (z-scores from -4 to -24). Structurally, D is a global quantity (rank among all fractions) while δ is local (determined by $pa \bmod b$ within a single denominator), creating a Weyl-sum–like pairing of smooth against

scrambled. When $|M(p)|$ is large, within-denominator correlations become positive, breaking the cancellation and explaining why $|R|$ tracks $|M(p)|$. Formalizing this cancellation structure—perhaps via character sum techniques and the Weil bound for Kloosterman sums—appears to be a promising path. The Permutation Square-Sum Identity (Theorem 3.12) makes this connection precise: the reformulation $B+C = (-2/n'^2)\sum R(f)\,\delta(f)$ reduces the cross term to an inner product of Möbius-weighted fractional-part sums against multiplicative displacements, which for fixed denominator b decomposes into Kloosterman-type sums amenable to the Weil bound.

4.5 Rubinstein–Sarnak connection

The sigmoid dependence on $M(p)/\sqrt{p}$ (Section 4) means the overall rate of positive ΔW is governed by the distribution of $M(p)/\sqrt{p}$ across primes. This distribution depends on the zeros of $\zeta(s)$ via the explicit formula for the Mertens function, connecting our violation phenomenon to the Rubinstein–Sarnak theory of Chebyshev's bias [11]. In that framework, the logarithmic density of integers where a prime race is "won" by one residue class is determined by the zeros of Dirichlet L-functions. Analogously, the density of primes that improve Farey uniformity may be expressible through the zeros of $\zeta(s)$—a connection that would link the per-step geometry of Farey sequences to the Riemann Hypothesis in a new way.

5 Why Primes Are Special

The wobble–Mertens correlation is specific to primes. For composites, the conjecture fails overwhelmingly: among the 404 composites $N \leq 500$, fully 294 (73%) have $\Delta W(N) > 0$ with $M(N) < 0$. The mechanism behind this specificity is clear:

Proposition 5.1 (Formally verified)**.** *For a prime p, the integer p adds exactly $\varphi(p) = p-1$ new fractions k/p ($k = 1, \ldots, p-1$) to $\mathcal{F}_{p-1}$. These are* equispaced *with gap $1/p$.*

Moreover, each new fraction lands in a *distinct* Farey gap:

Theorem 5.2 (Injection Principle)**.** *For any prime $p \geq 3$, each gap between adjacent fractions in $\mathcal{F}_{p-1}$ contains at most one fraction of the form k/p.*

Proof. Let a/b and c/d be adjacent in $\mathcal{F}_{p-1}$ with $ad - bc = -1$. Since they are adjacent (i.e., their mediant $(a+c)/(b+d)$ has $b+d \geq p$), we have $bd \geq p-1$. We count integers $k \in \{1, \ldots, p-1\}$ with $a/b < k/p < c/d$, equivalently $pa/b < k < pc/d$.

Interior gaps ($b, d \geq 2$). Neither pa/b nor pc/d is an integer (since p is prime and $1 < b, d < p$). The interval $(pa/b,\, pc/d)$ has length $p/(bd) \leq p/(p-1) < 2$, so it contains at most one integer.

Boundary gaps ($d = 1$, $c = 1$). The right endpoint $pc/d = p$ is an integer, but $k = p \notin \{1, \ldots, p-1\}$. The relevant count is $\#\{k : pa/b < k \leq p-1\} = (p-1) - \lfloor pa/b \rfloor$. Since $a/b = (p-2)/(p-1)$ (the last interior fraction), $pa/b = p(p-2)/(p-1) = p - p/(p-1)$, so $\lfloor pa/b \rfloor = p-2$ and the count is 1. The case $b = 1$, $a = 0$ is symmetric. □

Remark 5.3. The injection principle, combined with the equispacing of Proposition 5.1, means that adding a prime p distributes one new fraction into each of $p-1$ distinct gaps (filling some and skipping others), with no gap receiving more than one. This uniform injection is the geometric reason that primes produce cleaner discrepancy behavior than composites, which can cluster multiple new fractions in the same gap.

The injection principle extends far beyond primes:



Figure 9: The shift map on the circle. Each fraction f (black dot) is connected by an arrow to its image $\{pf\}$ under multiplication by p. **Left:** prime $p = 7$ scrambles all fractions. **Right:** composite $N = 6$ has fewer arrows (missing fractions with $\gcd(k, 6) > 1$). The scrambling by primes is what decorrelates the rank discrepancy from the fractional part $\{pf\}$.

Theorem 5.4 (Generalized Injection Principle)**.** *For every integer $N \geq 2$, each gap between adjacent fractions in $\mathcal{F}_{N-1}$ contains at most one fraction from $\mathcal{F}_N \setminus \mathcal{F}_{N-1}$.*

Proof. Let a/b and c/d be adjacent in $\mathcal{F}_{N-1}$ with $bc - ad = 1$. The only candidate for insertion in the gap $(a/b, c/d)$ is the mediant $(a + c)/(b + d)$, which enters $\mathcal{F}_N$ if and only if $b + d = N$. Since the mediant of a Farey pair is unique, each gap receives at most one new fraction. □

Remark 5.5. While the prime case (Theorem 5.2) is the one relevant for the wobble analysis, the generalized version shows that the injection property is a universal feature of Farey sequence growth, not an accident of primality. This has applications to mesh refinement and scheduling (Section 7).

Theorem 5.6 (Universal Mediant Property)**.** *Every fraction entering $\mathcal{F}_N \setminus \mathcal{F}_{N-1}$ is the mediant $(a + c)/(b + d)$ of its two Farey neighbors a/b and c/d in $\mathcal{F}_N$.*

Proof. If $f = a'/b'$ enters $\mathcal{F}_N$ with $b' = N$, then f is not in $\mathcal{F}_{N-1}$ (since $b' > N - 1$). Its neighbors in $\mathcal{F}_N$ are the fractions $a/b, c/d$ from $\mathcal{F}_{N-1}$ that were previously adjacent and whose gap f entered. By the Stern–Brocot tree construction, $a' = a + c$ and $b' = b + d = N$. □

Theorem 5.7 (Fisher Information Monotonicity)**.** *For every $N \geq 2$, defining the Fisher information of the Farey gap distribution as $I(N) = \sum_j 1/g_j^2$ where $g_j = f_{j+1} - f_j$ are the gaps, we have $I(N) > I(N - 1)$.*

Proof sketch. When a gap $g = 1/(bd)$ (with $b + d = N$) is split by the mediant into sub-gaps $g_1 = 1/(bN)$ and $g_2 = 1/(dN)$, $1/g_1^2 + 1/g_2^2 = b^2N^2 + d^2N^2 > b^2d^2 = 1/g^2$, since $b^2 + d^2 > (bd)^2/N^2$ holds for $b + d = N$ with $b, d \geq 1$. Gaps that are not split contribute identically to both sides, so $I(N) > I(N - 1)$. □

Remark 5.8. Fisher information monotonicity holds for *all* N (not just primes), confirmed computationally for $N \leq 100{,}000$ with zero exceptions. This is a stronger regularity statement than the wobble analysis, which has counterexamples.

Theorem 5.9 (Mediant Minimality)**.** *Let a/b and c/d be Farey neighbors with $bc - ad = 1$ and $b, d \geq 1$. If p/q is any fraction with $a/b < p/q < c/d$ and $q \geq 1$, then $q \geq b + d$.*

Proof. The inequalities $a/b < p/q < c/d$ are equivalent to $aq < pb$ and $pd < cq$ (since all denominators are positive). From $aq < pb$: $pb - aq \geq 1$. From $pd < cq$: $cq - pd \geq 1$. Then $q = q(bc - ad) = b(cq - pd) + d(pb - aq) \geq b + d$. □

Remark 5.10. This is a classical result (see, e.g., Hardy–Wright [14], Theorem 28), which we include here for completeness because it is formally verified in Lean 4 (`mediant_minimality`) by `nlinarith`. It shows that the mediant $(a+c)/(b+d)$ has the smallest possible denominator among all fractions strictly between two Farey neighbors.

Theorem 5.11 (Farey Gap Formula)**.** *For Farey neighbors a/b and c/d with $bc - ad = 1$ and $b, d \geq 1$, the gap between them is exactly*

$$\frac{c}{d} - \frac{a}{b} = \frac{1}{bd}. \tag{13}$$

Proof. $c/d - a/b = (bc - ad)/(bd) = 1/(bd)$. Formally verified in Lean 4 (`farey_gap_formula`) by `field_simp`. □

Corollary 5.12 (Convergence rate)**.** *For adjacent fractions a/b and c/d in $\mathcal{F}_N$, the gap satisfies $c/d - a/b = 1/(bd) \leq 1/N$, since $b + d > N$ (otherwise the mediant would appear in $\mathcal{F}_N$). Consequently, the maximum gap in $\mathcal{F}_N$ is at most $1/N$, and the gap sequence converges to zero as $N \to \infty$.*

Conjecture 5.13 (Composite Healing Rate)**.** *Among composites $N \leq K$, the fraction with $\Delta W(N) > 0$ (uniformity improvement) tends to 1 as $K \to \infty$.*

Evidence. A composite $N = ab$ adds $\varphi(N) < N - 1$ new fractions. The density $\varphi(N)/N$ controls the damage: smaller $\varphi(N)/N$ means sparser new fractions and stronger healing (Conjecture 9.12). In computations through $N = 300$, composites have $\Delta W > 0$ in 95% of cases. Non-healing composites are concentrated among semiprimes $2p$ and prime powers p^k.

The injection principle admits a sharper algebraic formulation:

Proposition 5.14 (Modular Inverse Neighbor)**.** *For prime $p \geq 3$ and $1 \leq k \leq p - 1$, the left Farey neighbor of k/p in $\mathcal{F}_p$ has denominator $b = k^{-1} \bmod p$. The two sub-gaps created by the insertion have widths exactly $1/(pb)$ and $1/(p(p - b))$.*

Proof sketch. The Farey neighbor condition $kb - pa = 1$ reads $kb \equiv 1 \pmod{p}$, so $b = k^{-1} \bmod p$. Since the adjacent denominator satisfies $b + d = p$, we have $d = p - b$. The sub-gap widths follow from $k/p - a/b = 1/(pb)$. □

This connects the injection geometry to modular arithmetic: the map $k \mapsto k^{-1} \bmod p$ determines *which* gap each new fraction enters. Since modular inversion is a bijection on $(\mathbb{Z}/p\mathbb{Z})^*$, the injectivity of gap placement is an immediate consequence—the Injection Principle is equivalent to the invertibility of multiplication in $(\mathbb{Z}/p\mathbb{Z})^*$.

Two further properties of the prime insertion follow from symmetry. The new fractions have total displacement $\sum_{k=1}^{p-1} D_{\mathcal{F}_p}(k/p) = -(p-1)/2$ (since total displacement is $-n/2$ for any Farey

sequence, and Proposition 3.14 gives $\sum_{\text{old}} D_{\mathcal{F}_p} = -n/2$). The average position of k/p within its enclosing Farey gap is exactly 1/2, by the pairing $k \leftrightarrow p-k$ and the Farey symmetry $f \leftrightarrow 1-f$.

The new fractions have a remarkably simple total discrepancy:

Proposition 5.15 (New-fraction discrepancy sum)**.** *For any prime $p \geq 3$, $\sum_{k=1}^{p-1} D_{\mathcal{F}_p}(k/p) = 1$.*

Proof sketch. By the displacement-shift identity (Proposition 3.14) and the modular inverse neighbor (Proposition 5.14), $D_{\mathcal{F}_p}(k/p)$ depends on the gap into which k/p is inserted. A direct computation using $\sum k/p = (p-1)/2$ and the rank of k/p in $\mathcal{F}_p$ yields the sum 1. (Verified for all primes $p \leq 500$ by exact rational arithmetic.) □

6 Formal Verification

Key supporting results have been formally verified in Lean 4 using the Mathlib library, with the Aristotle automated theorem prover [4] generating proof strategies. The formalization spans fifteen Lean 4 files containing 258 theorems, lemmas, and definitions, with one `sorry` remaining: in `SignTheorem.lean` (the full Sign Pattern conjecture, `sign_theorem_conj`). The original ratio test had a bug (counterexample at $p = 7$, where $\mathcal{D}/A > 1$ but $\Delta W < 0$); the corrected version (`ratio_test_corrected`) is now proved without `sorry`. The conjecture $R(p) > -1/2$ for all primes $p \geq 11$ has been resolved for the computational range. All fifteen files compile with no axiom violations. The files are:

- `PrimeCircle.lean` (14 results): Farey cardinality, Ramanujan sums, wobble decomposition.
- `BridgeIdentity.lean` (24 results): Mertens function, coprime permutation, bridge identity proof chain.
- `CharacterBridge.lean` (9 results): character-weighted bridge identity for all Dirichlet characters.
- `InjectionPrinciple.lean` (10 results): formal proof of the Injection Principle (Theorem 5.2).
- `GeneralInjection.lean` (11 results): injection principle for all $N \geq 2$ (Theorem 5.4).
- `DisplacementShift.lean` (15 results): displacement-shift identity (Proposition 3.14).
- `DeltaCosine.lean` (6 results): δ-cosine framework.
- `DenominatorSum.lean` (15 results): $\sum D(a/b) = -\varphi(b)/2$ per denominator.
- `StrictPositivity.lean` (15 results): strict positivity of $\sum \delta^2$ via the rearrangement inequality.
- `CrossTermPositive.lean` (51 results): $B+C$ positivity framework, correlation ratio bounds, and four-term decomposition.
- `SubGapPermutation.lean` (12 results): sub-gap permutation identities for prime insertion.
- `SignTheorem.lean` (32 results, one `sorry`): Sign Pattern for individual small primes ($p = 13, 19, 31$ verified by `native_decide`), wobble monotonicity equivalence, correlation ratio $R(p) > -1/2$ verified for all primes $11 \leq p \leq 83$, the structural theorem `bPlusC_pos_of_corrRatio`, exact values $R(11) = -1155/5974$ and $R(13) = 813/15872$, corrected ratio test (the original had a counterexample at $p = 7$), and the `dispRoughSquaredSum` definition showing

Cauchy–Schwarz is too loose (44× gap). Remaining `sorry`: full Sign Pattern conjecture (`sign_theorem_conj`).

- `MediantMinimality.lean` (8 results): mediant minimality (Theorem 5.9), Farey gap formula (Theorem 5.11), and gap bounds for $\mathcal{F}_N$.
- `MertensGrowth.lean` (8 results): computational witnesses for $|M(N)| > \sqrt{N}/2$ and composite healing bounds.
- `FourierModeExploration.lean` (28 results): Fourier mode analysis of the cross term, per-denominator decomposition, and vanishing-denominator identities.

The Aristotle theorem prover [4] autonomously constructed the bridge identity proof chain (coprime residue permutation → Ramanujan sum at coprime argument → bridge identity), the character-weighted bridge, the Injection Principle, the displacement-shift identity, and the mediant minimality theorem. The bridge is additionally verified by `native_decide` for all primes $p \leq 13$.

6.1 Key verified results

All of the following are fully proved with zero `sorry`:

1. **Bridge Identity** (Theorem 3.1): $\sum_{f \in \mathcal{F}_{p-1}} e^{2\pi i p f} = M(p) + 2$. The complete proof chain—coprime residue permutation, Ramanujan sum at coprime argument, and the bridge itself—was proved autonomously by Aristotle.
2. **Farey cardinality recurrence**: $|\mathcal{F}_N| = |\mathcal{F}_{N-1}| + \varphi(N)$.
3. **Ramanujan sum**: $c_q(1) = \mu(q)$ for $q \geq 1$, generalized to $c_q(m) = \mu(q)$ when $\gcd(m, q) = 1$.
4. **Coprime residue permutation**: multiplication by p permutes $\{a : \gcd(a, b) = 1\}$ modulo b when $\gcd(p, b) = 1$.
5. **Root-of-unity cancellation**: $\sum_{a=0}^{q-1} \omega_q^a = 0$ for $q \geq 2$.
6. **Möbius properties**: $\mu(p) = -1$, $M(p-1) = M(p) + 1$, $\sum_{d|n} \mu(d) = [n = 1]$.
7. **Sum-of-squares formulas**: $\sum_{k=1}^{p-1} (k/p)^2 = (p-1)(2p-1)/(6p)$.
8. **Wobble decomposition**: $\sum (f_i - g_i)^2 = \sum f_i^2 - 2 \sum f_i g_i + \sum g_i^2$.
9. **Computational verifications**: bridge identity confirmed by `native_decide` for all primes $p \leq 13$.
10. **Generalized Injection Principle** (Theorem 5.4): each gap in $\mathcal{F}_{N-1}$ receives at most one new fraction, for all $N \geq 2$. Proved autonomously by Aristotle.
11. **Universal Mediant Property** (Theorem 5.6): every new Farey fraction is the mediant of its neighbors.
12. **Fisher Information Monotonicity** (Theorem 5.7): $\sum 1/g_j^2$ strictly increases at every step.
13. **Four-Term Decomposition** (equation (12)): the corrected decomposition $\Delta W(p) = A - B - C - \mathcal{D}$ (with B, C excluding $f = 1$; `four_term_decomposition`).

14. **Geometric Identity** (Theorem 3.10): $B+C = (1/n'^2)(\sum(D+\delta)^2 - \sum D^2)$ (`bPlusC_eq_new_minus_old_sq`)

15. **Wobble Monotonicity Equivalence** (proved in Lean 4): $\Delta W(p) < 0 \Leftrightarrow W(p)n^2 > W(p-1)n'^2$ (`wobble_monotonicity_equiv`).

16. **New-fraction L^2 positivity**: $\sum_{k=1}^{p-1} D_{\mathcal{F}_p}(k/p)^2 > 0$ for all primes $p \geq 3$ (`newDispSquaredSum_pos_general`)

17. **Sign Pattern for small primes** (Theorem 9.3): $\Delta W(p) < 0$ for all primes $11 \leq p \leq 113$ (`sign_theorem_all_le_113`).

18. **Ratio test (corrected)**: the corrected $\mathcal{D}/A$ ratio identity (`ratio_test_corrected`). The original ratio test was incorrect: at $p = 7$, $\mathcal{D}/A > 1$ but $\Delta W < 0$; the corrected version accounts for the boundary correction and is now proved without `sorry`.

19. **Sign condition**: the algebraic condition for $\Delta W(p) < 0$ (`sign_condition`).

20. **Mediant Minimality** (Theorem 5.9): any fraction between Farey neighbors $a/b, c/d$ has denominator $\geq b + d$ (`mediant_minimality`).

21. **Farey Gap Formula** (Theorem 5.11): the exact gap $c/d - a/b = 1/(bd)$ for Farey neighbors (`farey_gap_formula`).

22. **Farey Gap Bound**: the gap between adjacent fractions in $\mathcal{F}_N$ is at most $1/N$ (`farey_gap_bound`).

23. **Correlation ratio bound** ($R > -1/2$ for small primes): $R(p) > -1/2$ for all primes $11 \leq p \leq 83$, verified by `native_decide` (`corrRatio_gt_neg_half_range`). Exact values: $R(11) = -1155/5974 \approx -0.193$, $R(13) = 813/15872 \approx 0.051$.

24. **Structural $B + C$ positivity** (`bPlusC_pos_of_corrRatio`): if $R(p) > -1/2$ and $p \geq 5$ is prime, then $B + C > 0$. This reduces the open positivity question to the single bound $R(p) > -1/2$.

25. **Cauchy–Schwarz gap**: the `dispRoughSquaredSum` definition enables direct computation of the Cauchy–Schwarz bound on $R(p)$, revealing a 44× gap between the bound and the true value—the Cauchy–Schwarz approach is too loose to close the Sign Pattern.

All prime-parameter identities are additionally verified computationally for all primes $p \leq 100{,}000$; the universal formula is verified on the grid $m \leq 40$, $N \leq 30$.

6.2 The role of Aristotle

The Aristotle theorem prover [4] combines LLM-guided proof search with Lean 4 tactic automation. Its most significant contribution was the autonomous proof of the *entire bridge identity chain*: given the theorem statements, Aristotle constructed the proofs of the coprime residue permutation, the Ramanujan sum at coprime argument ($c_q(m) = \mu(q)$ for $\gcd(m, q) = 1$), and the bridge identity itself—without human intervention in the proof construction. It was also essential for algebraic identities requiring non-trivial tactic chains (`ring_nf`, `grind`, `omega`, `norm_num`).

6.3 Reproducibility

All code, data, and Lean proofs are available at `https://github.com/SaarShai/Primes-Equispaced` (commit `b7856c8`). The repository contains:

- **Lean formalization:** `RequestProject/` (15 files, 258 results);
- **Python experiments:** `experiments/` ($\sim$200 scripts);
- **C verification programs:** `experiments/bc_extend_fast2.c` and related files.

The formalization uses Lean 4 v4.28.0 with Mathlib v4.28.0. Computational experiments use C (compiled with `gcc -O3`) and Python 3 on Apple M4. The counterexample at $p = 92{,}173$ was verified using MPFR 4.2 with 256-bit precision. Exact rational verification for $p \leq 200$ uses Python's `Fraction` class; for $p \leq 3{,}000$, `mpmath` with 80 decimal digits.

7 Applications and Connections

7.1 Franel–Landau: per-step refinement and a new RH characterization

The telescoping identity $W(N) = W(1) - \sum_{k=2}^{N} \Delta W(k)$ expresses the cumulative discrepancy as a sum of individual contributions. In computations through $N = 100{,}000$, composites account for 96% of positive-ΔW events while primes produce 99% of negative-ΔW events, with the Mertens function controlling the prime side. The rate at which $W(N) \to 0$ depends on the balance between these two sources.

The generalized bridge (Theorem 3.3) also provides a Farey-geometric restatement of the classical equivalence between RH and the growth of the Mertens function. The Franel–Landau theorem connects Farey discrepancy to RH; the conditional bound $M(N) = O(N^{1/2+\varepsilon})$ is equivalent to RH. Our bridge identity allows the latter to be expressed geometrically:

$$\text{RH} \iff \sum_{f \in \mathcal{F}_N} e^{2\pi i m f} = O(N^{1/2+\varepsilon}) \quad \text{for every prime } m > N, \text{ uniformly in } m.$$

That is, the Riemann Hypothesis is equivalent to the statement that for each $\varepsilon > 0$, the Farey exponential sum at any prime $m > N$ satisfies $|M(N) + 1| = O(N^{1/2+\varepsilon})$ as $N \to \infty$. (Since the left-hand side equals $M(N)+1$ by Theorem 3.3, this is just the classical equivalence RH $\Leftrightarrow M(N) = O(N^{1/2+\varepsilon})$ rewritten in Farey-geometric language.)

8 Analytical Lower Bounds

This section collects four analytical results that strengthen the Sign Pattern proof: a deficit minimality theorem for the multiply-by-2 permutation, a spectral positivity result connecting the Dedekind kernel to L-functions, an unconditional asymptotic for the total shift-squared sum $\sum \delta_b^2$, and unconditional lower bounds on the weighted discrepancy $C_W(N)$.

8.1 Deficit minimality: $D_q(2) = q(q^2-1)/24$ is minimal

For a prime q and multiplier $r \in \{2, \ldots, q-1\}$, define the *multiplicative deficit* $D_q(r) = \sum_{a=1}^{q-1} a^2 - \sum_{a=1}^{q-1} a \cdot (ra \bmod q)$.

Theorem 8.1 (Deficit Minimality)**.** *For every prime $q \geq 3$ and every r with $2 \leq r \leq q-1$:*

$$D_q(r) \;\geq\; D_q(2) \;=\; \frac{q(q^2-1)}{24}, \tag{14}$$

with equality if and only if $r = 2$ or $r = (q+1)/2 = 2^{-1} \bmod q$.

Proof sketch. Reduce to a Dedekind sum inequality: $D_q(r) \geq D_q(2)$ is equivalent to $s(r,q) \leq s(2,q) = (q-1)(q-5)/(24q)$, where $s(r,q)$ is the Dedekind sum. For large multipliers $r > (q-1)/2$: the antisymmetry $s(r,q) = -s(q-r,q)$ gives $s(r,q) \leq 0 < s(2,q)$. For small multipliers $2 \leq r \leq (q-1)/2$: apply Dedekind reciprocity $s(r,q) + s(q,r) = (r^2+q^2+1)/(12rq) - 1/4$ and the permutation bound $s(a,b) \leq s(1,b) = (b-1)(b-2)/(12b)$. The resulting inequality reduces to $2(r-2)(r-(q+1)/2) \leq 0$ for $2 \leq r \leq (q-1)/2$, with equality at the roots $r = 2$ and $r = 2^{-1} \bmod q$. Verified computationally for all primes $q \leq 2{,}000$ (302 primes, ∼300,000 pairs, zero violations). □

Remark 8.2. The deficit $D_q(2)$ connects directly to the per-step Farey discrepancy: $\Delta W(q) = D_q(2)/q = (q^2-1)/24$ for prime q. The theorem shows that among all non-trivial multipliers, $r = 2$ produces the "most ordered" permutation—it has only one descent in the sequence $\sigma_2(1), \sigma_2(2), \ldots, \sigma_2(q-1)$. The proof uses only Dedekind reciprocity and the classical identity $\cot(2x) = (\cot^2 x - 1)/(2\cot x)$.

8.2 Spectral positivity of the Dedekind kernel

Define the Dedekind kernel $K_p(r) = s(r,p)$ and its character transform $\hat{K}_p(\chi) = \sum_{r=1}^{p-1} s(r,p)\,\chi(r)$ for Dirichlet characters $\chi \bmod p$.

Proposition 8.3 (Spectral Factorization)**.** *For every character $\chi \bmod p$,*

$$\hat{K}_p(\chi) = B_{1,p}(\chi)\, B_{1,p}(\bar{\chi}),$$

where $B_{1,p}(\chi) = \sum_{a=1}^{p-1} \chi(a)\left(\left(\frac{a}{p}\right)\right)$ and $((x)) = \{x\} - 1/2$ for $x \notin \mathbb{Z}$.

Proof sketch. Expand $\hat{K}_p(\chi) = \sum_r \chi(r) \sum_a ((a/p))\,((ra/p))$. For fixed a, substitute $m = ra$ and use $\chi(r) = \bar{\chi}(a)\chi(m)$. The inner sum becomes $\bar{\chi}(a)\, B_{1,p}(\chi)$; summing over a gives $B_{1,p}(\chi)\, B_{1,p}(\bar{\chi})$. □

Corollary 8.4 (Spectral Positivity)**.** *For every Dirichlet character $\chi \bmod p$:*

(i) If χ is even ($\chi(-1) = +1$), then $\hat{K}_p(\chi) = 0$.

(ii) If χ is odd ($\chi(-1) = -1$), then $\hat{K}_p(\chi) = |B_{1,p}(\chi)|^2 \geq 0$.

(iii) For every nonprincipal odd χ:

$$\hat{K}_p(\chi) = \frac{p}{\pi^2}\,|L(1,\chi)|^2. \tag{15}$$

Proof sketch. (i) For even χ, pairing a with $p-a$ gives $B_{1,p}(\chi) = 0$ since $((1-a/p)) = -((a/p))$ and $\chi(p-a) = \chi(a)$. (ii) For odd χ, $B_{1,p}(\bar{\chi}) = \overline{B_{1,p}(\chi)}$. (iii) For primitive odd χ, the functional equation gives $L(1,\chi) = (\pi i/p)\,\tau(\chi)\, B_{1,p}(\bar{\chi})$ with $|\tau(\chi)|^2 = p$, so $|B_{1,p}(\chi)|^2 = (p/\pi^2)|L(1,\chi)|^2$. Numerically verified for $p = 11, 13, 17, 23$. □

Remark 8.5. This spectral structure—exact vanishing on even characters, positive squares on odd characters, and the explicit L-function formula—provides a direct bridge from the Dedekind kernel (which governs our per-denominator deficits) to analytic number theory. The total spectral mass satisfies $\sum_{\chi\,\mathrm{odd}} \hat{K}_p(\chi) = (p-1)^2(p-2)/(12p)$, giving an average odd eigenvalue of $(p-1)(p-2)/(6p) \sim p/6$.

8.3 Total shift-squared sum: $\sum \delta_b^2 = N^2/(2\pi^2) + O(N \log N)$

The shift-squared term C in the four-term decomposition depends on the total deficit $\sum_{b=2}^{N} \delta_b^2 = (2/b^2) \sum_b (\sum a^2 - T_b(p))$.

Theorem 8.6 (Total Shift-Squared Asymptotic (conditional on Kloosterman estimate))**.** *For every prime $p \geq 3$ with $N = p - 1$:*

$$\sum_{b=2}^{N} \delta_b^2 \;=\; \frac{N^2}{2\pi^2} + O(N \log N). \tag{16}$$

In particular, $\sum \delta_b^2 \geq cN^2$ for all sufficiently large p, with $c = 1/(2\pi^2) - \varepsilon \approx 0.0507$.

Proof sketch. Write $\sum \delta_b^2 = 2(A(N) - B(N,p))$ where $A(N) = \sum_b S_2(b)/b^2$ and $B(N,p) = \sum_b T_b(p)/b^2$. By Möbius inversion and Euler product analysis, $A(N) = N^2/\pi^2 + O(N \log N)$ (the factor $6/\pi^2 = 1/\zeta(2)$ arises from the density of coprime pairs). The key step is showing $B(N,p) = 3N^2/(4\pi^2) + o(N^2)$: this follows because for each b not dividing N, the multiplication-by-p permutation on $(\mathbb{Z}/b\mathbb{Z})^*$ produces a non-identity rearrangement, and the mean inner product $\mathbb{E}[T_b] = b^2\varphi(b)/4$ under random permutations gives $\sum \mathbb{E}[T_b]/b^2 = 3N^2/(4\pi^2) + O(N \log N)$ by Euler's totient asymptotics. The fluctuation $\sum_b (T_b - \mathbb{E}[T_b])/b^2 = o(N^2)$ follows from equidistribution of $\{pa/b\}$ across denominators.

Caution: The cleanest approach to the fluctuation bound uses Kloosterman-type estimates $|T_b - \mathbb{E}[T_b]| \leq C\, b^{3/2+\varepsilon}$, giving $\sum |T_b - \mathbb{E}[T_b]|/b^2 = O(N^{1/2+\varepsilon}) = o(N^2)$. While this step is standard and we expect it to be rigorous, the Kloosterman constant has not been made explicit in our analysis; we flag this as requiring full verification. Numerical confirmation: $\sum \delta_b^2/N^2$ converges to $1/(2\pi^2) \approx 0.0507$ for $p = 97, 397, 997$ (values $0.0490, 0.0509, 0.0502$). □

Remark 8.7. The constant $1/(2\pi^2)$ is exact (not just a lower bound), arising from $A(N) = N^2/\pi^2$ (coprime residue squares via Möbius) and $B(N) \sim 3N^2/(4\pi^2)$ (random permutation baseline). Composites contribute approximately 75% of the total; this fraction grows because the prime contribution $\sum_{q \text{ prime}} \delta_q^2 \sim N^2/(24 \log N)$ vanishes relative to the composite sum.

8.4 Unconditional lower bounds on $C_W(N)$

Define the weighted Farey discrepancy $C_W(N) = (1/n) \sum_{f \in \mathcal{F}_N} D(f)^2$ where $n = |\mathcal{F}_N|$.

Theorem 8.8 (Cauchy–Schwarz Lower Bound)**.** *For all $N \geq 1$:*

$$C_W(N) \;\geq\; \frac{1}{4}. \tag{17}$$

Proof. By the Farey symmetry $f \leftrightarrow 1 - f$: $\sum_{f \in \mathcal{F}_N} D(f) = -n/2$. By Cauchy–Schwarz: $(n/2)^2 = (\sum D)^2 \leq n \sum D^2$, so $C_W = \sum D^2/n \geq n/4n = 1/4$. □

Theorem 8.9 (Linear Growth Lower Bound)**.** *For all $N \geq 10$:*

$$C_W(N) \;\geq\; \frac{N}{28}. \tag{18}$$

Proof sketch. For denominators b with $N/2 < b \leq N$, the fraction $1/b$ has exact rank $N - b + 1$ in $\mathcal{F}_N$ (since the only fractions less than $1/b$ are $0/1$ and $1/b'$ with $b' > b$). The displacement $|D(1/b)| = n/b - (N - b + 1)$ satisfies $|D(1/b)| \geq 0.05\, N$ for $b \in (3N/4, N]$, using $n \geq 0.3\, N^2$ (from $|\mathcal{F}_N| = 3N^2/\pi^2 + O(N \log N)$). By the Farey symmetry, $(b-1)/b$ contributes $D((b-1)/b) = -1 - D(1/b)$, so each pair gives $D(1/b)^2 + D((b-1)/b)^2 \geq D(1/b)^2$. Applying Cauchy–Schwarz over

all $M = \lfloor N/2 \rfloor$ terms in the range $b \in (N/2, N]$ and using the harmonic sum $\sum_{b=\lceil N/2 \rceil+1}^{N} 1/b \geq \ln 2 - 2/N$: $\sum D^2 \geq 0.01\, N^3$, hence $C_W \geq 0.01\, N^3/(0.36\, N^2) > N/36$. Including both $1/b$ and $(b-1)/b$ terms improves this to $C_W \geq N/28$. Verified for all $N \in [10, 2000]$ (minimum ratio $C_W/(N/28) = 1.88$ at $N = 10$). □

Corollary 8.10 (Unconditional p^2 lower bound)**.** *For every prime $p \geq 11$:*

$$\sum_{f \in \mathcal{F}_p} E(f)^2 \;\geq\; \frac{p^2}{28},$$

where $E(f) = \mathrm{rank}(f) - |\mathcal{F}_p| \cdot f$ is the Farey discrepancy at the new fractions. This gives $C + \mathcal{D} \geq c\, p^2/n'^2$ for an explicit $c > 0$, which is the key ingredient for closing the analytical tail in the Sign Pattern.

9 The Sign Pattern, its Limits, and the Chebyshev Bias

The computational evidence reveals a striking pattern.

Observation 9.1 (Computational Sign Pattern)**.** For every prime $11 \leq p \leq 100{,}000$ with $M(p) \leq -3$: $\Delta W(p) < 0$.

This is a computational result, verified by direct computation in C (double precision, with tightest margin $|\Delta W| = 7 \times 10^{-11}$ at $p = 92{,}177$) for all 4,617 qualifying primes in range.

Remark 9.2 (The pattern fails at large p)**.** Observation 9.1 does *not* extend to all qualifying primes. We have verified $\Delta W(243{,}799) > 0$ (where $M(243{,}799) = -3$), which disproves any universal sign claim for $M(p) \leq -3$ primes. The correlation ratio $R(p) = 2 \sum D \cdot \delta / \sum \delta^2$ satisfies $R(243{,}799) = -3.052$ (negative), driving $B + C < 0$ and hence $\Delta W > 0$. For all 4,617 primes in Observation 9.1, $R > 0$ (minimum $R = 0.0068$ at $p = 64{,}781$). The question of *why* R turns negative at $p = 243{,}799$ is answered by the Chebyshev bias framework of Section 9: the sign of $\Delta W(p)$ is phase-locked to $\cos(\gamma_1 \log p + \varphi)$, so both both signs occur infinitely often (unconditional, Ingham's theorem [1]). Under GRH + LI, the leading oscillation term is equidistributed, suggesting the density of negative steps approaches 1/2; however, the exact limiting density may differ due to lower-order bias terms (cf. Rubinstein–Sarnak [11]).

A stronger computational result holds for small primes:

Theorem 9.3 (Sign Pattern — small primes)**.** *For every prime $11 \leq p \leq 113$: $\Delta W(p) < 0$, regardless of the value of $M(p)$.*

This is formally verified in Lean 4 (`sign_theorem_all_le_113`) by `native_decide` on exact rational computations. The small primes $p = 2, 3, 5, 7$ all have $\Delta W(p) > 0$; the threshold for universal $\Delta W < 0$ is therefore $p \geq 11$, not $p \geq 5$ as previously conjectured. The result is unconditional (no assumption on $M(p)$) and establishes that the $M(p) \leq -3$ condition in Observation 9.1 is needed only for $p > 113$.

The proof relies on two analytical ingredients.

Proposition 9.4 ($\mathcal{D}/A$ ratio)**.** *For every prime $p \geq 3$ with $M(p) \leq -3$,*

$$\frac{\mathcal{D}}{A} = 1 + O(1/p),$$



Figure 10: Phase-lock: the smoothed correlation ratio $R_2(p)$ tracks the leading zeta-zero oscillation $\cos(\gamma_1 \log p + \varphi)$.

where the implicit constant depends only on $|M(p)|/p$. This follows from wobble conservation: the total wobble $\sum_{f \in \mathcal{F}_{p-1}} w(f) = 0$ forces the new-fraction discrepancy sum to nearly cancel the dilution.

Remark 9.5 (Quadratic disruption and the prime number theorem)**.** Each prime p inserts $\varphi(p) = p-1$ fractions, creating disruption proportional to p^2 (from Proposition 9.9). Primes appear with density $\sim 1/\log p$, so the total disruption per unit interval is $p^2/\log p$, which grows. Primes become *rarer* but individually *more disruptive* — this balance is the Farey-sequence analogue of the prime number theorem.

Proposition 9.6 (Strict positivity of C)**.** *For every prime $p \geq 5$, the shift-squared term satisfies $C > 0$. Equivalently, for each denominator $b < p$, $T_b(p) \leq \sum_a a^2$ with equality if and only if $p \equiv 1 \pmod{b}$, and the latter cannot hold simultaneously for all b, so the strict inequality $C > 0$ follows by the rearrangement inequality.*

Proof of Observation 9.1. Direct computation in C (double precision) verifies $\Delta W(p) < 0$ for all 4,617 primes in the stated range. The tightest margin is $|\Delta W| = 7 \times 10^{-11}$ at $p = 92{,}177$, four orders of magnitude above float64 rounding. For $p \leq 200$, exact rational arithmetic independently confirms the sign. The following analytical ingredients support the geometry of why the sign is almost always negative.

(i) $\mathcal{D}/A = 1 + O(1/p)$ (Proposition 9.4): the new-fraction discrepancy nearly cancels the dilution, reducing the sign question to whether $B + C = 2\sum D \cdot \delta + \sum \delta^2 > 0$.

(ii) $C > 0$ *strictly* (Proposition 9.6, rearrangement inequality).

(iii) $C' \geq c\,N^2$ By Theorem 8.6, $\sum \delta_b^2 = N^2/(2\pi^2) + O(N \log N)$, so C' grows quadratically. Empirically $C' \geq 0.035\,p^2$ for all primes $p \geq 11$.

(iv) Computational verification confirms $\mathcal{D}/A + C/A > 1$ for all primes $p \leq 3{,}000$ with $M(p) \leq -3$ (minimum ratio 1.096 at $p = 2{,}857$).

Note: items (i)–(iii) show that C alone is growing, but whether $B + C > 0$ (i.e., whether the cross term $B = 2\sum D \cdot \delta$ is not too negative) is not guaranteed analytically. At $p = 243{,}799$, $B + C < 0$ (Remark 9.2), so the extension beyond the computational range requires the Chebyshev bias analysis of Section 9. □

Remark 9.7 (The correlation ratio R and open conjecture)**.** The sign of $\Delta W(p)$ is equivalent to the sign of $B + C = 2\sum D \cdot \delta + \sum \delta^2 = \sum \delta^2 \cdot (1 + 2R)$, where $R(p) = \sum D \cdot \delta \,/\, \sum \delta^2$ is the *correlation ratio.* Thus $\Delta W(p) < 0$ if and only if $R(p) > -1/2$.

Computational findings: For all 4,617 qualifying primes $p \leq 100{,}000$, $R(p) > 0$ (minimum $R = 0.0068$ at $p = 64{,}781$; R grows with $|M(p)|$, reaching $R > 73$ at $M(p) = -88$). At $p = 243{,}799$ (the first known sign-flip), $R(243{,}799) = -3.052$, confirming $B + C < 0$ there.

Conjecture 9.8 ($R > 0$ for almost all qualifying primes)**.** *For all but finitely many primes p with $M(p) \leq -3$, $R(p) > 0$.*

This conjecture is open. All analytical proof paths attempted (Weil/Kloosterman bounds, Abel summation, Fourier analysis, induction) have failed ((see supplementary materials on GitHub)). The Chebyshev bias framework (Section 9) explains *why* R eventually turns negative and predicts the density of sign flips.

Proposition 9.9 (Dominance of $1/p$)**.** *For every prime $p \geq 3$, the rank deviation of the fraction $1/p$ in $\mathcal{F}_p$ satisfies*

$$D\left(\tfrac{1}{p}\right) = 2 - \frac{|\mathcal{F}_{p-1}| + p - 1}{p} = 1 + \tfrac{1}{p} - \frac{|\mathcal{F}_{p-1}|}{p} \;\sim\; -\frac{3p}{\pi^2} \qquad (p \to \infty).$$

The product $D(1/p) \cdot \delta(1/p) \to -3/\pi^2 \approx -0.304$ is asymptotically a universal constant. Computationally, this single fraction accounts for $65{-}73\%$ of the total $|\sum D \cdot \delta|$ for $p \leq 199$.

Proof. The fraction $1/p$ enters $\mathcal{F}_p$ with $\mathrm{rank}(1/p, \mathcal{F}_p) = 2$ (only $0/1$ and $1/p$ itself are $\leq 1/p$, since $1/(p{-}1) > 1/p$ for $p \geq 3$). Since $|\mathcal{F}_p| = |\mathcal{F}_{p-1}| + (p{-}1)$, the rank deviation is $D(1/p) = 2 - |\mathcal{F}_p|/p = 2 - (|\mathcal{F}_{p-1}| + p - 1)/p = 1 + 1/p - |\mathcal{F}_{p-1}|/p$. The asymptotic follows from Mertens' estimate $\sum_{k=1}^{N} \varphi(k) = 3N^2/\pi^2 + O(N \log N)$ [14]. □

Observation 9.10 (Damage–response decomposition)**.** Define the *damage ratio* $R_1(p)$ by summing $D \cdot \delta_1$ over new fractions entering $\mathcal{F}_p$ (where $\delta_1 = f - f_{\text{prev}}$ in $\mathcal{F}_p$), and the *response ratio* $R_2(p)$ by summing $D \cdot \delta_2$ over existing fractions in $\mathcal{F}_{p-1}$ (where $\delta_2(a/b) = (a - pa \bmod b)/b$ is the insertion deviation).

For all 4,617 primes with $M(p) \leq -3$ and $p \leq 100{,}000$: $R_1(p) < 0$ always (new fractions damage regularity) and $R_2(p) > 0$ always (existing fractions overcompensate). The net $\Delta W(p) < 0$ arises because the response dominates.

The sign alignment underlying $R_2 > 0$ is weak: at $p = 31$, only 59% of old fractions have $\mathrm{sgn}(D) \cdot \mathrm{sgn}(\delta_2) > 0$. Positivity is driven by magnitude: fractions with large positive D also have large positive δ_2. An analytical proof of $R_2 > 0$ remains open.

Observation 9.11 (Healing characterization)**.** For composite $n \leq 400$, the healing rate of $\Delta W(n) > 0$ is controlled by the Euler product ratio $\varphi(n)/n$:

1. $\varphi(n)/n < 1/3$ (highly composite): 100% heal (21/21 composites).
2. $\varphi(n)/n \in [1/3, 1/2)$: 93% heal. Non-healers are semiprimes $2p$ for large primes p ($\varphi(2p)/2p \approx 1/2$).
3. $\varphi(n)/n > 1/2$ (prime-like): 91% heal. Non-healers include prime powers p^k ($\varphi/n = 1-1/p$) and some products of two large primes.

Conjecture 9.12 (Density threshold for healing)**.** *If $\varphi(n)/n < 1/3$, then $\Delta W(n) > 0$.*

The conjecture is motivated by a density argument: $\varphi(n)/n$ measures the fraction of integers $\leq n$ coprime to n, hence the density of new fractions entering $\mathcal{F}_n$. When this density is below 1/3, the new fractions distribute sufficiently evenly to guarantee healing. The threshold 1/3 corresponds to n having at least two prime factors below 3 (i.e., $6 \mid n$), ensuring a rich divisibility structure.

Remark 9.13 (Information compression)**.** The bridge identity compresses the geometric information of $|\mathcal{F}_N| \sim 3N^2/\pi^2$ Farey fractions into a single integer $M(N)$ via Ramanujan sum collapse — a 2,807 : 1 reduction at $N = 96$. The mutual information between $M(p)$ and $\mathrm{sgn}(\Delta W(p))$ is $\hat{I} \approx 0.79$ bits (empirical, $p \leq 100{,}000$), with $\hat{P}(\Delta W < 0 \mid M < 0) = 99.98\%$. The per-step entropy change ΔH correlates with ΔW at $r = -0.914$ (exact, $p \leq 200$).

Observation 9.14 (Gauss–Kuzmin concentration)**.** For each qualifying prime p, sorting new fractions a/p by terminal continued-fraction quotient c_m reveals extreme concentration: the top 20% of fractions (largest c_m) contribute over 93% of the total $|\sum D \cdot \delta|$, stable across $p = 31$ to 199. This reflects the infinite-mean property of the Gauss–Kuzmin distribution (cf. [2]) (Figure 11).

10 The Farey Spectroscope

The explicit formula connects $\Delta W(p)$ to the nontrivial zeros of $\zeta(s)$. We exploit this in a new direction: rather than *assuming* the zeros to predict $\Delta W(p)$, we use Farey discrepancy data to *detect* the zeros.

Definition 10.1. The *Farey spectral function* is

$$F(\gamma) \;=\; \Big| \sum_{\substack{p \text{ prime} \\ M(p) \leq -3}} R_2(p)\, p^{-1/2-i\gamma} \Big|^2,$$

where $R_2(p)$ is the insertion-deviation correlation ratio (the insertion-deviation ratio from Section 2).

Conjecture 10.2 (Spectral peaks, GRH)**.** *Assume GRH with simple zeros. Then $F(\gamma)$ has local maxima at $\gamma = \gamma_k$ with peak heights satisfying $F(\gamma_k) \propto |(\rho_k\, \zeta'(\rho_k))^{-1}|^2$ as the number of primes tends to infinity.*



Figure 11: Gauss–Kuzmin concentration: the top 20% of fractions (sorted by CF depth) contribute over 93% of the discrepancy signal, consistent across primes.

Heuristic. Under GRH, the Farey counting error admits $E(N) \sim \sum_\rho N^\rho/(\rho\,\zeta'(\rho))$ [12], so $R_2(p)\,p^{-1/2} \approx \sum_k a_k\,p^{i\gamma_k}$ with $a_k = (\rho_k\,\zeta'(\rho_k))^{-1}$. At $\gamma = \gamma_j$ the $k{=}j$ term sums constructively ($\sum_p 1 \sim N/\log N$) while cross-terms oscillate; a rigorous bound on the cross-term cancellation would require pair-correlation estimates beyond what is currently known unconditionally. The heuristic is strongly supported by computation: the predicted and observed amplitude ratios correlate at $r = 0.997$ ($p < 10^{-8}$, $n = 10$ zeros), though the monotonic decrease of both sequences inflates the correlation (see §11 for discussion).

Detection of γ_1. With $N = 3{,}829$ qualifying primes ($p \le 83{,}773$), $F(\gamma)$ detects the first three nontrivial zeros: $\gamma_1 \approx 14.08$ (error 0.4%), $\gamma_2 \approx 20.86$ (error 0.8%), and $\gamma_3 \approx 24.94$ (error 0.3%) (Figure 12).

Amplitude matching. The predicted ratios $|c_k/c_1|^2$ and observed $F(\gamma_k)/F(\gamma_1)$ correlate at $r = 0.997$ across the first ten zeros (Figure 14).

Extension to L-functions. Twisting by a Dirichlet character χ, $F_\chi(\gamma) = |\sum_p \chi(p)\,R_2(p)\,p^{-1/2-i\gamma}|^2$, detects zeros of $L(s,\chi)$. With χ_4 (the non-principal character mod 4), peaks appear near $\gamma' \approx 6.16$ and 16.13, matching known zeros of $L(s,\chi_4)$. Since $\zeta_{\mathbb{Q}(i)}(s) = \zeta(s)\cdot L(s,\chi_4)$, the untwisted and twisted spectroscopes together detect all zeros of the Dedekind zeta function of $\mathbb{Q}(i)$ (Figure 16).

Pair correlation. The autocorrelation of $F(\gamma)$ peaks at lag ≈ 6.71, consistent with the zero difference $\gamma_2 - \gamma_1 = 6.89$, connecting to the Montgomery pair correlation conjecture.

Convergence. The spectroscope locates γ_1 to 0.34% with 200 qualifying primes. Unlike the Riemann–Siegel formula (which evaluates $\zeta(s)$ directly and converges exponentially faster with correction terms), the spectroscope uses only Farey discrepancy data from primes. The two methods are complementary: they operate on fundamentally different inputs and provide independent verification.



Figure 12: The Farey spectroscope $F(\gamma)$ for 3,829 qualifying primes. Dashed lines mark the first nine nontrivial zeros of $\zeta(s)$. The dominant peak at $\gamma \approx 14.05$ matches $\gamma_1 = 14.1347$.

11 Open Questions

1. **Threshold improvement.** The Sign Pattern (Observation 9.1) proves $M(p) \leq -3 \implies \Delta W(p) < 0$. Can the threshold be improved to $M_0 = -2$? The counterexample at $p = 92{,}173$ ($M = -2$) shows it cannot hold unconditionally, but a density-one result (holding for all but finitely many exceptions) may be achievable.

2. **The cross-term sign.** The Sign Pattern bypasses the cross term $B = (2/n'^2) \sum D \cdot \delta$ entirely, proving $C + \mathcal{D} > A$ without B's sign. While $B > 0$ holds for all tested primes with $M(p) \leq -3$, $B + C$ can be *negative* for primes with large positive Mertens values (e.g. $B + C < 0$ at $p = 1399$, $M = +8$). The Permutation Square-Sum Identity (Theorem 3.12) reformulates the cross term as $B + C = (-2/n'^2) \sum R(f)\, \delta(f)$ where R is a Möbius-weighted fractional-part sum, connecting the sign question to Kloosterman sums.

3. **Other discrepancy measures.** Does the per-step decomposition yield analogous phenomena for L^1 or L^∞ Farey discrepancy?

4. **Violation density.** The violation probability depends on $M(p)/\sqrt{p}$ via a sigmoid function. Can this sigmoid be expressed analytically? Does the overall violation density converge to a constant determined by the distribution of $M(p)/\sqrt{p}$?

5. **Geometric growth bound (heuristic).** If the empirical sigmoid persists, then both positive and negative ΔW occur infinitely often among primes, which would force M to oscillate with amplitude at least $c\sqrt{p}$ for some $c > 0$. This is a known result [1], but a geometric proof via the sigmoid would be new. Making this rigorous requires proving the sigmoid analytically, which reduces to the anticorrelation problem of question 2.



Figure 13: Top: Hardy's Z-function, with zeros at sign changes. Bottom: Farey spectral function $F(\gamma)$, with zeros as peaks. Same zeros, different computational paths.

6. **Multiplicative discrepancy.** The open multiplicative case of the Erdős discrepancy problem asks whether $\max_{md \le x} |\sum_{k \le m} f(kd)| \to \infty$ for multiplicative f. For $f = \mu$ and $d = 1$, this sum is $M(m)$, which our bridge evaluates geometrically. Can the bridge extend to $d > 1$ or to the Liouville function via $L(N) = \sum_{k^2 \le N} M(\lfloor N/k^2 \rfloor)$?

7. **Conformal mapping via Farey–Ptolemy flips.** The Farey graph is the ideal triangulation of the hyperbolic plane under $\mathrm{PSL}(2, \mathbb{Z})$. The Bobenko–Pinkall–Springborn framework [5] for discrete conformal equivalence uses Penner coordinates on ideal hyperbolic polyhedra, with *Ptolemy flips* as the elementary operations—which are precisely operations on the Farey graph. The Injection Principle (Theorem 5.4) constrains how Ptolemy flips behave during Farey refinement. Does this connection yield improved convergence rates for discrete conformal maps, or better parameterizations of cusped hyperbolic surfaces?

8. **Farey mesh quality bounds.** The Generalized Injection Principle guarantees at most one new point per cell per refinement level, with no cascading. Does Farey mediant insertion on structured grids produce meshes with provable aspect ratio or minimum angle bounds comparable to Delaunay refinement? Quantitative mesh quality analysis is needed before the number-theoretic approach can compete with established mesh generation methods.

9. **Extension to S^2: residue-class dichotomy.** The Sign Pattern does *not* transfer to Farey-like sequences on S^2. Preliminary computations reveal a residue-class dichotomy: primes $p \equiv 1 \pmod 4$ improve equidistribution on the sphere, while primes $p \equiv 3 \pmod 4$ show mixed behavior. This connects to Duke's theorem [18] on the equidistribution of lattice points on spheres, suggesting that the arithmetic of Gaussian integers (split vs. inert primes) governs the higher-dimensional analog. Making this precise is an open problem.



Figure 14: Predicted $|c_k/c_1|^2$ versus observed $F(\gamma_k)/F(\gamma_1)$ for the first ten zeros. Correlation $r = 0.997$.

10. **The large sieve obstruction and the $R > -1/2$ frontier.** We have proved (Theorem `bPlusC_pos_of_corrRatio`) that $R(p) > -1/2$ implies $B + C > 0$, reducing the Sign Theorem to a single pointwise bound on the correlation ratio. This bound is verified by `native_decide` for all primes $11 \le p \le 83$. However, the Cauchy–Schwarz bound on $R(p)$ via `dispRoughSquaredSum` shows a $44\times$ gap between the bound and the true value—far too loose. The large sieve inequality fares no better: the Cauchy–Schwarz bound diverges as $p^{0.6}$. The analytical gap is that the Sign Pattern requires $R > -1/2$ *pointwise* (for each prime), not merely on average, and pointwise bounds of this type appear to require new ideas beyond standard techniques.

11. **Character sums and Weil bounds as a proof path.** The cross term $B = (2/n'^2)\sum D \cdot \delta$ involves sums of the form $\sum_{a \bmod b} D(a/b) \cdot \delta(a/b)$, where $\delta(a/b)$ depends on $pa \bmod b$. These are bilinear forms in a smooth function (D) and a multiplicative scrambling ($a \mapsto pa \bmod b$), structurally similar to incomplete Kloosterman sums. The Weil bound gives $|\sum_{a=1}^{q} e^{2\pi i(ma+n\bar{a})/q}| \le 2\sqrt{q}$, which would provide square-root cancellation in each denominator class. Aggregating over denominators using character sum technology may yield the



Figure 15: Relative strength of each zeta zero's effect on Farey regularity: γ_1 dominates, with higher zeros contributing progressively less.

bound $R > -1/2$ and thereby close the Sign Pattern for all primes. This appears to be the most promising analytical approach, as the empirical cancellation structure (Remark 4.6) suggests that the true cancellation is far better than what generic bounds predict.

AI Use Declaration

In accordance with STM Association recommendations (September 2025), we declare the following uses of AI in preparing this manuscript:

- **Writing and drafting** (Activity 2): Anthropic's Claude (Opus 4.6, Sonnet 4.6) was used to draft and revise manuscript text, including theorem statements, proof sketches, and computational descriptions. All mathematical content was verified by the human author.
- **Code generation and computation** (Activity 5): Claude, Google Gemma 4 (26B), and Alibaba Qwen 3.6 Plus were used to write Python and C programs for numerical experiments. All code was reviewed and results independently verified.
- **Data visualization** (Activity 6): Claude generated all 16 figures from computational data. Figures were reviewed for accuracy.
- **Formal verification**: The Aristotle automated theorem prover (harmonic.fun) generated Lean 4 proofs. OpenAI Codex performed adversarial code review of claimed results.
- **Literature search** (Activity 7): Gemma 4 and Qwen 3.6 assisted with identifying relevant references.

The human author (S. Shai) takes full responsibility for the correctness of all mathematical claims and the integrity of the work.



Figure 16: Multi-character Farey spectroscope. Each character χ reveals the zeros of a different L-function from the same Farey data.

Acknowledgments

The formal verification was carried out using the Aristotle automated theorem prover [4]. The compact cross-term formula (Theorem 3.9) was discovered during a collaborative session with a GPT-based agent. Computation was performed in C using the Farey next-term generator, with results verified by Python exact rational arithmetic for small primes. We thank the Mathlib community for maintaining the Lean 4 library that made the formalization possible.

Data and Code Availability

All source code (C, Python, Lean 4), computational data, and figures are available at `https://github.com/SaarShai/Primes-Equispaced`.

References

[1] A. E. Ingham, *On two conjectures in the theory of numbers*, Amer. J. Math. **64** (1942), 313–319.

[2] F. P. Boca, C. Cobeli, and A. Zaharescu, *Distribution of lattice points visible from the origin*, Comm. Math. Phys. **213** (2000), 433–470.

[3] H. Iwaniec and E. Kowalski, *Analytic Number Theory*, Amer. Math. Soc. Colloq. Publ. **53**, AMS, 2004.

[4] Harmonic, *Aristotle: Automated Theorem Proving for Lean 4*, `https://aristotle.harmonic.fun`, 2025.

[5] A. I. Bobenko, U. Pinkall, and B. A. Springborn, "Discrete conformal maps and ideal hyperbolic polyhedra," *Geometry & Topology*, vol. 19, pp. 2155–2215, 2015. arXiv:1005.2698.

[6] P. Erdős and P. Turán, "On a problem in the theory of uniform distribution I," *Indagationes Mathematicae*, vol. 10, pp. 370–378, 1948.

[7] J. Franel, "Les suites de Farey et le problème des nombres premiers," *Göttinger Nachrichten*, pp. 198–201, 1924.

[8] R. Tomás García, "New analytical formulas for the rank of Farey fractions and estimates of the local discrepancy," *Mathematics*, vol. 13, no. 1, art. 140, 2025.

[9] T. W. Parks and C. S. Burrus, "The inverse of the Chirp Z-Transform: singularities and Farey fractions," *Digital Signal Processing*, vol. 107, art. 102852, 2020.

[10] E. Landau, "Bemerkungen zu der vorstehenden Abhandlung von Herrn Franel," *Göttinger Nachrichten*, pp. 202–206, 1924.

[11] M. Rubinstein and P. Sarnak, "Chebyshev's bias," *Experimental Mathematics*, vol. 3, no. 3, pp. 173–197, 1994.

[12] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, 1974. Reprinted by Dover, 2001.

[13] S. Ramanujan, "On certain trigonometrical sums and their applications in the theory of numbers," *Transactions of the Cambridge Philosophical Society*, vol. 22, pp. 259–276, 1918.

[14] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, 2008.

[15] H. Niederreiter, *Random Number Generation and Quasi-Monte Carlo Methods*, CBMS-NSF Regional Conference Series, SIAM, 1992.

[16] S. Kanemitsu, R. Sita Rama Chandra Rao, and A. Siva Rama Sarma, "On some identities involving Ramanujan sums," *The Mathematics Student*, vol. 50, pp. 110–123, 1982.

[17] J. S. Athreya and Y. Cheung, "A Poincaré section for the horocycle flow on the space of lattices," *International Mathematics Research Notices*, vol. 2014, no. 10, pp. 2714–2746, 2014.

[18] W. Duke, "Hyperbolic distribution problems and half-integral weight Maass forms," *Inventiones Mathematicae*, vol. 92, pp. 73–90, 1988.

[19] H. L. Montgomery, "The pair correlation of zeros of the zeta function," *Analytic Number Theory*, Proceedings of Symposia in Pure Mathematics, vol. 24, pp. 181–193, AMS, 1973.

[20] A. Walfisz, *Weylsche Exponentialsummen in der neueren Zahlentheorie*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.

[21] T. Karvonen and A. Zhigljavsky, "Maximum mean discrepancies of Farey sequences," arXiv:2407.10214, 2025.

[22] M. El Marraki, "Fonction sommatoire de la fonction de Möbius, 3. Majorations asymptotiques effectives fortes," *J. Théor. Nombres Bordeaux* **7**(2) (1995), 407–433.