



Figure 4 | Scaling of the 4D-like response with the tilt angle θ . The linear dependence on θ reveals the nonlinear character of the response, demonstrating that it is induced by two independent perturbations, $\partial\varphi_x/\partial t$ and θ . The slope δI_y^{gs} is determined as a function of θ at $\varphi_y^{(0)} = 0.500(5)\pi$ by measuring the double-well imbalance when pumping along x , as described in Fig. 3 and using the same lattice depths. The solid line shows the slope that is expected for an ideal system. The fit errors for δI_y^{gs} are smaller than the size of the data points and the insets show two examples of the measurement of $I_y(\varphi_x)$ (for the values of θ indicated by the grey shading), as in Fig. 3a.

We verify this by measuring the peak slope δI_y^{gs} at $\varphi_y^{(0)} = \pi/2$ as a function of θ (Fig. 4). Doing so also provides another way of obtaining the second Chern number, by determining the slope of $\delta I_y^{\text{gs}}(\theta)$ (see Methods). This linear fit gives $\nu_2^{\text{exp}} = 1.01(8)$, where the error is determined as described above. Furthermore, we confirm that the peak slope at fixed θ scales with the depth of the short y lattice $V_{s,y}$ as expected (Extended Data Fig. 1, Methods). In particular, the direction of the nonlinear response is independent of $V_{s,y}$, indicating its robustness against perturbations of the system.

In conclusion, we present an observation of a dynamical 4D quantum Hall effect, opening up a route to studying higher-dimensional quantum Hall physics experimentally. Extending our work, additional density-type nonlinear responses that are implied by the intrinsic 4D symmetry of a 2D charge pump can be measured⁶. By adding a spin-dependent Yang–Mills gauge field, a dynamical version of the time-reversal-symmetric 4D quantum Hall effect, which exhibits a ground state with $\text{SO}(5)$ symmetry, could be realized⁶. Including interactions may yield intriguing fractional phases that originate in the 4D fractional quantum Hall effect⁴, similarly to previous proposals for 1D charge pumps²⁸, and might enable the study of open questions in the context of Floquet engineering¹⁵. Going beyond the limit of weak perturbations, quantized electric quadrupole moments could be observed in spatially frustrated systems with $\theta = \pi/4$ (ref. 29). Furthermore, a quantum Hall system with four extended dimensions might be realized with cold atoms²⁰ using recently demonstrated techniques for creating synthetic dimensions^{18,19}. In finite systems, this would permit the observation of boundary phenomena such as isolated Weyl points³⁰. Ultimately, the ability to experimentally realize 4D quantum Hall systems could provide insight into lattice quantum chromodynamics models based on the Yang–Mills theory⁷, and even quantum gravity⁴.

We note that, simultaneously with this work, complementary results on topological edge states in 2D photonic pumps have been obtained³¹.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 23 May; accepted 31 October 2017.

1. Klitzing, K. v., Dorda, G. & Pepper, M. New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance. *Phys. Rev. Lett.* **45**, 494–497 (1980).
2. Thouless, D. J., Kohmoto, M., Nightingale, M. P. & den Nijs, M. Quantized Hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.* **49**, 405–408 (1982).

3. Fröhlich, J. & Pedrini, B. in *Mathematical Physics 2000* (eds Fokas, A. et al.) 9–47 (Imperial College Press, 2000).
4. Zhang, S.-C. & Hu, J. A four-dimensional generalization of the quantum Hall effect. *Science* **294**, 823–828 (2001).
5. Thouless, D. J. Quantization of particle transport. *Phys. Rev. B* **27**, 6083–6087 (1983).
6. Kraus, Y. E., Ringel, Z. & Zilberberg, O. Four-dimensional quantum Hall effect in a two-dimensional quasicrystal. *Phys. Rev. Lett.* **111**, 226401 (2013).
7. Yang, C. N. & Mills, R. L. Conservation of isotopic spin and isotopic gauge invariance. *Phys. Rev.* **96**, 191–195 (1954).
8. Qi, X.-L. & Zhang, S.-C. Topological insulators and superconductors. *Rev. Mod. Phys.* **83**, 1057–1110 (2011).
9. Nayak, C., Simon, S. H., Stern, A., Freedman, M. & Das Sarma, S. Non-Abelian anyons and topological quantum computation. *Rev. Mod. Phys.* **80**, 1083–1159 (2008).
10. Laughlin, R. B. Quantized Hall conductivity in two dimensions. *Phys. Rev. B* **23**, 5632–5633 (1981).
11. Lu, L. et al. Experimental observation of Weyl points. *Science* **349**, 622–624 (2015).
12. Xu, S.-Y. et al. Discovery of a Weyl fermion semimetal and topological Fermi arcs. *Science* **349**, 613–617 (2015).
13. Hsieh, D. et al. A topological Dirac insulator in a quantum spin Hall phase. *Nature* **452**, 970–974 (2008).
14. Qi, X.-L., Hughes, T. L. & Zhang, S.-C. Topological field theory of time-reversal invariant insulators. *Phys. Rev. B* **78**, 195424 (2008).
15. Goldman, N., Budich, J. C. & Zoller, P. Topological quantum matter with ultracold gases in optical lattices. *Nat. Phys.* **12**, 639–645 (2016).
16. Lu, L., Joannopoulos, J. D. & Soljacic, M. Topological photonics. *Nat. Photon.* **8**, 821–829 (2014).
17. Sugawa, S., Salces-Carcoba, F., Perry, A. R., Yue, Y. & Spielman, I. B. Observation of a non-abelian Yang monopole: from new Chern numbers to a topological transition. Preprint at <http://arxiv.org/abs/1610.06228> (2016).
18. Mancini, M. et al. Observation of chiral edge states with neutral fermions in synthetic Hall ribbons. *Science* **349**, 1510–1513 (2015).
19. Stuhl, B. K., Lu, H.-I., Ayccock, L. M., Genkina, D. & Spielman, I. B. Visualizing edge states with an atomic Bose gas in the quantum Hall regime. *Science* **349**, 1514–1518 (2015).
20. Price, H. M., Zilberberg, O., Ozawa, T., Carusotto, I. & Goldman, N. Four-dimensional quantum Hall effect with ultracold atoms. *Phys. Rev. Lett.* **115**, 195303 (2015).
21. Price, H. M., Zilberberg, O., Ozawa, T., Carusotto, I. & Goldman, N. Measurement of Chern numbers through center-of-mass responses. *Phys. Rev. B* **93**, 245113 (2016).
22. Kraus, Y. E. & Zilberberg, O. Topological equivalence between the Fibonacci quasicrystal and the Harper model. *Phys. Rev. Lett.* **109**, 116404 (2012).
23. Switkes, M., Marcus, C. M., Campman, K. & Gossard, A. C. An adiabatic quantum electron pump. *Science* **283**, 1905–1908 (1999).
24. Kraus, Y. E., Lahini, Y., Ringel, Z., Verbin, M. & Zilberberg, O. Topological states and adiabatic pumping in quasicrystals. *Phys. Rev. Lett.* **109**, 106402 (2012).
25. Lohse, M., Schweizer, C., Zilberberg, O., Aidelsburger, M. & Bloch, I. A Thouless quantum pump with ultracold bosonic atoms in an optical superlattice. *Nat. Phys.* **12**, 350–354 (2016).
26. Nakajima, S. et al. Topological Thouless pumping of ultracold fermions. *Nat. Phys.* **12**, 296–300 (2016).
27. Rice, M. J. & Mele, E. J. Elementary excitations of a linearly conjugated diatomic polymer. *Phys. Rev. Lett.* **49**, 1455–1459 (1982).
28. Zhu, S.-L., Wang, Z.-D., Chan, Y.-H. & Duan, L.-M. Topological Bose–Mott insulators in a one-dimensional optical superlattice. *Phys. Rev. Lett.* **110**, 075303 (2013).
29. Benalcazar, W. A., Bernevig, B. A. & Hughes, T. L. Quantized electric multipole insulators. *Science* **357**, 61–66 (2017).
30. Hosur, P. & Qi, X. Recent developments in transport phenomena in Weyl semimetals. *C. R. Phys.* **14**, 857–870 (2013).
31. Zilberberg, O. et al. Photonic topological boundary pumping as a probe of 4D quantum Hall physics. *Nature* **553**, <https://doi.org/10.1038/nature25011> (2018).

Acknowledgements We acknowledge discussions with M. Aidelsburger and I. Carusotto. This work was funded by the European Commission (UQUAM, SIQS), the Deutsche Forschungsgemeinschaft (DF, FOR2414) and the Nanosystems Initiative Munich. M.L. was additionally supported by the Elitenetzwerk Bayern (EXQM), H.M.P. by the European Commission (FET Proactive, grant no. 640800 ‘AQuS’, and Marie Skłodowska–Curie Action, grant no. 656093 ‘SynOptic’) and the Autonomous Province of Trento (SiQuero), and O.Z. by the Swiss National Science Foundation.

Author Contributions M.L. and C.S. performed the experiment and data analysis. O.Z. proposed the experiment. All authors contributed to the theoretical analysis and to writing the paper. I.B. supervised the project.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. Correspondence and requests for materials should be addressed to I.B. (immanuel.bloch@physik.uni-muenchen.de).