

smaller than a hundredth of the RM_{src} value of FRB 121102 (Fig. 3), which is also about 500 times larger than those previously detected in fast radio bursts²⁷. The five other known fast radio bursts with polarimetric measurements present a heterogeneous picture, with a range of polarization fractions and rotation measures³. As previously considered²⁸, the large Faraday rotation of FRB 121102 further suggests that fast radio bursts with no detectable linear polarization may actually have very large $|RM|$, higher than 10^4 – 10^5 rad m⁻², that was undetectable because of the limited frequency resolution (0.4-MHz channels at 1.4 GHz) of the observations.

Monitoring the rotation measure and polarization angle of FRB 121102 with time, along with searches for polarization and Faraday rotation from the persistent source, can help differentiate among competing models. FRB 121102 is unusual not only because of its large rotation measure but also because it is the only known repeating fast radio burst. This may indicate that FRB 121102 is a fundamentally different type of source compared to the rest of the fast radio burst population; future measurements may investigate a possible correlation between fast radio burst repetition and rotation measure. Perhaps the markedly higher activity level of FRB 121102 compared to other known fast radio bursts is predominantly a consequence of its environment; for example, because these magnetized structures can also boost the detectability of the bursts via plasma lensing²⁹.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 1 November; accepted 20 November 2017.

- Lorimer, D. R., Bailes, M., McLaughlin, M. A., Narkevic, D. J. & Crawford, F. A bright millisecond radio burst of extragalactic origin. *Science* **318**, 777–780 (2007).
- Thornton, D. *et al.* A population of fast radio bursts at cosmological distances. *Science* **341**, 53–56 (2013).
- Petroff, E. *et al.* FRBCAT: the fast radio burst catalogue. *Publ. Astron. Soc. Aust.* **33**, e045 (2016).
- Spitler, L. G. *et al.* Fast radio burst discovered in the Arecibo pulsar ALFA survey. *Astrophys. J.* **790**, 101 (2014).
- Spitler, L. G. *et al.* A repeating fast radio burst. *Nature* **531**, 202–205 (2016).
- Petroff, E. *et al.* A survey of FRB fields: limits on repeatability. *Mon. Not. R. Astron. Soc.* **454**, 457–462 (2015).
- Chatterjee, S. *et al.* A direct localization of a fast radio burst and its host. *Nature* **541**, 58–61 (2017).
- Tendulkar, S. P. *et al.* The host galaxy and redshift of the repeating fast radio burst FRB 121102. *Astrophys. J.* **834**, L7 (2017).
- Bassa, C. G. *et al.* FRB 121102 is coincident with a star-forming region in its host galaxy. *Astrophys. J.* **843**, L8 (2017).
- Marcote, B. *et al.* The repeating fast radio burst FRB 121102 as seen on millisecond angular scales. *Astrophys. J.* **834**, L8 (2017).
- Bower, G. C., Wright, M. C. H., Falcke, H. & Backer, D. C. Interferometric detection of linear polarization from Sagittarius A* at 230 GHz. *Astrophys. J.* **588**, 331–337 (2003).
- Marrone, D. P., Moran, J. M., Zhao, J.-H. & Rao, R. An unambiguous detection of Faraday rotation in Sagittarius A*. *Astrophys. J.* **654**, L57–L60 (2007).
- Metzger, B. D., Berger, E. & Margalit, B. Millisecond magnetar birth connects frb 121102 to superluminous supernovae and long-duration gamma-ray bursts. *Astrophys. J.* **841**, 14 (2017).
- Piro, A. L. The impact of a supernova remnant on fast radio bursts. *Astrophys. J.* **824**, L32 (2016).
- Scholz, P. *et al.* The repeating fast radio burst FRB 121102: multi-wavelength observations and additional bursts. *Astrophys. J.* **833**, 177 (2016).
- Scholz, P. *et al.* Simultaneous X-ray, gamma-ray, and radio observations of the repeating fast radio burst FRB 121102. *Astrophys. J.* **846**, 80 (2017).
- Oppermann, N. *et al.* Estimating extragalactic Faraday rotation. *Astron. Astrophys.* **575**, A118 (2015).
- Akahori, T., Ryu, D. & Gaensler, B. M. Fast radio bursts as probes of magnetic fields in the intergalactic medium. *Astrophys. J.* **824**, 105 (2016).
- O'Sullivan, S. P. *et al.* Complex Faraday depth structure of active galactic nuclei as revealed by broad-band radio polarimetry. *Mon. Not. R. Astron. Soc.* **421**, 3300–3315 (2012).
- Haverkorn, M. in *Magnetic Fields in Diffuse Media* (eds Lazarian, A. *et al.*) 483–506 (Springer, 2015).
- Hunt, L. K. & Hirashita, H. The size-density relation of extragalactic H II regions. *Astron. Astrophys.* **507**, 1327–1343 (2009).
- Quataert, E., Narayan, R. & Reid, M. J. What is the accretion rate in Sagittarius A*? *Astrophys. J.* **517**, L101–L104 (1999).
- Eatough, R. P. *et al.* A strong magnetic field around the supermassive black hole at the centre of the Galaxy. *Nature* **501**, 391–394 (2013).
- Davidson, K. & Fesen, R. A. Recent developments concerning the Crab Nebula. *Annu. Rev. Astron. Astrophys.* **23**, 119–146 (1985).
- Harvey-Smith, L. *et al.* Faraday rotation of the supernova remnant G296.5+10.0: evidence for a magnetized progenitor wind. *Astrophys. J.* **712**, 1157–1165 (2010).
- Zhang, B. A 'Cosmic comb' model of fast radio bursts. *Astrophys. J.* **836**, L32 (2017).
- Masui, K. *et al.* Dense magnetized plasma associated with a fast radio burst. *Nature* **528**, 523–525 (2015).
- Petroff, E. *et al.* A real-time fast radio burst: polarization detection and multiwavelength follow-up. *Mon. Not. R. Astron. Soc.* **447**, 246–255 (2015).
- Cordes, J. M. *et al.* Lensing of fast radio bursts by plasma structures in host galaxies. *Astrophys. J.* **842**, 35 (2017).
- Manchester, R. N., Hobbs, G. B., Teoh, A. & Hobbs, M. The Australia telescope national facility pulsar catalogue. *Astron. J.* **129**, 1993–2006 (2005).

Acknowledgements We thank the staff of the Arecibo Observatory and the Green Bank Observatory for their help with our observations. We also thank B. Adebahr, L. Connor, G. Desvignes, R. Eatough, R. Fender, M. Haverkorn, A. Karastergiou, R. Morganti, E. Petroff, F. Vieyro and J. Weisberg for suggestions and comments on the manuscript. The Arecibo Observatory is operated by SRI International under a cooperative agreement with the National Science Foundation (AST-1100968), and in alliance with Ana G. Méndez-Universidad Metropolitana and the Universities Space Research Association. The Green Bank Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. Breakthrough Listen (BL) is managed by the Breakthrough Initiatives, sponsored by the Breakthrough Prize Foundation (<http://www.breakthroughinitiatives.org>). The research leading to these results received funding from the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007–2013). J.W.T.H. is a Netherlands Organisation for Scientific Research (NWO) Vidi Fellow and, together with D.M., K.G. and C.G.B., acknowledges funding for this work from ERC Starting Grant DRAGNET under contract number 337062. L.G.S. acknowledges financial support from the ERC Starting Grant BEACON under contract number 279702, as well as the Max Planck Society. A.M.A. is an NWO Veni Fellow. S.C., J.M.C., P.D., T.J.W.L., M.A.M. and S.M.R. are partially supported by the NANOGrav Physics Frontiers Center (NSF award 1430284). V.M.K. holds the Lorne Trottier Chair in Astrophysics & Cosmology and a Canada Research Chair and receives support from an NSERC Discovery Grant and Herzberg Prize, from an R. Howard Webster Foundation Fellowship from the Canadian Institute for Advanced Research (CIFAR), and from the FRQNT Centre de Recherche en Astrophysique du Québec. C.J.L. acknowledges support from NSF award 1611606. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. B.M. acknowledges support from the Spanish Ministerio de Economía y Competitividad (MINECO) under grants AYA2016-76012-C3-1-P and MDM-2014-0369 of ICCUB (Unidad de Excelencia 'María de Maeztu'). S.M.R. is a CIFAR Senior Fellow. P.S. holds a Covington Fellowship at DRAC.

Author Contributions A.S. led the development of the Arecibo observing functionality used here and discovered the first bursts near 4.5 GHz. L.G.S. is Principal Investigator of the Arecibo monitoring campaign. D.M. discovered the rotation measure and analysed the burst properties in detail. K.G. searched all Arecibo datasets near 4.5 GHz for bursts. J.W.T.H. led the discussion on the interpretation of the results and writing of the manuscript. A.M.A. guided the development of the rotation measure fitting code. G.H.H. and C.S. performed the rotation measure synthesis and deconvolution analysis. G.C.B., S.C., J.M.C., V.G., V.M.K., C.J.L., M.A.M. and D.M. also contributed to the writing of the manuscript and analysis. V.G. observed, searched and detected bursts from the GBT at 6.5 GHz as a part of the BL monitoring campaign of known fast radio bursts. A.P.V.S. is the Principal Investigator of the BL project. C.B. helped with the polarization calibration of the test pulsar. G.H. wrote a code to splice raw voltages across computer nodes. All other co-authors contributed to the interpretation of the analysis results and to the final version of the manuscript.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. Correspondence and requests for materials should be addressed to J.W.T.H. (J.W.T.Hessels@uva.nl).

Reviewer Information Nature thanks H. Falcke and the other anonymous reviewer(s) for their contribution to the peer review of this work.