

and Ni2. The Ni1–O1–Ni2 bond angle is about 145° and the length of the Ni2–O2 bond is approximately 2.5 Å, similar to that of bulk SNO. The proton first rotates about O1 while being bound to it, so that it enters the sub-surface layer of the SNO slab and reduces the O2–H distance to about 2.6 Å (image I₂ in Fig. 3c). This rotation of the O1–H bond distorts the surface layer considerably, which manifests itself in an increased Ni2–O2 distance of about 3.43 Å and a substantial change of the Ni1–O1–Ni2 angle (about 153°). This surface distortion process is associated with a barrier of about 0.9 eV. Further rotation of the O1–H bond brings the proton close to O2 (O2–H distance of approximately 1.5 Å; image I₃ in Fig. 3c) while increasing the inter-octahedral angle Ni1–O1–Ni2 to about 175°, and leads to the concurrent healing of the Ni2–O2 bond (image I₄ in Fig. 3c), which thus preserves the SNO framework upon intake of protons. Once the proton is intercalated into the SNO lattice, intra-octahedral proton hopping occurs even at room temperature, as seen in the *ab initio* simulations (Supplementary Video 3), indicating facile proton diffusion within SNO. The calculated energy barrier associated with the migration of protons within the SNO lattice is about 0.27 eV, which is low compared to those of other proton-conducting oxides²⁸ (0.4–0.6 eV). Volume expansion is observed in simulations after proton intercalation (Extended Data Fig. 10a–g and Supplementary Information section 7), which is in agreement with reflectometry measurements.

Density functional theory calculations (Fig. 3d) show that as each hydrogen atom is added to the supercell, its electron is transferred to a previously unoccupied Ni–O orbital of pristine SNO. The transfer of the electron can also be seen in the difference between the charge density of the combined HSNO system and that of the sum of SNO and H, which shows charge depletion around the H and a corresponding charge accumulation around the adjacent Ni and O (Extended Data Fig. 10h, i). Owing to electron–electron correlations, this state is shifted down into the valence band, while the remaining unoccupied states of that Ni are shifted up in energy. The bandgap remains almost unchanged until an electron has been added to each Ni atom. Both of these observations are consistent with the experimentally observed changes in X-ray absorption spectra and the increase in the electrical resistance of SNO that enables effective sensing.

Online Content Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

Received 2 May; accepted 31 October 2017.

Published online 18 December 2017.

- Chen, Y. W. *et al.* Atomic layer-deposited tunnel oxide stabilizes silicon photoanodes for water oxidation. *Nat. Mater.* **10**, 539–544 (2011).
- Uhlig, H. H. in *Uhlig's Corrosion Handbook* 3rd edn (ed. Revie, R. W.) Ch. 51 (John Wiley & Sons, 2011).
- Yuh, J. Design and control of autonomous underwater robots: a survey. *Auton. Robots* **8**, 7–24 (2000).
- Robison, B. H. Deep pelagic biology. *J. Exp. Mar. Biol. Ecol.* **300**, 253–272 (2004).
- Medarde, M. L. Structural, magnetic and electronic properties of RNiO₃ perovskites (R = rare earth). *J. Phys. Condens. Matter* **9**, 1679–1707 (1997).
- Middey, S. *et al.* Physics of ultrathin films and heterostructures of rare-earth nickelates. *Annu. Rev. Mater. Res.* **46**, 305–334 (2016).
- Catalan, G. Progress in perovskite nickelate research. *Phase Transit.* **81**, 729–749 (2008).
- Zhou, Y. *et al.* Strongly correlated perovskite fuel cells. *Nature* **534**, 231–234 (2016).
- Kritzer, P. Corrosion in high-temperature and supercritical water and aqueous solutions: a review. *J. Supercrit. Fluids* **29**, 1–29 (2004).
- Pelloquin, D., Barrier, N., Maignan, A. & Caignaert, V. Reactivity in air of the Sr₃Co₂O_{7-δ} RP = 2 phase: formation of the hydrated Sr₃Co₂O₅(OH)₂·xH₂O cobaltite. *Solid State Sci.* **7**, 853–860 (2005).
- Bedore, C. N. & Kajiwara, S. M. Bioelectric fields of marine organisms: voltage and frequency contributions to detectability by electroreceptive predators. *Physiol. Biochem. Zool.* **86**, 298–311 (2013).
- Baron, V. D. Electric discharges of two species of stargazers from the South China Sea (Uranoscopidae, Perciformes). *J. Ichthyol.* **49**, 1065–1072 (2009).
- Hirota, M. A method to measure ship's underwater electric field from deck. In *Proc. 2000 Int. Symp. on 'Underwater Technology'* 224–228 (IEEE, 2000); <http://ieeexplore.ieee.org/document/852547/>.

- Kim, J.-G., Joo, J.-H. & Koo, S.-J. Development of high-driving potential and high-efficiency Mg-based sacrificial anodes for cathodic protection. *J. Mater. Sci. Lett.* **19**, 477–479 (2000).
- Bennett, M. V. L., Wurzel, M. & Grundfest, H. The electrophysiology of electric organs of marine electric fishes. *J. Gen. Physiol.* **44**, 757–804 (1961).
- Mathewson, R., Mauro, A., Amatrik, E. & Grundfest, H. Morphology of main and accessory electric organs of *Narcine brasiliensis* (Olfers) and some correlations with their electrophysiological properties. *Biol. Bull.* **115**, 126–135 (1958).
- Kalmijn, A. J. The electric sense of sharks and rays. *J. Exp. Biol.* **55**, 371–383 (1971).
- Kalmijn, A. J. Electric and magnetic field detection in elasmobranch fishes. *Science* **218**, 916–918 (1982).
- Bellono, N. W., Leitch, D. B. & Julius, D. Molecular basis of ancestral vertebrate electroreception. *Nature* **543**, 391–396 (2017).
- Fields, R. D. The shark's electric sense. *Sci. Am.* **297**, 74–81 (2007).
- Josberger, E. E. *et al.* Proton conductivity in ampullae of Lorenzini jelly. *Sci. Adv.* **2**, e1600112 (2016).
- Brown, B. R. Sensing temperature without ion channels. *Nature* **421**, 495 (2003).
- Bard, A. J., Faulkner, L. R., Leddy, J. & Zoski, C. G. *Electrochemical Methods: Fundamentals and Applications* 1st edn, Vol. 2 (Wiley, 1980).
- Dailant, J. & Gibaud, A. (eds) *X-ray and Neutron Reflectivity: Principles and Applications* Vol. 770 (Springer, 2008).
- Medarde, M. *et al.* RNiO₃ perovskites (R = Pr, Nd): nickel valence and the metal-insulator transition investigated by x-ray-absorption spectroscopy. *Phys. Rev. B* **46**, 14975–14984 (1992).
- Mizokawa, T. *et al.* Electronic structure of PrNiO₃ studied by photoemission and x-ray-absorption spectroscopy: band gap and orbital ordering. *Phys. Rev. B* **52**, 13865–13873 (1995).
- Li, Z. *et al.* Correlated perovskites as a new platform for super-broadband-tunable photonics. *Adv. Mater.* **28**, 9117–9125 (2016).
- Kreuer, K. D. Proton-conducting oxides. *Annu. Rev. Mater. Res.* **33**, 333–359 (2003).

Supplementary Information is available in the online version of the paper.

Acknowledgements S.R. thanks K. Catania (Vanderbilt University) for discussions on bioelectric fields in marine organisms and B. Robinson and K. Benoit-Bird of the Monterey Bay Aquarium Research Institute for technical discussions on electroreception in sharks. We acknowledge financial support from the Army Research Office (W911NF-16-1-0289, W911NF-16-1-0042), National Science Foundation (DMR-1609898, DMR-1610215), Defense Advanced Research Projects Agency (grant D15AP00111), Office of Naval Research (grants N00014-16-1-2442 and N00014-12-1040) and Air Force Office of Scientific Research (grants FA9550-16-1-0159 and FA9550-14-1-0389). Use of the Center for Nanoscale Materials, an Office of Science user facility, was supported by the US Department of Energy (DOE), the Office of Science, Office of Basic Energy Sciences under contract number DE-AC02-06CH11357. This research used resources of the Advanced Photon Source, a US DOE Office of Science User Facility operated by Argonne National Laboratory under contract number DE-AC02-06CH11357. This research used resources of the National Energy Research Scientific Computing Center, which is supported by the Office of Science of the US DOE under contract number DE-AC02-05CH11231. An award of computer time was provided by the Innovative and Novel Computational Impact on Theory and Experiment (INCITE) programme. This research used resources of the Argonne Leadership Computing Facility at Argonne National Laboratory, which is supported by the Office of Science of the US DOE under contract DE-AC02-06CH11357. S.S.N. acknowledges support from the University of Massachusetts-Amherst through start-up funding. Part of the research described in this paper was performed at the Canadian Light Source, which is supported by the Canada Foundation for Innovation, Natural Sciences and Engineering Research Council of Canada, the University of Saskatchewan, the Government of Saskatchewan, Western Economic Diversification Canada, the National Research Council Canada and the Canadian Institutes of Health Research.

Author Contributions Z.Z., D.S. and S.R. conceived the study. Z.Z. and D.S. fabricated the SNO thin-film devices and performed the electrical, stability, sensing and electrochemical experiments. B.N., M.C. and S.K.R.S.S. performed the AIMD simulations and the nudged elastic band calculations to compute activation barriers. M.K. and K.M.R. performed first-principles electronic structure calculations. D.S. and J.A.D. performed neutron reflectivity measurements. Y.S., D.S. and H.Z. performed X-ray diffraction and X-ray reflectivity measurements. J.W.F., J.L., R.S., F.H. and R.C. performed X-ray absorption measurements. C.W. and N.Y. performed optical measurements and analysis. J.Z. and S.S.N. performed cross-sectional conducting AFM studies. K.R. and Z.Z. performed in-plane conducting AFM studies. Z.Z., D.S., B.N., S.K.R.S.S. and S.R. wrote the manuscript. All authors discussed the results and commented on the manuscript.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. Correspondence and requests for materials should be addressed to S.R. (shriram@purdue.edu).

Reviewer Information Nature thanks M. Lyons and the other anonymous reviewer(s) for their contribution to the peer review of this work.