

treated mice. Decreasing the number of cells entering the genomic DNA amplification and sequencing process increased the observed editing percentage to as high as 10% *Tmc1*<sup>Bth</sup> allele disruption, but also elevated background *Tmc1* indel rates of untreated mice to an average of  $0.82 \pm 0.57\%$  and a maximum of 1.6%, probably reflecting increased noise from processing of minute quantities of genomic DNA. No indel frequencies above that of untreated controls at any of the above-identified off-target sites were observed in Cas9–*Tmc1*–mut3–lipid-treated tissues (Extended Data Fig. 9c). Together, these observations confirm that Cas9–*Tmc1*–mut3–lipid treatment *in vivo* edits the *Tmc1* locus with no detected editing at GUIDE-seq-identified off-target loci.

An analysis of indel-containing *Tmc1* sequencing reads from treated *Tmc1*<sup>Bth/+</sup> mice allowed us to directly assess the allele specificity of Cas9–*Tmc1*–mut3 *in vivo*. Of 11,694 sequencing reads containing indels from four treated organ of Corti samples, 6,118 (52%) contained an intact nucleotide at *Tmc1* position 1,235. Of these, 5,736 (94%) contained modification of the mutant *Tmc1*<sup>Bth</sup> allele, whereas only 382 (6%) contained modification of the wild-type *Tmc1* allele (Fig. 4b). Therefore, samples after treatment on average contained 15-fold higher modification of the *Tmc1*<sup>Bth</sup> allele over the wild-type allele (Fig. 4b, c). These results demonstrate selective disruption of the *Tmc1*<sup>Bth</sup> allele in *Tmc1*<sup>Bth/+</sup> mice, consistent with observed hearing phenotypes, even though the *Tmc1*<sup>Bth</sup> and wild-type *Tmc1* alleles differ only at a single base pair.

This work shows that cationic lipid-mediated Cas9–sgRNA complex delivery *in vivo* can achieve allele-specific gene disruption in a mouse model of a human genetic disease, resulting in amelioration of the disease phenotype. Our results suggest that this approach has potential for the treatment of autosomal-dominant hearing loss related to hair cell dysfunction, and provide a complementary strategy to other approaches that use antisense oligos (ASOs) or RNA interference<sup>6,25</sup>. The genome editing strategy developed here may inform the future development of a DNA-free, virus-free, one-time treatment for certain genetic hearing loss disorders.

**Online Content** Methods, along with any additional Extended Data display items and Source Data, are available in the online version of the paper; references unique to these sections appear only in the online paper.

**Received 4 April; accepted 24 November 2017.**

**Published online 20 December 2017.**

- Angeli, S., Lin, X. & Liu, X. Z. Genetics of hearing and deafness. *Anat. Rec. (Hoboken)* **295**, 1812–1829 (2012).
- Marazita, M. L. *et al.* Genetic epidemiological studies of early-onset deafness in the U.S. school-age population. *Am. J. Med. Genet.* **46**, 486–491 (1993).
- Morton, C. C. & Nance, W. E. Newborn hearing screening—a silent revolution. *N. Engl. J. Med.* **354**, 2151–2164 (2006).
- Géléoc, G. S. & Holt, J. R. Sound strategies for hearing restoration. *Science* **344**, 1241062 (2014).
- Müller, U. & Barr-Gillespie, P. G. New treatment options for hearing loss. *Nat. Rev. Drug Discov.* **14**, 346–365 (2015).
- Shibata, S. B. *et al.* RNA interference prevents autosomal-dominant hearing loss. *Am. J. Hum. Genet.* **98**, 1101–1113 (2016).
- Pan, B. *et al.* Gene therapy restores auditory and vestibular function in a mouse model of Usher syndrome type 1c. *Nat. Biotechnol.* **35**, 264–272 (2017).
- Ahmed H., Shubina-Oleinik O. & Holt, J. R. Emerging gene therapies for genetic hearing loss. *J. Assoc. Res. Otolaryngol.* **18**, 649–670 (2017).
- Sacheli, R., Delacroix, L., Vandenackerveken, P., Nguyen, L. & Malgrange, B. Gene transfer in inner ear cells: a challenging race. *Gene Ther.* **20**, 237–247 (2013).

- Komor, A. C., Badran, A. H. & Liu, D. R. CRISPR-based technologies for the manipulation of eukaryotic genomes. *Cell* **168**, 20–36 (2017).
- Long, C. *et al.* Postnatal genome editing partially restores dystrophin expression in a mouse model of muscular dystrophy. *Science* **351**, 400–403 (2016).
- Yang, Y. *et al.* A dual AAV system enables the Cas9-mediated correction of a metabolic liver disease in newborn mice. *Nat. Biotechnol.* **34**, 334–338 (2016).
- Yin, H. *et al.* Therapeutic genome editing by combined viral and non-viral delivery of CRISPR system components *in vivo*. *Nat. Biotechnol.* **34**, 328–333 (2016).
- Kim, S., Kim, D., Cho, S. W., Kim, J. & Kim, J. S. Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome Res.* **24**, 1012–1019 (2014).
- Zuris, J. A. *et al.* Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing *in vitro* and *in vivo*. *Nat. Biotechnol.* **33**, 73–80 (2015).
- Pan, B. *et al.* TMC1 and TMC2 are components of the mechanotransduction channel in hair cells of the mammalian inner ear. *Neuron* **79**, 504–515 (2013).
- Camp, G. V. & Smith, R. *Hereditary Hearing Loss Homepage* <http://hereditaryhearingloss.org> (2017).
- Zhao, Y. *et al.* A novel DFNA36 mutation in TMC1 orthologous to the Beethoven (Bth) mouse associated with autosomal dominant hearing loss in a Chinese family. *PLoS One* **9**, e97064 (2014).
- Kurima, K. *et al.* Dominant and recessive deafness caused by mutations of a novel gene, TMC1, required for cochlear hair-cell function. *Nat. Genet.* **30**, 277–284 (2002).
- Kawashima, Y., Kurima, K., Pan, B., Griffith, A. J. & Holt, J. R. Transmembrane channel-like (TMC) genes are required for auditory and vestibular mechanosensation. *PLoS Arch.* **467**, 85–94 (2015).
- Vreugde, S. *et al.* Beethoven, a mouse model for dominant, progressive hearing loss DFNA36. *Nat. Genet.* **30**, 257–258 (2002).
- Fu, Y., Sander, J. D., Reyon, D., Cascio, V. M. & Joung, J. K. Improving CRISPR-Cas nuclease specificity using truncated guide RNAs. *Nat. Biotechnol.* **32**, 279–284 (2014).
- Tsai, S. Q. *et al.* GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat. Biotechnol.* **33**, 187–197 (2015).
- Zhang, F. *CRISPR Design* <http://crispr.mit.edu> (2015).
- Lentz, J. J. *et al.* Rescue of hearing and vestibular function by antisense oligonucleotides in a mouse model of human deafness. *Nat. Med.* **19**, 345–350 (2013).

**Supplementary Information** is available in the online version of the paper.

**Acknowledgements** This work was supported by DARPA HR0011-17-2-0049 (to D.R.L.), US NIH R01 EB022376 (to D.R.L.), R35 GM118062 (to D.R.L.), R01 DC006908 (to Z.-Y.C.), P30 DC05209 (to M.C.L.), R01 DC00138 (to M.C.L.), and R01 DC013521 (to J.R.H.). We are grateful for support from the David-Shulsky Foundation (to Z.-Y.C.), a Frederick and Ines Yeatts Hair Cell Regeneration grant (to Y.T., V.L., M.H., and Y.S.), the Bertarelli Foundation and the Jeff and Kimberly Barber Fund (to J.R.H.), the Broad Institute (to D.R.L. and Z.-Y.C.), and the HHMI (to D.R.L.). We thank H. Rees, S. Tsai, M. Packer, K. Zhao and D. Usanov for assistance.

**Author Contributions** X.G. and Y.T. designed the research, performed genome editing (X.G.) and hearing biology (Y.T.) experiments, analysed data, and wrote the manuscript. V.L., M.H., W.-H.Y., B.P., Y.-J.H., and H.W. designed experiments, performed hearing biology experiments and analysed data. D.B.P., M.C.L., and W.-J.K. designed hearing experiments and analysed data. Y.S. and S.Y. supported hearing biology experiments. J.H.H. analysed GUIDE-seq data. D.B.T. supported genome editing experiments. Y.L. and Q.X. designed and synthesized lipids. J.R.H., Z.-Y.C. and D.R.L. designed and supervised the research, and wrote the manuscript. All authors edited the manuscript.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare competing financial interests: details are available in the online version of the paper. Readers are welcome to comment on the online version of the paper. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations. Correspondence and requests for materials should be addressed to D.R.L. ([drliu@fas.harvard.edu](mailto:drliu@fas.harvard.edu)) and Z.Y.C. ([Zheng-Yi\\_Chen@meei.harvard.edu](mailto:Zheng-Yi_Chen@meei.harvard.edu)).