

# Mathematics of Symmetric Key Cryptography

## Unit-II

J. Bhuvana, ASP / CSE

August 29, 2020

# Session Objectives

- To learn about the modular arithmetic operations

# Session Outcomes

At the end of this session, participants will be able to discuss

- Modular arithmetic operations

# Agenda

## 1 Modular Arithmetic

# Presentation Outline

## 1 Modular Arithmetic

# Modular Arithmetic

- define modulo operator " $a \bmod n$ " to be remainder when  $a$  is divided by  $n$  where integer  $n$  is called the modulus
- $b$  is called a residue of  $a \bmod n$  since with integers can always write:  
 $a = qn + b$
- usually chose smallest positive remainder as residue  
ie.  $0 \leq b \leq n - 1$
- process is known as modulo reduction  
eg.  $-12 \bmod 7 = -5 \bmod 7 = 2 \bmod 7 = 9 \bmod 7$
- $a$  &  $b$  are congruent if:  **$a \bmod n = b \bmod n$**  when divided by  $n$ ,  $a$  &  $b$  have same remainder  
eg.  $100 \bmod 11 \cong 34 \bmod 11$  so 100 is congruent to 34 mod 11

# Modular Arithmetic Operations

$$\textcircled{1} \quad [(a \bmod n) + (b \bmod n)] \bmod n = (a + b) \bmod n$$

$$\textcircled{2} \quad [(a \bmod n) - (b \bmod n)] \bmod n = (a - b) \bmod n$$

$$\textcircled{3} \quad [(a \bmod n) \times (b \bmod n)] \bmod n = (a \times b) \bmod n$$

$$\begin{aligned} \text{e.g.} \quad & [(11 \bmod 8) + (15 \bmod 8)] \bmod 8 = 10 \bmod 8 = 2 ; \\ & (11 + 15) \bmod 8 = 26 \bmod 8 = 2 \end{aligned}$$

$$\begin{aligned} & [(11 \bmod 8) - (15 \bmod 8)] \bmod 8 = -4 \bmod 8 = 4 \\ & (11 - 15) \bmod 8 = -4 \bmod 8 = 4 \end{aligned}$$

$$\begin{aligned} & [(11 \bmod 8) \times (15 \bmod 8)] \bmod 8 = 21 \bmod 8 = 5 \\ & (11 \times 15) \bmod 8 = 165 \bmod 8 = 5 \end{aligned}$$

# Modulo 8 Addition Example

+	0	1	2	3	4	5	6	7
0	0	1	2	3	4	5	6	7
1	1	2	3	4	5	6	7	0
2	2	3	4	5	6	7	0	1
3	3	4	5	6	7	0	1	2
4	4	5	6	7	0	1	2	3
5	5	6	7	0	1	2	3	4
6	6	7	0	1	2	3	4	5
7	7	0	1	2	3	4	5	6



# Modulo 8 Multiplication

+	0	1	2	3	4	5	6	7
0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7
2	0	2	4	6	0	2	4	6
3	0	3	6	1	4	7	2	5
4	0	4	0	4	0	4	0	4
5	0	5	2	7	4	1	6	3
6	0	6	4	2	0	6	4	2
7	0	7	6	5	4	3	2	1

# Modular Arithmetic Properties

Property	Expression
Commutative Laws	$(w + x) \bmod n = (x + w) \bmod n$ $(w \times x) \bmod n = (x \times w) \bmod n$
Associative Laws	$[(w + x) + y] \bmod n = [w + (x + y)] \bmod n$ $[(w \times x) \times y] \bmod n = [w \times (x \times y)] \bmod n$
Distributive Law	$[w \times (x + y)] \bmod n = [(w \times x) + (w \times y)] \bmod n$
Identities	$(0 + w) \bmod n = w \bmod n$ $(1 \times w) \bmod n = w \bmod n$
Additive Inverse $(-w)$	For each $w \in \mathbb{Z}_n$ , there exists a $z$ such that $w + z \equiv 0 \bmod n$

# Additive and Multiplicative Inverses

- Additive inverse: For any integer  $a \in Z_m$ ,  $b \in Z_m$  is the additive inverse of  $a$  if
$$(a + b) \bmod m \cong 0$$
- Multiplicative inverse: For any integer  $a \in Z_m$ , integer  $b$  is the multiplicative inverse if
$$ab \cong 1 \bmod m$$

# Summary

- Modular arithmetic with integers