


Week 11,


MAT 110 DIFFERENTIAL CALCULUS & COORDINATE GEOMETRY

INTRODUCTION TO POLAR COORDINATES

Cartesian Coordinate




Polar Coordinate



We will get infinite number of polar coordinates for a specific "r" value such as:

$$\theta = \frac{\pi}{6} \pm 2n\pi$$



$n = 1, 2, 3, \dots$
+ve integers

$$(r, \frac{\pi}{6}) = (r, -\frac{11\pi}{6})$$

Consider as $-\frac{11\pi}{6}$ as it is rotating clockwise.

$$\frac{\pi}{6} + 2\pi = \frac{13\pi}{6}$$

Adding or subtracting $2n\pi$ from the original "θ" does not change our polar coordinate pt

O - pole

OX - polar axis

P - any point in the plane

OP ~ r ~ radius vector or radius


$$\angle XOP = \theta$$

Vectorial angle


Polar coordinate point :

$$P(r, \theta)$$

r → distance and always +ve.



$$(r, \frac{\pi}{6}) = (r, \frac{13\pi}{6})$$



$$\sin \theta = \frac{y}{r} \longrightarrow \operatorname{cosec} \theta = \frac{1}{\sin \theta} = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \longrightarrow \sec \theta = \frac{1}{\cos \theta} = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \longrightarrow \cot \theta = \frac{1}{\tan \theta} = \frac{x}{y}$$

$$x = r \cos \theta \quad \text{--- (i)} \quad y = r \sin \theta \quad \text{--- (ii)}$$


$$\text{Sq eqn (i)} \rightarrow x^2 = r^2 \cos^2 \theta$$

$$\text{Sq eqn (ii)} \rightarrow y^2 = r^2 \sin^2 \theta$$

$$\text{Add} \rightarrow x^2 + y^2 = r^2 (\cos^2 \theta + \sin^2 \theta)$$

$$r^2 = x^2 + y^2$$

POLAR EQUATIONS FOR CONIC SECTIONS



OP, OP_0, PP_0 form a triangle

$OP = r$
 $OP_0 = r_0$
 PP_0


} Arms of the triangle

$\triangle OPP_0$ is a right angle triangle

$$\cos(\theta - \theta_0) = \frac{r_0}{r} \quad \left[\cos \theta = \frac{x}{r} \right]$$

$r \cos(\theta - \theta_0) = r_0$ → Standard Eqn of Line in Polar coordinates

θ_0, r_0 are fixed



→ Given $r_0 = 2$ Eqn of L is:

$$\theta_0 = \frac{\pi}{3}$$

$$r \cos(\theta - \theta_0) = r_0$$

$$r \cos\left(\theta - \frac{\pi}{3}\right) = 2$$

We know,


$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\therefore r \cos\left(\theta - \frac{\pi}{3}\right) = \underbrace{r \cos \theta}_x \underbrace{\cos \frac{\pi}{3}}_{\frac{1}{2}} + \underbrace{r \sin \theta}_y \underbrace{\sin \frac{\pi}{3}}_{\frac{\sqrt{3}}{2}} = 2$$

$$\Rightarrow x \left(\frac{1}{2}\right) + y \left(\frac{\sqrt{3}}{2}\right) = 2$$

$\Rightarrow x + \sqrt{3}y = 4$ → Eqn of Line in cartesian coordinate for the given line in polar coordinate


CIRCLE



If the center of the circle is at the origin $(0,0)$, then the equation of the circle is defined as:
 $r = a$; while a is the radius of the circle

Center: $(r_0, \theta_0) \rightarrow P_0$; radius = a
 P is any pt on the circle with coordinate (r, θ)

Cosine formula for triangle:




$$c = \sqrt{a^2 + b^2 - 2ab \cdot \cos \theta}$$

For the $\triangle OPP_0$

$$a^2 = r^2 + r_0^2 - 2rr_0 \cos(\theta - \theta_0) \quad \text{--- (i)}$$

Eqn of circle in Polar coordinate system



If the circle passes through the origin then $r_0 = a$
 $a \rightarrow$ radius of the circle


\therefore Eqn of circle in polar coordinate system:

$$a^2 = r^2 + a^2 - 2ra \cos(\theta - \theta_0)$$

$$0 = r^2 - 2ar \cos(\theta - \theta_0)$$

$$r^2 = 2ar \cos(\theta - \theta_0)$$

$$r = 2a \cos(\theta - \theta_0) \quad \text{--- (ii)}$$



Center of circle P_0 is on "x" axis
 It passes through the origin
 It is on the right side of "x" axis

Eqn of circle:


$$r = 2a \cos(\theta - \theta_0) \rightarrow \text{Ref (ii) } \circ \circ \text{ circle passes through the origin}$$

$$\boxed{r = 2a \cos \theta} \quad \text{--- (iii) } \circ \circ \theta_0 = 0$$

As "x" axis on the negative direction

$$-r = 2a \cos \theta$$

$$\boxed{r = -2a \cos \theta} \quad \text{--- (iii) } \circ \circ$$



The circle is on the +ve "y" axis

radius = a

Angular coordinate = $\frac{\pi}{2}$

Eqn of circle:

$$r = 2a \cos\left(\theta - \frac{\pi}{2}\right) \rightarrow \text{Refer (i)}$$

$$r = 2a \sin \theta$$

$$\circ \circ \boxed{r = 2a \sin \theta}$$

The circle is on the -ve "y" axis


$$\boxed{r = -2a \sin \theta}$$

Recall $\cos(A-B)$

$$\begin{aligned} \cos\left(\theta - \frac{\pi}{2}\right) &= \cos \theta \cos \frac{\pi}{2} + \sin \theta \sin \frac{\pi}{2} \\ &= \cos \theta (0) + \sin \theta (1) \end{aligned}$$

$$\cos\left(\theta - \frac{\pi}{2}\right) = \sin \theta$$

Combine Eqn of Parabola, Ellipse, Hyperbola



Focus directrix eqn:

$$PF = e PD$$

$$r = e (OF - BF)$$

$$r = e (k - r \cos \theta)$$

$$r + e r \cos \theta = k e$$


$$r(1 + e \cos \theta) = k e$$

$$r = \frac{k e}{1 + e \cos \theta}$$

$\because PF \propto PD$

the ratio of PF & PD is a constant "e"

e \rightarrow eccentricity of the conic section



We can reach the eqn of parabola, ellipse, hyperbola from this general eqn above.

Parabola: $e = 1$, $r = \frac{k}{1 + \cos \theta}$


Ellipse: $0 < e < 1$, take $e = \frac{1}{2}$, $r = \frac{k/2}{1 + \frac{\cos \theta}{2}}$

$$r = \frac{k}{2 + \cos \theta}$$

Hyperbola: $e > 1$

Let's take $e = 2$

$$r = \frac{2k}{1+2\cos\theta}$$



Focus directrix eqn:

$$PF = e PD$$

$$r = e(OF - BF)$$

$$= e(k - r\cos\theta)$$

$$= e(k + r\cos\theta)$$

$$r - er\cos\theta = ke$$

$$r(1 - e\cos\theta) = ke$$

$$r = \frac{ke}{1 - e\cos\theta}$$

Parabola: $e = 1$, $r = \frac{k}{1 - \cos\theta}$


Ellipse: $0 < e < 1$, take $e = \frac{1}{2}$, $r = \frac{k/2}{1 - \frac{\cos\theta}{2}} = \frac{k}{2 + \cos\theta}$

Hyperbola: $e > 1$

take $e = 2$

$$r = \frac{2k}{1 - 2\cos\theta}$$

Cylindrical Coordinates



3-D
Space

$P(r, \theta, z)$
Coordinate point

$x = x_0 \Rightarrow$ It is a plane } 3-D
through x -axis




$x = x_0 \Rightarrow$ a straight line } 2-D
(vertical)

$x = x_0 \Rightarrow$ This plane can be extended infinitely in the direction of y -axis & in the direction of z -axis.

Similarly $y = y_0 \Rightarrow$ A plane through y -axis \rightarrow 3-D
 \Rightarrow A horizontal line \rightarrow 2-D


$y = y_0 \Rightarrow$ This plane can be extended infinitely in the direction of x -axis & in the direction of z -axis.

$Z = z_0 \Rightarrow$ A plane parallel to the horizon
or parallel to xy -plane.




2-D (Circle)


$\theta \in (0, 2\pi)$
full circle



3-D (Cylinder)

A right circular cylinder is a cylinder that has a closed circular surface having two parallel bases on both ends and whose elements are perpendicular \perp to its base. It is also called a right cylinder.





All the points in the green line will correspond to $\theta = \theta_0$, $r = r_0$.
 Similarly if we introduce the height $z = z_0$, it will correspond to the green line as well.


Rectangular to Cylindrical

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ z &= z \end{aligned}$$

Cylindrical to Rectangular

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ \tan \theta &= \frac{y}{x} \\ z &= z \end{aligned}$$

Spherical Coordinate



Eqn of sphere, center $(0,0,0)$
 $[x^2 + y^2 + z^2 = \rho^2]$

(i) $\rho \rightarrow \text{rho}$

$\rho = |\vec{OP}| \rightarrow \text{radius of sphere}$

$\rho \geq 0$

(ii) $\theta \rightarrow \text{angle from } x \text{ axis to the projection of the point } P \text{ on } xy\text{-plane}$

$\rightarrow \text{Vertical projection of } P \text{ on } xy\text{-plane}$

(iii) $\rightarrow \text{Let's call the projecting point on } xy\text{-plane } P' \text{ from the point } P.$


$P' \rightarrow (r, \theta) \quad r = d\{(0,0), P'\}$


$\theta \in (0, 2\pi)$

(iv) $\phi \rightarrow \text{angle between } OP \text{ to } z\text{-axis. It starts to rotate from } z\text{-axis towards } xy \text{ plane. } \therefore \phi \in (0, \pi/2)$

Then it rotates from $xy\text{-plane}$ towards the $-ve$ side of $z\text{-axis}$. $\therefore \phi \in (\pi/2, \pi)$

Hence the complete turn of ϕ is $(0, \pi)$.





$z \perp$ on xy -plane
 so z axis to the line is 90° .

$$r = \rho \sin \phi \quad \begin{cases} x = r \cos \theta = \rho \sin \phi \cos \theta \\ y = r \sin \theta = \rho \sin \phi \sin \theta \end{cases}$$

$$z = \rho \cos \phi$$

Rectangular to Spherical:

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi$$

Spherical to Rectangular

$$\rho^2 = r^2 + z^2$$

$$\rho^2 = x^2 + y^2 + z^2$$

$$\tan \theta = \frac{y}{x}$$

$$\cos \phi = \frac{z}{\rho} = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$