


A.


PCA – Full Cohort
PC2 (4.7%)
PC1 (6.1%)
Female Amyloid+
Female Amyloid−
Male Amyloid+
Male Amyloid-


B.


PLS–DA – Full Cohort
Component 2 (3.9%)
Component 1 (3.1%)
Female Amyloid+
Female Amyloid−
Male Amyloid+
Male Amyloid-


C.


PLS–DA – Female
Component 2 (3.9%)
Component 1 (3.7%)
Positive
Negative


D.


PLS–DA – Male
Component 2 (5.4%)
Component 1 (3.9%)
Positive
Negative


E.


Loading on Component 1 (Female)
Positive = elevated in amyloid+; Negative = elevated in amyloid−
threo−1−(p−Hydroxyphenyl)propylene glycol 4'−glucoside.8
MRC1
mupirocin H
Chrycolide
5−dehydro−4−deoxy−2−O−sulfo−D−glucuronate
SRGN
histidine methyl ester
1−deoxyxylonojirimycin
6−methylhept−4−enoate.1
N−methyl−L−glutamic Acid.1
Methylthioglycolate
Mecamylamine
PON1
D−galactopyranosyl trimethylammonium bromide
N−acetyl−L−2−aminoadipate(2−).1
NID2
Quinic acid
Mannosamine
3−Isopropyl−2−methoxy−5−methylpyrazine
DG(22:5/22:6/0:0)


F.


Loading on Component 1 (Male)
Positive = elevated in amyloid+; Negative = elevated in amyloid−
Pyrroline hydroxycarboxylic acid
OMD
Citroside A.4
CTSS
N2−(gamma−Glutamyl)−4−carboxyphenylhydrazine
Cys/Asp/Asp/Thr/Thr.1
Stoloniferone F
PKHD1L1
1−Aminocyclopropanecarboxylic acid
Hydroxyoctanoic acid
N−hydroxyisoleucine.1
LPL
LPA
7beta−acetoxy−gorgostan−3beta_5alpha_6beta−triol
6beta−acetoxy−24−methylcholestan−3beta_5alpha_22R_24−tetrol
TNXB
epithio−seco−cholestadienetriol S_S−dioxide
cyclopentanehexanoic acid
Methylimidazole acetaldehyde
threo−1−(p−Hydroxyphenyl)propylene glycol 4'−glucoside.8


G.


Full Cohort
Pyrroline hydroxycarboxylic acid
{[−3−phenylprop−2−en−1−yl]oxy}sulfonic acid
Methylthioglycolate
Etodolac acyl glucuronide
Asn/Met/Asp/Gln
2_4−Dihydroxyacetophenone 5−sulfate
5−(3_4−dihydroxyphenyl)−4−(sulfooxy)pentanoic acid
IGKV3D−15
KRT1
KRT9
−Log2(P−Value)
Log2 Fold Change

H.


Female Specific
threo−1−(p−Hydroxyphenyl)propylene glycol 4'−glucoside.8
DG(22:5/22:6/0:0)
Quinic acid
Methylthioglycolate
2_5−Dihydroxypyridine
3−hydroxypyridine
IGHV3−64
KRT1
KRT9
D−desosamine.1
−Log2(P−Value)
Log2 Fold Change

I.


Male Specific
Pyrroline hydroxycarboxylic acid
N−hydroxyisoleucine.1
{[−3−phenylprop−2−en−1−yl]oxy}sulfonic acid
LPA
PM20D1
4−Butyl−5−ethylthiazole.1
IGKV1
CFHR2
2_4−Dihydroxyacetophenone 5−sulfate
4−Butyl−5−ethylthiazole
−Log2(P−Value)
Log2 Fold Change

J.

Intersection size
299
261
95
66
57
20
12
3
Female (Volcano)
Male (Volcano)
Female (PLS−DA)
Male (PLS−DA)
Set size