
Loading on Component 1 (Female)
Positive = elevated in amyloid+; Negative = elevated in amyloid−
threo−1−(p−Hydroxyphenyl)propylene glycol 4'−glucoside.8
MRC1
mupirocin H
Chrycolide
5−dehydro−4−deoxy−2−O−sulfo−D−glucuronate
SRGN
histidine methyl ester
1−deoxyxylonojirimycin
6−methylhept−4−enoate.1
N−methyl−L−glutamic Acid.1
Methylthioglycolate
Mecamylamine
PON1
D−galactopyranosyl trimethylammonium bromide
N−acetyl−L−2−aminoadipate(2−).1
NID2
Quinic acid
Mannosamine
3−Isopropyl−2−methoxy−5−methylpyrazine
DG(22:5/22:6/0:0)
−0.05
0.00
0.05
0.10
Loading on Component 1 (Male)
Positive = elevated in amyloid+; Negative = elevated in amyloid−
Pyrroline hydroxycarboxylic acid
OMD
Citroside A.4
CTSS
N2−(gamma−Glutamyl)−4−carboxyphenylhydrazine
Cys/Asp/Asp/Thr/Thr.1
Stoloniferone F
PKHD1L1
1−Aminocyclopropanecarboxylic acid
Hydroxyoctanoic acid
N−hydroxyisoleucine.1
LPL
LPA
7beta−acetoxy−gorgostan−3beta_5alpha_6beta−triol
6beta−acetoxy−24−methylcholestan−3beta_5alpha_22R_24−tetrol
TNXB
epithio−seco−cholestadienetriol S_S−dioxide
cyclopentanehexanoic acid
Methylimidazole acetaldehyde
threo−1−(p−Hydroxyphenyl)propylene glycol 4'−glucoside.8
−0.05
0.00
0.05
0.10
Association
Higher in Amyloid−
Higher in Amyloid+
Higher in Amyloid−
Higher in Amyloid+