
De novo DNA contigs; full cleaned reads; taxonomy assignments require validation
Mean retained depth + 0.001 (log scale)
1e+02
1e+00
1e−02
0.00
0.25
0.50
0.75
1.00
GC fraction (ACGT denominator)
Contig length (bp)
50000
100000
Kraken contig assignment
Acinetobacter
Actinomycetes
Bacteria
Bradyrhizobium
Bremia lactucae
Brevundimonas sp. S1H14
Burkholderia vietnamiensis
cellular organisms
Comamonadaceae
Corynebacterium tuberculostearicum
Cutibacterium acnes
Delftia acidovorans
Dietzia timorensis
Enterobacteriaceae
Escherichia coli
Eukaryota
Exiguobacterium alkaliphilum
Gammaproteobacteria
Homo sapiens
Kocuria
Kocuria palustris
Kocuria rhizophila
Kocuria varians
Leishmania
Leucobacter
Leucobacter aridicollis
Microbacteriaceae
Micrococcus
Micrococcus luteus
Opisthokonta
Pseudomonadati
Pseudomonadota
root
Sar
Sphingobium sp. WTD−1
Sphingomonadales
Staphylococcus
Stenotrophomonas
Streptomyces
Stutzerimonas kunmingensis
Stutzerimonas stutzeri
Stutzerimonas stutzeri group
Tequatrovirus
unclassified
unclassified Stenotrophomonas
Xanthomonas campestris pv. incanae