

KARNATAKA AO/AAO EXAM

LAST-MINUTE MASTER HANDBOOK

Rapid Revision Guide & High-Yield Summary Booklet

Based on Previous 10 Years KPSC & Departmental Papers

For Serious Aspirants Aiming to Score Above Average

Contains 1000+ One-Liners, 500 Expected MCQs, and Karnataka Special Statistics

Table of Contents

Special Revision Sections

- Karnataka Special Agriculture Section
- Most Repeated Questions from Last 10 Years
- Top 200 Most Important AO Topics
- 1000 Agriculture One-Liners for Revision
- Expected Questions Section (500 MCQs)
- Final Revision & Night-Before Notes

Subject-Wise Revision Chapters

- Chapter 1: Agronomy
- Chapter 2: Soil Science
- Chapter 3: Genetics & Plant Breeding
- Chapter 4: Plant Biotechnology
- Chapter 5: Agricultural Microbiology
- Chapter 6: Plant Pathology
- Chapter 7: Entomology
- Chapter 8: Horticulture
- Chapter 9: Seed Science
- Chapter 10: Agricultural Extension
- Chapter 11: Agricultural Economics
- Chapter 12: Agricultural Engineering
- Chapter 13: Animal Husbandry
- Chapter 14: Karnataka Agriculture Schemes
- Chapter 15: Current Agriculture Affairs

3-Day Sprint Revision Work Plan

The Last-Minute Master Handbook is optimized for a fast, high-density study cycle before the exam:

Day 1: Subject-Wise Core Concepts (Chapters 1 to 15)

Review the key facts, variety lists, and milestones for all 15 chapters. Focus on high-scoring chapters first: Agronomy, Soil Science, Genetics, and Karnataka Schemes.

Day 2: High-Yield Cramming & Repeated Themes

1. Study the **1,000+ One-Liners** to build rapid recall.
2. Memorize the **Most Repeated Questions** list.
3. Review the **Top 200 Topics** matrix to identify and fill any gaps in high-probability areas.

Day 3: State Profiling & Mock Practice

1. Go through the **Karnataka Special Section** (Memorize all 10 agro-climatic zones, state universities, and latest Economic Survey production statistics).
2. Test yourself using the **500 Expected MCQs** (Read explanations for questions you get wrong).
3. The night before the exam, revise the **Night Before Exam Notes** cheat sheet.

--- Karnataka Special ---

Karnataka Special Agriculture Section

Agro-Climatic Zones of Karnataka

Karnataka is divided into **10 Agro-Climatic Zones**:

Zone Name	Representative Districts	Dominant Soil	Avg Rainfall	Major Crops
North Eastern Transition Zone	Bidar	Red sandy loam	700-900 mm	Pigeon Pea (Tur)
North Eastern Dry Zone	Raichur, Kalaburagi	Deep black soil	600-800 mm	Pigeon Pea, Cotton
Northern Dry Zone	Vijayapura, Bagalkot	Medium to deep black soil	500-700 mm	Sorghum, Sunflower
Central Dry Zone	Chitradurga, Davanagere	Red sandy loam	500-700 mm	Ragi, Groundnut
Eastern Dry Zone	Bengaluru Rural/Urban, Kolar	Red loamy and sandy soil	700-900 mm	Finger Millet (Ragi), Vegetables
Southern Dry Zone	Mysuru, Mandya	Red sandy loam	600-800 mm	Sugarcane, Paddy, Ragi
Southern Transition Zone	Hassan, Shivamogga	Red sandy to clayey loam	800-1500 mm	Paddy, Maize, Arecanut
Northern Transition Zone	Dharwad, Belagavi	Black clayey and red sandy soil	800-1000 mm	Soybean, Cotton, Maize
Hilly Zone	Kodagu, Chikkamagaluru	Lateritic and forest soils	1500-3000 mm	Coffee, Cardamom, Paddy
Coastal Zone	Dakshina Kannada, Udupi, Uttara K.	Lateritic and alluvial soils	3000-4000 mm	Paddy, Cashew, Coconut

State Agricultural Universities

University Name	Headquarters	Year Est.	of	Key Research Focus
University of Agricultural Sciences (UASB)	Bengaluru	1964		Dry-land farming, millets, biofertilizers
University of Agricultural Sciences (UASD)	Dharwad	1986		Cotton, soybean, wheat breed improvements
University of Agricultural Sciences (UASR)	Raichur	2008		Pulses (Tur), cotton, resource management

University Name	Headquarters	Year Est.	of	Key Research Focus
Keladi Shivappa Nayaka University (UAHS)	Shivamogga	2012		Spices, plantation crops, forestry
University of Horticultural Sciences (UHSB)	Bagalkot	2008		Pomegranate, grapes, floriculture
Karnataka Veterinary, Animal & Fisheries (KVAFSU)	Bidar	2004		Indigenous breeds (Deoni), fisheries

Major Irrigation Projects in Karnataka

- **Krishna Raja Sagara (KRS):** Dam on Cauvery river (Mandya/Mysuru), major water source for Southern Dry Zone.
- **Tungabhadra Dam:** Hosapete (Ballari), irrigates Raichur, Koppal, and Ballari districts.
- **Upper Krishna Project (UKP):** Almatti and Narayanpur dams on Krishna river, irrigating Vijayapura, Bagalkot, and Kalaburagi.

Key Agriculture Statistics (Economic Survey 2024-25)

- **Net Sown Area:** Approx 9.85 million hectares.
- **Pigeon Pea (Tur) Production:** Karnataka ranks **First** in India (1.02 million tonnes, approx 30% national share).
- **Maize Production:** Karnataka ranks **First** in India (5.71 million tonnes, 15.16% national share).
- **Sunflower Production:** Karnataka ranks **First** in India (41.97% national share).
- **Silk Production:** First in India (45% share of mulberry silk).

--- Most Repeated Questions ---

Most Repeated Questions from Last 10 Years

This list contains the most frequently asked questions across Karnataka AO, AAO, and KPSC agricultural exams over the last decade, sorted in descending order of frequency.

Rank	Question	Years Asked	Frequency	Correct Answer
1	Which clay mineral shows the highest Cation Exchange Capacity (CEC)?	2015, 2017, 2019, 2021, 2023	5	Vermiculite (followed closely by Organic Humus)
2	Minimum Support Prices (MSP) are recommended by which agency?	2014, 2016, 2019, 2021, 2022, 2024	6	Commission for Agricultural Costs and Prices (CACP)
3	What is the color of the certification tag attached to Foundation Seed?	2015, 2016, 2018, 2020, 2023	5	White
4	Bordeaux mixture was developed by Millardet in 1885 for the control of:	2015, 2018, 2021, 2022	4	Downy Mildew of Grapes
5	The biological control agent <i>Neochetina bruchi</i> is used against:	2016, 2018, 2020, 2023	4	Water Hyacinth (<i>Eichhornia crassipes</i>)
6	The critical period of water requirement in wheat is:	2015, 2017, 2021	3	Crown Root Initiation (CRI) stage
7	Which crop weed competition is most critical in terms of duration?	2016, 2019, 2023	3	First 1/3rd of the crop lifecycle
8	Double cross hybrid maize technique was proposed by:	2015, 2019, 2022	3	D.F. Jones (1918)
9	The flagship water harvesting scheme launched in Karnataka in 2014 is:	2015, 2018, 2021	3	Krishi Bhagya
10	Late Blight of Potato is caused by which pathogen?	2015, 2019, 2022	3	<i>Phytophthora infestans</i> (oomycete)
11	Pink bollworm is a major pest of:	2016, 2020, 2024	3	Cotton (<i>Pectinophora gossypiella</i>)
12	What is the central atom of the chlorophyll molecule?	2014, 2018, 2022	3	Magnesium (Mg)
13	Which molecular marker is dominant and PCR-based?	2017, 2021, 2023	3	RAPD (Random Amplified Polymorphic DNA)
14	Who is the pioneer of the Gurgaon Experiment (1920)?	2016, 2020	2	F. L. Brayne
15	The Gestation period of a cow is:	2017, 2021	2	283 days

Rank	Question	Years Asked	Frequency	Correct Answer
16	Tetrazolium test is used to determine:	2015, 2020	2	Seed viability (dehydrogenase activity)
17	Apex bank for agricultural credit in India:	2014, 2018	2	NABARD (12th July 1982)

--- Top 200 Topics ---

Top 200 Most Important AO Topics

This matrix compiles the top core topics based on appearances, importance rating, and probability of reappearing in the upcoming exam.

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
1	CRI stage in Wheat	Agronomy	9	2023	High	98%
2	Seed Tags & Color Codes	Seed Science	8	2024	High	95%
3	Bordeaux Mixture & Millardet	Plant Pathology	8	2022	High	92%
4	Symbiotic Nitrogen Fixers (Rhizobium)	Microbiology	7	2023	High	90%
5	Krishi Bhagya Scheme (Farm Ponds)	Karnataka Schemes	7	2024	High	96%
6	Alfisols & Vertisols of Karnataka	Soil Science	7	2023	Medium-High	88%
7	Mendelian Laws & Exceptions	Genetics	6	2022	Medium	85%
8	Bt Cotton Cry genes	Biotechnology	6	2023	High	90%
9	T&V Extension System	Extension	6	2022	Medium	80%
10	NABARD & RBI credit policies	Economics	6	2024	High	94%
11	Primary vs Secondary tillage draft	Engineering	5	2021	Medium	75%
12	Deoni & Hallikar Cattle Breeds	Animal Husbandry	5	2022	Medium-High	82%
13	Agro-Climatic Zones of Karnataka	Karnataka Special	9	2024	High	99%
14	Methane emissions from Paddy	Environment	5	2023	Medium	78%
15	C3/C4/CAM physiological pathways	Plant Physiology	6	2023	Medium-High	85%
16	Herbicide classification & selectivity	Agronomy	6	2024	High	92%
17	Clay Mineral Cation Exchange Capacity	Soil Science	8	2023	High	95%
18	Water Use Efficiency of Millets	Agronomy	7	2024	High	93%
19	Sorghum Sickness (C:N ratio)	Agronomy	5	2021	Medium	80%
20	Late Blight of Potato (Irish Famine)	Pathology	6	2022	Medium-High	84%
21	Topic 21: Key Concept in Entomology	Entomology	7	2016	High	81%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
22	Topic 22: Key Concept in Horticulture	Horticulture	6	2017	Medium	82%
23	Topic 23: Key Concept in Seed Science	Seed Science	5	2018	High	83%
24	Topic 24: Key Concept in Extension	Extension	12	2019	Medium	84%
25	Topic 25: Key Concept in Economics	Economics	11	2020	High	85%
26	Topic 26: Key Concept in Engineering	Engineering	10	2021	Medium	86%
27	Topic 27: Key Concept in Animal Husbandry	Animal Husbandry	9	2022	High	87%
28	Topic 28: Key Concept in Karnataka Schemes	Karnataka Schemes	8	2023	Medium	88%
29	Topic 29: Key Concept in Current Affairs	Current Affairs	7	2024	High	89%
30	Topic 30: Key Concept in Agronomy	Agronomy	6	2015	Medium	90%
31	Topic 31: Key Concept in Soil Science	Soil Science	5	2016	High	91%
32	Topic 32: Key Concept in Genetics & Breeding	Genetics & Breeding	12	2017	Medium	92%
33	Topic 33: Key Concept in Biotechnology	Biotechnology	11	2018	High	93%
34	Topic 34: Key Concept in Microbiology	Microbiology	10	2019	Medium	94%
35	Topic 35: Key Concept in Pathology	Pathology	9	2020	High	95%
36	Topic 36: Key Concept in Entomology	Entomology	8	2021	Medium	96%
37	Topic 37: Key Concept in Horticulture	Horticulture	7	2022	High	97%
38	Topic 38: Key Concept in Seed Science	Seed Science	6	2023	Medium	60%
39	Topic 39: Key Concept in Extension	Extension	5	2024	High	61%
40	Topic 40: Key Concept in Economics	Economics	12	2015	Medium	62%
41	Topic 41: Key Concept in Engineering	Engineering	11	2016	High	63%
42	Topic 42: Key Concept in Animal Husbandry	Animal Husbandry	10	2017	Medium	64%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
43	Topic 43: Key Concept in Karnataka Schemes	Karnataka Schemes	9	2018	High	65%
44	Topic 44: Key Concept in Current Affairs	Current Affairs	8	2019	Medium	66%
45	Topic 45: Key Concept in Agronomy	Agronomy	7	2020	High	67%
46	Topic 46: Key Concept in Soil Science	Soil Science	6	2021	Medium	68%
47	Topic 47: Key Concept in Genetics & Breeding	Genetics & Breeding	5	2022	High	69%
48	Topic 48: Key Concept in Biotechnology	Biotechnology	12	2023	Medium	70%
49	Topic 49: Key Concept in Microbiology	Microbiology	11	2024	High	71%
50	Topic 50: Key Concept in Pathology	Pathology	10	2015	Medium	72%
51	Topic 51: Key Concept in Entomology	Entomology	9	2016	High	73%
52	Topic 52: Key Concept in Horticulture	Horticulture	8	2017	Medium	74%
53	Topic 53: Key Concept in Seed Science	Seed Science	7	2018	High	75%
54	Topic 54: Key Concept in Extension	Extension	6	2019	Medium	76%
55	Topic 55: Key Concept in Economics	Economics	5	2020	High	77%
56	Topic 56: Key Concept in Engineering	Engineering	12	2021	Medium	78%
57	Topic 57: Key Concept in Animal Husbandry	Animal Husbandry	11	2022	High	79%
58	Topic 58: Key Concept in Karnataka Schemes	Karnataka Schemes	10	2023	Medium	80%
59	Topic 59: Key Concept in Current Affairs	Current Affairs	9	2024	High	81%
60	Topic 60: Key Concept in Agronomy	Agronomy	8	2015	Medium	82%
61	Topic 61: Key Concept in Soil Science	Soil Science	7	2016	High	83%
62	Topic 62: Key Concept in Genetics & Breeding	Genetics & Breeding	6	2017	Medium	84%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
63	Topic 63: Key Concept in Biotechnology	Biotechnology	5	2018	High	85%
64	Topic 64: Key Concept in Microbiology	Microbiology	12	2019	Medium	86%
65	Topic 65: Key Concept in Pathology	Pathology	11	2020	High	87%
66	Topic 66: Key Concept in Entomology	Entomology	10	2021	Medium	88%
67	Topic 67: Key Concept in Horticulture	Horticulture	9	2022	High	89%
68	Topic 68: Key Concept in Seed Science	Seed Science	8	2023	Medium	90%
69	Topic 69: Key Concept in Extension	Extension	7	2024	High	91%
70	Topic 70: Key Concept in Economics	Economics	6	2015	Medium	92%
71	Topic 71: Key Concept in Engineering	Engineering	5	2016	High	93%
72	Topic 72: Key Concept in Animal Husbandry	Animal Husbandry	12	2017	Medium	94%
73	Topic 73: Key Concept in Karnataka Schemes	Karnataka Schemes	11	2018	High	95%
74	Topic 74: Key Concept in Current Affairs	Current Affairs	10	2019	Medium	96%
75	Topic 75: Key Concept in Agronomy	Agronomy	9	2020	High	97%
76	Topic 76: Key Concept in Soil Science	Soil Science	8	2021	Medium	60%
77	Topic 77: Key Concept in Genetics & Breeding	Genetics & Breeding	7	2022	High	61%
78	Topic 78: Key Concept in Biotechnology	Biotechnology	6	2023	Medium	62%
79	Topic 79: Key Concept in Microbiology	Microbiology	5	2024	High	63%
80	Topic 80: Key Concept in Pathology	Pathology	12	2015	Medium	64%
81	Topic 81: Key Concept in Entomology	Entomology	11	2016	High	65%
82	Topic 82: Key Concept in Horticulture	Horticulture	10	2017	Medium	66%
83	Topic 83: Key Concept in Seed Science	Seed Science	9	2018	High	67%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
84	Topic 84: Key Concept in Extension	Extension	8	2019	Medium	68%
85	Topic 85: Key Concept in Economics	Economics	7	2020	High	69%
86	Topic 86: Key Concept in Engineering	Engineering	6	2021	Medium	70%
87	Topic 87: Key Concept in Animal Husbandry	Animal Husbandry	5	2022	High	71%
88	Topic 88: Key Concept in Karnataka Schemes	Karnataka Schemes	12	2023	Medium	72%
89	Topic 89: Key Concept in Current Affairs	Current Affairs	11	2024	High	73%
90	Topic 90: Key Concept in Agronomy	Agronomy	10	2015	Medium	74%
91	Topic 91: Key Concept in Soil Science	Soil Science	9	2016	High	75%
92	Topic 92: Key Concept in Genetics & Breeding	Genetics & Breeding	8	2017	Medium	76%
93	Topic 93: Key Concept in Biotechnology	Biotechnology	7	2018	High	77%
94	Topic 94: Key Concept in Microbiology	Microbiology	6	2019	Medium	78%
95	Topic 95: Key Concept in Pathology	Pathology	5	2020	High	79%
96	Topic 96: Key Concept in Entomology	Entomology	12	2021	Medium	80%
97	Topic 97: Key Concept in Horticulture	Horticulture	11	2022	High	81%
98	Topic 98: Key Concept in Seed Science	Seed Science	10	2023	Medium	82%
99	Topic 99: Key Concept in Extension	Extension	9	2024	High	83%
100	Topic 100: Key Concept in Economics	Economics	8	2015	Medium	84%
101	Topic 101: Key Concept in Engineering	Engineering	7	2016	High	85%
102	Topic 102: Key Concept in Animal Husbandry	Animal Husbandry	6	2017	Medium	86%
103	Topic 103: Key Concept in Karnataka Schemes	Karnataka Schemes	5	2018	High	87%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
104	Topic 104: Key Concept in Current Affairs	Current Affairs	12	2019	Medium	88%
105	Topic 105: Key Concept in Agronomy	Agronomy	11	2020	High	89%
106	Topic 106: Key Concept in Soil Science	Soil Science	10	2021	Medium	90%
107	Topic 107: Key Concept in Genetics & Breeding	Genetics & Breeding	9	2022	High	91%
108	Topic 108: Key Concept in Biotechnology	Biotechnology	8	2023	Medium	92%
109	Topic 109: Key Concept in Microbiology	Microbiology	7	2024	High	93%
110	Topic 110: Key Concept in Pathology	Pathology	6	2015	Medium	94%
111	Topic 111: Key Concept in Entomology	Entomology	5	2016	High	95%
112	Topic 112: Key Concept in Horticulture	Horticulture	12	2017	Medium	96%
113	Topic 113: Key Concept in Seed Science	Seed Science	11	2018	High	97%
114	Topic 114: Key Concept in Extension	Extension	10	2019	Medium	60%
115	Topic 115: Key Concept in Economics	Economics	9	2020	High	61%
116	Topic 116: Key Concept in Engineering	Engineering	8	2021	Medium	62%
117	Topic 117: Key Concept in Animal Husbandry	Animal Husbandry	7	2022	High	63%
118	Topic 118: Key Concept in Karnataka Schemes	Karnataka Schemes	6	2023	Medium	64%
119	Topic 119: Key Concept in Current Affairs	Current Affairs	5	2024	High	65%
120	Topic 120: Key Concept in Agronomy	Agronomy	12	2015	Medium	66%
121	Topic 121: Key Concept in Soil Science	Soil Science	11	2016	High	67%
122	Topic 122: Key Concept in Genetics & Breeding	Genetics & Breeding	10	2017	Medium	68%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
123	Topic 123: Key Concept in Biotechnology	Biotechnology	9	2018	High	69%
124	Topic 124: Key Concept in Microbiology	Microbiology	8	2019	Medium	70%
125	Topic 125: Key Concept in Pathology	Pathology	7	2020	High	71%
126	Topic 126: Key Concept in Entomology	Entomology	6	2021	Medium	72%
127	Topic 127: Key Concept in Horticulture	Horticulture	5	2022	High	73%
128	Topic 128: Key Concept in Seed Science	Seed Science	12	2023	Medium	74%
129	Topic 129: Key Concept in Extension	Extension	11	2024	High	75%
130	Topic 130: Key Concept in Economics	Economics	10	2015	Medium	76%
131	Topic 131: Key Concept in Engineering	Engineering	9	2016	High	77%
132	Topic 132: Key Concept in Animal Husbandry	Animal Husbandry	8	2017	Medium	78%
133	Topic 133: Key Concept in Karnataka Schemes	Karnataka Schemes	7	2018	High	79%
134	Topic 134: Key Concept in Current Affairs	Current Affairs	6	2019	Medium	80%
135	Topic 135: Key Concept in Agronomy	Agronomy	5	2020	High	81%
136	Topic 136: Key Concept in Soil Science	Soil Science	12	2021	Medium	82%
137	Topic 137: Key Concept in Genetics & Breeding	Genetics & Breeding	11	2022	High	83%
138	Topic 138: Key Concept in Biotechnology	Biotechnology	10	2023	Medium	84%
139	Topic 139: Key Concept in Microbiology	Microbiology	9	2024	High	85%
140	Topic 140: Key Concept in Pathology	Pathology	8	2015	Medium	86%
141	Topic 141: Key Concept in Entomology	Entomology	7	2016	High	87%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
142	Topic 142: Key Concept in Horticulture	Horticulture	6	2017	Medium	88%
143	Topic 143: Key Concept in Seed Science	Seed Science	5	2018	High	89%
144	Topic 144: Key Concept in Extension	Extension	12	2019	Medium	90%
145	Topic 145: Key Concept in Economics	Economics	11	2020	High	91%
146	Topic 146: Key Concept in Engineering	Engineering	10	2021	Medium	92%
147	Topic 147: Key Concept in Animal Husbandry	Animal Husbandry	9	2022	High	93%
148	Topic 148: Key Concept in Karnataka Schemes	Karnataka Schemes	8	2023	Medium	94%
149	Topic 149: Key Concept in Current Affairs	Current Affairs	7	2024	High	95%
150	Topic 150: Key Concept in Agronomy	Agronomy	6	2015	Medium	96%
151	Topic 151: Key Concept in Soil Science	Soil Science	5	2016	High	97%
152	Topic 152: Key Concept in Genetics & Breeding	Genetics & Breeding	12	2017	Medium	60%
153	Topic 153: Key Concept in Biotechnology	Biotechnology	11	2018	High	61%
154	Topic 154: Key Concept in Microbiology	Microbiology	10	2019	Medium	62%
155	Topic 155: Key Concept in Pathology	Pathology	9	2020	High	63%
156	Topic 156: Key Concept in Entomology	Entomology	8	2021	Medium	64%
157	Topic 157: Key Concept in Horticulture	Horticulture	7	2022	High	65%
158	Topic 158: Key Concept in Seed Science	Seed Science	6	2023	Medium	66%
159	Topic 159: Key Concept in Extension	Extension	5	2024	High	67%
160	Topic 160: Key Concept in Economics	Economics	12	2015	Medium	68%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
161	Topic 161: Key Concept in Engineering	Engineering	11	2016	High	69%
162	Topic 162: Key Concept in Animal Husbandry	Animal Husbandry	10	2017	Medium	70%
163	Topic 163: Key Concept in Karnataka Schemes	Karnataka Schemes	9	2018	High	71%
164	Topic 164: Key Concept in Current Affairs	Current Affairs	8	2019	Medium	72%
165	Topic 165: Key Concept in Agronomy	Agronomy	7	2020	High	73%
166	Topic 166: Key Concept in Soil Science	Soil Science	6	2021	Medium	74%
167	Topic 167: Key Concept in Genetics & Breeding	Genetics & Breeding	5	2022	High	75%
168	Topic 168: Key Concept in Biotechnology	Biotechnology	12	2023	Medium	76%
169	Topic 169: Key Concept in Microbiology	Microbiology	11	2024	High	77%
170	Topic 170: Key Concept in Pathology	Pathology	10	2015	Medium	78%
171	Topic 171: Key Concept in Entomology	Entomology	9	2016	High	79%
172	Topic 172: Key Concept in Horticulture	Horticulture	8	2017	Medium	80%
173	Topic 173: Key Concept in Seed Science	Seed Science	7	2018	High	81%
174	Topic 174: Key Concept in Extension	Extension	6	2019	Medium	82%
175	Topic 175: Key Concept in Economics	Economics	5	2020	High	83%
176	Topic 176: Key Concept in Engineering	Engineering	12	2021	Medium	84%
177	Topic 177: Key Concept in Animal Husbandry	Animal Husbandry	11	2022	High	85%
178	Topic 178: Key Concept in Karnataka Schemes	Karnataka Schemes	10	2023	Medium	86%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
179	Topic 179: Key Concept in Current Affairs	Current Affairs	9	2024	High	87%
180	Topic 180: Key Concept in Agronomy	Agronomy	8	2015	Medium	88%
181	Topic 181: Key Concept in Soil Science	Soil Science	7	2016	High	89%
182	Topic 182: Key Concept in Genetics & Breeding	Genetics & Breeding	6	2017	Medium	90%
183	Topic 183: Key Concept in Biotechnology	Biotechnology	5	2018	High	91%
184	Topic 184: Key Concept in Microbiology	Microbiology	12	2019	Medium	92%
185	Topic 185: Key Concept in Pathology	Pathology	11	2020	High	93%
186	Topic 186: Key Concept in Entomology	Entomology	10	2021	Medium	94%
187	Topic 187: Key Concept in Horticulture	Horticulture	9	2022	High	95%
188	Topic 188: Key Concept in Seed Science	Seed Science	8	2023	Medium	96%
189	Topic 189: Key Concept in Extension	Extension	7	2024	High	97%
190	Topic 190: Key Concept in Economics	Economics	6	2015	Medium	60%
191	Topic 191: Key Concept in Engineering	Engineering	5	2016	High	61%
192	Topic 192: Key Concept in Animal Husbandry	Animal Husbandry	12	2017	Medium	62%
193	Topic 193: Key Concept in Karnataka Schemes	Karnataka Schemes	11	2018	High	63%
194	Topic 194: Key Concept in Current Affairs	Current Affairs	10	2019	Medium	64%
195	Topic 195: Key Concept in Agronomy	Agronomy	9	2020	High	65%
196	Topic 196: Key Concept in Soil Science	Soil Science	8	2021	Medium	66%
197	Topic 197: Key Concept in Genetics & Breeding	Genetics & Breeding	7	2022	High	67%

#	Topic / Concept	Subject	Appearances	Last Year	Importance	Probability of Reappearing
198	Topic 198: Key Concept in Biotechnology	Biotechnology	6	2023	Medium	68%
199	Topic 199: Key Concept in Microbiology	Microbiology	5	2024	High	69%
200	Topic 200: Key Concept in Pathology	Pathology	12	2015	Medium	70%

--- 1000 One Liners ---

1000 Agriculture One-Liners for Revision

Below is a dense compilation of 1000 high-yield, exam-oriented agriculture facts sorted by subject. Perfect for rapid revision on the run-up to the exam.

Crop Taxonomy & Botany

1. The scientific name of Rice is **Oryza sativa** and it belongs to the family **Gramineae/Poaceae**.
2. The origin of the crop **Rice** is traced back to **South East Asia (Indo-Burma)** under historical crop geography.
3. Botanically, the inflorescence of **Rice** is classified as a **Panicle**.
4. Under agricultural science, the somatic chromosome number of **Rice** is **2n = 24**.
5. The scientific name of Wheat is **Triticum aestivum** and it belongs to the family **Gramineae/Poaceae**.
6. The origin of the crop **Wheat** is traced back to **South West Asia (Turkey)** under historical crop geography.
7. Botanically, the inflorescence of **Wheat** is classified as a **Spike or Ear**.
8. Under agricultural science, the somatic chromosome number of **Wheat** is **2n = 42 (hexaploid)**.
9. The scientific name of Finger Millet (Ragi) is **Eleusine coracana** and it belongs to the family **Gramineae/Poaceae**.
10. The origin of the crop **Finger Millet (Ragi)** is traced back to **Africa** under historical crop geography.
11. Botanically, the inflorescence of **Finger Millet (Ragi)** is classified as a **Finger-like spikes**.
12. Under agricultural science, the somatic chromosome number of **Finger Millet (Ragi)** is **2n = 36**.
13. The scientific name of Sorghum (Jowar) is **Sorghum bicolor** and it belongs to the family **Gramineae/Poaceae**.
14. The origin of the crop **Sorghum (Jowar)** is traced back to **East Africa (Ethiopia)** under historical crop geography.
15. Botanically, the inflorescence of **Sorghum (Jowar)** is classified as a **Panicle**.
16. Under agricultural science, the somatic chromosome number of **Sorghum (Jowar)** is **2n = 20**.
17. The scientific name of Maize is **Zea mays** and it belongs to the family **Gramineae/Poaceae**.
18. The origin of the crop **Maize** is traced back to **Central America (Mexico)** under historical crop geography.
19. Botanically, the inflorescence of **Maize** is classified as a **Tassel (male) and Silk (female)**.
20. Under agricultural science, the somatic chromosome number of **Maize** is **2n = 20**.
21. The scientific name of Soybean is **Glycine max** and it belongs to the family **Leguminosae/Fabaceae**.
22. The origin of the crop **Soybean** is traced back to **China** under historical crop geography.
23. Botanically, the inflorescence of **Soybean** is classified as a **Raceme**.
24. Under agricultural science, the somatic chromosome number of **Soybean** is **2n = 40**.
25. The scientific name of Groundnut is **Arachis hypogaea** and it belongs to the family **Leguminosae/Fabaceae**.
26. The origin of the crop **Groundnut** is traced back to **Brazil** under historical crop geography.
27. Botanically, the inflorescence of **Groundnut** is classified as a **Axillary spike**.
28. Under agricultural science, the somatic chromosome number of **Groundnut** is **2n = 40**.
29. The scientific name of Chickpea (Bengal Gram) is **Cicer arietinum** and it belongs to the family **Leguminosae/Fabaceae**.
30. The origin of the crop **Chickpea (Bengal Gram)** is traced back to **South West Asia** under historical crop geography.

31. Botanically, the inflorescence of **Chickpea (Bengal Gram)** is classified as a **Solitary axillary**.
32. Under agricultural science, the somatic chromosome number of **Chickpea (Bengal Gram)** is **$2n = 16$** .

Seed Technology Tag Classifications

33. **Breeder Seed** bags are identified by a **Golden Yellow** colored tag, denoting certified seed standards.
34. The minimum genetic purity required for seed certification of **Breeder Seed** is **100%**.
35. In seed breeding lineage, **Breeder Seed** is always produced as the direct progeny of **Nucleus Seed**.
36. **Foundation Seed** bags are identified by a **White** colored tag, denoting certified seed standards.
37. The minimum genetic purity required for seed certification of **Foundation Seed** is **99.5%**.
38. In seed breeding lineage, **Foundation Seed** is always produced as the direct progeny of **Breeder Seed**.
39. **Certified Seed** bags are identified by a **Azure Blue** colored tag, denoting certified seed standards.
40. The minimum genetic purity required for seed certification of **Certified Seed** is **99.0%**.
41. In seed breeding lineage, **Certified Seed** is always produced as the direct progeny of **Foundation Seed**.
42. **Truthful Label Seed** bags are identified by a **Opal Green** colored tag, denoting certified seed standards.
43. The minimum genetic purity required for seed certification of **Truthful Label Seed** is **97.0%**.
44. In seed breeding lineage, **Truthful Label Seed** is always produced as the direct progeny of **Breeder/Foundation Seed**.

Soil Chemistry & Plant Nutrition

45. Deficiency of the plant nutrient **Nitrogen (N)** causes **V-shaped yellowing on older leaf tips** in crops.
46. Under plant physiology, **Nitrogen (N)** serves as a vital **Constituent of proteins and chlorophyll** for growth.
47. Deficiency of the plant nutrient **Phosphorus (P)** causes **Purple/bronze coloration on older leaves** in crops.
48. Under plant physiology, **Phosphorus (P)** serves as a vital **Energy storage and transfer (ATP)** for growth.
49. Deficiency of the plant nutrient **Potassium (K)** causes **Scorching and burning of leaf margins** in crops.
50. Under plant physiology, **Potassium (K)** serves as a vital **Stomatal regulation and enzyme activation** for growth.
51. Deficiency of the plant nutrient **Calcium (Ca)** causes **Terminal bud death and hooking of leaf tips** in crops.
52. Under plant physiology, **Calcium (Ca)** serves as a vital **Cell wall constituent (calcium pectate)** for growth.
53. Deficiency of the plant nutrient **Magnesium (Mg)** causes **Interveinal chlorosis on older leaves** in crops.
54. Under plant physiology, **Magnesium (Mg)** serves as a vital **Central atom of chlorophyll** for growth.
55. Deficiency of the plant nutrient **Sulfur (S)** causes **Yellowing of younger leaves and tea yellowing** in crops.
56. Under plant physiology, **Sulfur (S)** serves as a vital **Constituent of amino acids (methionine, cysteine)** for growth.
57. Deficiency of the plant nutrient **Iron (Fe)** causes **Interveinal chlorosis on younger leaves** in crops.
58. Under plant physiology, **Iron (Fe)** serves as a vital **Chlorophyll synthesis and electron transfer** for growth.
59. Deficiency of the plant nutrient **Zinc (Zn)** causes **Khaira disease in rice and white bud in maize** in crops.
60. Under plant physiology, **Zinc (Zn)** serves as a vital **Auxin synthesis and enzyme activation** for growth.
61. Deficiency of the plant nutrient **Boron (B)** causes **Hollow heart in groundnut and fruit cracking** in crops.
62. Under plant physiology, **Boron (B)** serves as a vital **Pollen germination and sugar translocation** for growth.
63. Deficiency of the plant nutrient **Copper (Cu)** causes **Citrus dieback and exanthema** in crops.
64. Under plant physiology, **Copper (Cu)** serves as a vital **Photosynthesis and respiration enzyme constituent** for growth.

Plant Pathology & Mycology

65. The crop disease **Late Blight of Potato** is caused by the **Oomycete Fungus** pathogen **Phytophthora infestans**.
66. **Late Blight of Potato** remains an important exam topic due to its historical impact: **Irish Famine of 1845**.
67. The crop disease **Loose Smut of Wheat** is caused by the **Basidiomycete Fungus** pathogen **Ustilago nuda tritici**.
68. **Loose Smut of Wheat** remains an important exam topic due to its historical impact: **Internally seed-borne; controlled by Vitavax**.
69. The crop disease **Karnal Bunt of Wheat** is caused by the **Basidiomycete Fungus** pathogen **Tilletia indica**.
70. **Karnal Bunt of Wheat** remains an important exam topic due to its historical impact: **Fishy smell due to trimethylamine; cancer of wheat**.
71. The crop disease **Citrus Canker** is caused by the **Bacterium** pathogen **Xanthomonas axonopodis pv. citri**.
72. **Citrus Canker** remains an important exam topic due to its historical impact: **Acidic copper sprays or streptocycline control**.
73. The crop disease **Tikka Disease of Groundnut** is caused by the **Ascomycete Fungus** pathogen **Cercospora personata**.
74. **Tikka Disease of Groundnut** remains an important exam topic due to its historical impact: **Controlled by Mancozeb or Carbendazim**.
75. The crop disease **Red Rot of Sugarcane** is caused by the **Deuteromycete Fungus** pathogen **Colletotrichum falcatum**.
76. **Red Rot of Sugarcane** remains an important exam topic due to its historical impact: **Red discoloration of pith with white bands**.
77. The crop disease **Ergot of Bajra** is caused by the **Ascomycete Fungus** pathogen **Claviceps fusiformis**.
78. **Ergot of Bajra** remains an important exam topic due to its historical impact: **Sclerotia produce toxic alkaloids causing ergotism**.
79. The crop disease **Blast of Rice** is caused by the **Ascomycete Fungus** pathogen **Pyricularia oryzae**.
80. **Blast of Rice** remains an important exam topic due to its historical impact: **Spindle-shaped lesions on leaves; neck blast stage**.
81. The crop disease **Panama Wilt of Banana** is caused by the **Soil Fungus** pathogen **Fusarium oxysporum f. sp. cubense**.
82. **Panama Wilt of Banana** remains an important exam topic due to its historical impact: **Vascular wilting; reclaimed by growing resistant varieties**.
83. The crop disease **Black Arm of Cotton** is caused by the **Bacterium** pathogen **Xanthomonas campestris pv. malvacearum**.
84. **Black Arm of Cotton** remains an important exam topic due to its historical impact: **Angular leaf spots and boll rotting**.

Agricultural Entomology

85. **Pink Bollworm** is scientifically classified as **Pectinophora gossypiella** and belongs to the insect order **Lepidoptera**.
86. The pest **Pink Bollworm** attacks **Cotton** crops where its **Larva feeds inside the bolls, destroying lint**.
87. **Brown Plant Hopper (BPH)** is scientifically classified as **Nilaparvata lugens** and belongs to the insect order **Hemiptera**.
88. The pest **Brown Plant Hopper (BPH)** attacks **Rice** crops where its **Causes hopper burn due to cell sap sucking**.
89. **Rice Gundhi Bug** is scientifically classified as **Leptocorisa acuta** and belongs to the insect order **Hemiptera**.
90. The pest **Rice Gundhi Bug** attacks **Rice** crops where its **Sucks milk from developing grains in milky stage**.
91. **Fall Armyworm** is scientifically classified as **Spodoptera frugiperda** and belongs to the insect order **Lepidoptera**.
92. The pest **Fall Armyworm** attacks **Maize** crops where its **Invasive pest first reported in Karnataka in 2018**.
93. **Shoot Fly** is scientifically classified as **Atherigona soccata** and belongs to the insect order **Diptera**.
94. The pest **Shoot Fly** attacks **Sorghum** crops where its **Causes dead heart symptoms in young seedlings**.
95. **Gram Pod Borer** is scientifically classified as **Helicoverpa armigera** and belongs to the insect order **Lepidoptera**.

96. The pest **Gram Pod Borer** attacks **Chickpea** crops where its **Larva feeds on leaves and bores into pods**.
97. **Whitefly** is scientifically classified as **Bemisia tabaci** and belongs to the insect order **Hemiptera**.
98. The pest **Whitefly** attacks **Cotton/Tomato** crops where its **Vector for Leaf Curl Viruses**.
99. **Mustard Aphid** is scientifically classified as **Lipaphis erysimi** and belongs to the insect order **Hemiptera**.
100. The pest **Mustard Aphid** attacks **Mustard** crops where its **Sucks sap causing curling and drying of leaves**.

Karnataka Agro-Climatic Profiles

101. **North Eastern Transition Zone** of Karnataka includes key crop growing areas of **Bidar**.
102. The dominant soil type in **North Eastern Transition Zone** is **Red sandy loam**, with average rainfall of **700-900 mm**.
103. **Pigeon Pea (Tur)** represents the major commercial crop cultivated in the **North Eastern Transition Zone**.
104. **North Eastern Dry Zone** of Karnataka includes key crop growing areas of **Raichur, Kalaburagi**.
105. The dominant soil type in **North Eastern Dry Zone** is **Deep black soil**, with average rainfall of **600-800 mm**.
106. **Pigeon Pea, Cotton** represents the major commercial crop cultivated in the **North Eastern Dry Zone**.
107. **Northern Dry Zone** of Karnataka includes key crop growing areas of **Vijayapura, Bagalkot**.
108. The dominant soil type in **Northern Dry Zone** is **Medium to deep black soil**, with average rainfall of **500-700 mm**.
109. **Sorghum, Sunflower** represents the major commercial crop cultivated in the **Northern Dry Zone**.
110. **Central Dry Zone** of Karnataka includes key crop growing areas of **Chitradurga, Davanagere**.
111. The dominant soil type in **Central Dry Zone** is **Red sandy loam**, with average rainfall of **500-700 mm**.
112. **Ragi, Groundnut** represents the major commercial crop cultivated in the **Central Dry Zone**.
113. **Eastern Dry Zone** of Karnataka includes key crop growing areas of **Bengaluru Rural/Urban, Kolar**.
114. The dominant soil type in **Eastern Dry Zone** is **Red loamy and sandy soil**, with average rainfall of **700-900 mm**.
115. **Finger Millet (Ragi), Vegetables** represents the major commercial crop cultivated in the **Eastern Dry Zone**.
116. **Southern Dry Zone** of Karnataka includes key crop growing areas of **Mysuru, Mandya**.
117. The dominant soil type in **Southern Dry Zone** is **Red sandy loam**, with average rainfall of **600-800 mm**.
118. **Sugarcane, Paddy, Ragi** represents the major commercial crop cultivated in the **Southern Dry Zone**.
119. **Southern Transition Zone** of Karnataka includes key crop growing areas of **Hassan, Shivamogga**.
120. The dominant soil type in **Southern Transition Zone** is **Red sandy to clayey loam**, with average rainfall of **800-1500 mm**.
121. **Paddy, Maize, Arecanut** represents the major commercial crop cultivated in the **Southern Transition Zone**.
122. **Northern Transition Zone** of Karnataka includes key crop growing areas of **Dharwad, Belagavi**.
123. The dominant soil type in **Northern Transition Zone** is **Black clayey and red sandy soil**, with average rainfall of **800-1000 mm**.
124. **Soybean, Cotton, Maize** represents the major commercial crop cultivated in the **Northern Transition Zone**.
125. **Hilly Zone** of Karnataka includes key crop growing areas of **Kodagu, Chikkamagaluru**.
126. The dominant soil type in **Hilly Zone** is **Lateritic and forest soils**, with average rainfall of **1500-3000 mm**.
127. **Coffee, Cardamom, Paddy** represents the major commercial crop cultivated in the **Hilly Zone**.
128. **Coastal Zone** of Karnataka includes key crop growing areas of **Dakshina Kannada, Udupi, Uttara K.**
129. The dominant soil type in **Coastal Zone** is **Lateritic and alluvial soils**, with average rainfall of **3000-4000 mm**.
130. **Paddy, Cashew, Coconut** represents the major commercial crop cultivated in the **Coastal Zone**.

State Agricultural Universities

131. **University of Agricultural Sciences (UASB)** was established in the year **1964** to cater to regional farming needs.
132. The administrative headquarters of **University of Agricultural Sciences (UASB)** is located at **Bengaluru**, focusing on **Dry-land farming, millets, biofertilizers**.
133. **University of Agricultural Sciences (UASD)** was established in the year **1986** to cater to regional farming needs.
134. The administrative headquarters of **University of Agricultural Sciences (UASD)** is located at **Dharwad**, focusing on **Cotton, soybean, wheat breed improvements**.
135. **University of Agricultural Sciences (UASR)** was established in the year **2008** to cater to regional farming needs.
136. The administrative headquarters of **University of Agricultural Sciences (UASR)** is located at **Raichur**, focusing on **Pulses (Tur), cotton, resource management**.
137. **Keladi Shivappa Nayaka University (UAHS)** was established in the year **2012** to cater to regional farming needs.
138. The administrative headquarters of **Keladi Shivappa Nayaka University (UAHS)** is located at **Shivamogga**, focusing on **Spices, plantation crops, forestry**.
139. **University of Horticultural Sciences (UHSB)** was established in the year **2008** to cater to regional farming needs.
140. The administrative headquarters of **University of Horticultural Sciences (UHSB)** is located at **Bagalkot**, focusing on **Pomegranate, grapes, floriculture**.
141. **Karnataka Veterinary, Animal & Fisheries (KVAFSU)** was established in the year **2004** to cater to regional farming needs.
142. The administrative headquarters of **Karnataka Veterinary, Animal & Fisheries (KVAFSU)** is located at **Bidar**, focusing on **Indigenous breeds (Deoni), fisheries**.

Karnataka Agricultural Schemes

143. The state-sponsored scheme **Krishi Bhagya** was launched by Karnataka in **2014**.
144. Under **Krishi Bhagya**, small farmers are supported through **Rainwater harvesting through farm ponds (Krishi Honda) and micro-irrigation in dry zones**.
145. The state-sponsored scheme **Ganga Kalyana** was launched by Karnataka in **1997**.
146. Under **Ganga Kalyana**, small farmers are supported through **Subsidized drilling of borewells and installation of pump sets for SC/ST and minority farmers**.
147. The state-sponsored scheme **Raitha Siri** was launched by Karnataka in **2019**.
148. Under **Raitha Siri**, small farmers are supported through **Financial incentive of Rs. 10,000 per hectare for growing minor millets (Siri Dhanya)**.
149. The state-sponsored scheme **Raitha Vidya Nidhi** was launched by Karnataka in **2021**.
150. Under **Raitha Vidya Nidhi**, small farmers are supported through **Scholarships for the children of farmers to pursue higher and technical education**.
151. The state-sponsored scheme **Savayava Bhagya** was launched by Karnataka in **2013**.
152. Under **Savayava Bhagya**, small farmers are supported through **Promotion of organic farming through model organic villages in each taluk**.
153. The state-sponsored scheme **Krusha Aranya Protsaha Yojane** was launched by Karnataka in **2011**.
154. Under **Krusha Aranya Protsaha Yojane**, small farmers are supported through **Agro-forestry survival incentive of Rs. 30 per plant for tree seedlings planted on farmlands**.

Animal Husbandry & Breeds

155. **Hallikar** is a notable **Cattle** breed native to the region of **Karnataka**.
156. The **Hallikar** breed is highly valued for being a **Famous draft breed characterized by long, vertical horns and alert disposition**.
157. **Deoni** is a notable **Cattle** breed native to the region of **Karnataka/Maharashtra**.

158. The **Deoni** breed is highly valued for being a **Dual-purpose breed showing Gir-like forehead and black/white coat**.

159. **Amritmahal** is a notable **Cattle** breed native to the region of **Karnataka**.

160. The **Amritmahal** breed is highly valued for being a **Premium draft breed used historically for royal baggage transport**.

161. **Krishna Valley** is a notable **Cattle** breed native to the region of **Karnataka/Maharashtra**.

162. The **Krishna Valley** breed is highly valued for being a **Draft breed native to the banks of the Krishna River**.

163. **Murrah** is a notable **Buffalo** breed native to the region of **Haryana**.

164. The **Murrah** breed is highly valued for being a **Jet black dairy buffalo with tightly curved horns and high milk fat yield**.

165. **Surti** is a notable **Buffalo** breed native to the region of **Gujarat**.

166. The **Surti** breed is highly valued for being a **Dairy buffalo showing distinct sickle-shaped horns and white brow bands**.

High-Yield Technical One-Liners

167. Water Use Efficiency (WUE) is highest in finger millet (13.4 kg/ha-mm) due to its specialized C4 photosynthetic pathway.

168. Crown Root Initiation (CRI) is the most critical stage for irrigation in wheat, occurring exactly 21 days after sowing.

169. Puddling in rice destroys the soil structure, reducing percolation loss and creating an impervious subsurface layer.

170. Nipping in chickpea involves the removal of the terminal buds to encourage lateral branching, leading to higher pod yield.

171. Pendimethalin is applied as a pre-emergence herbicide to control annual grasses and broad-leaved weeds in pulses and groundnut.

172. Sorghum sickness is caused by nitrogen immobilization due to the high C:N ratio of sorghum stubble remaining in the soil.

173. Relay cropping refers to the sowing of a second crop before the harvest of the standing first crop (e.g., sowing Lathyrus in standing rice).

174. Tetrazolium test determines seed viability by measuring the activity of dehydrogenase enzymes, turning live cells red.

175. Real Value of Seed (%) is calculated by multiplying the Purity Percentage by the Germination Percentage and dividing by 100.

176. Orthodox seeds can survive low temperatures and drying down to 5% moisture content without losing viability (e.g., Wheat, Rice).

177. Recalcitrant seeds lose viability rapidly if dried or stored at freezing temperatures (e.g., Cocoa, Mango, Rubber).

178. NABARD was established on 12th July 1982 based on the recommendations of the B. Sivaraman Committee.

179. Commission for Agricultural Costs and Prices (CACP) recommends the Minimum Support Prices (MSP) for 23 mandated crops in India.

180. Bordeaux mixture was discovered by P.M.A. Millardet in 1885 to control Downy Mildew of Grapes caused by Plasmopara viticola.

181. First hybrid variety of Cotton in the world is H-4, developed by Dr. C.T. Patel at Surat in 1970.

182. The dwarfing gene in wheat is Norin-10, which was introduced into Mexican varieties by Dr. Norman E. Borlaug.

183. The dwarfing gene in rice is Dee-geo-woo-gen (DGWG), which was used to develop the miracle rice variety IR-8.

184. Active soil acidity is due to hydrogen (H⁺) ions in the soil solution, whereas reserve acidity is due to aluminum (Al³⁺) on exchange sites.

185. Reclamation of sodic/alkali soil requires the application of Gypsum (calcium sulfate) to replace exchangeable sodium with calcium.

186. Reclamation of acid soils requires the application of Lime (calcium carbonate) to neutralize active hydrogen ions.

187. Central atom of the chlorophyll molecule is Magnesium (Mg), which is a mobile nutrient showing deficiency on older leaves first.

188. Terminal bud death in plants is a characteristic symptom of Calcium (Ca) and Boron (B) deficiency.
189. Yellowing of tea leaves is a classic symptom of Sulfur (S) deficiency.
190. Khaira disease of rice is caused by the deficiency of Zinc (Zn) and was first discovered by Y.L. Nene at Pantnagar.
191. Gestation period of a cow is 283 days, whereas for a buffalo it is 310 days.
192. Foot and Mouth Disease (FMD) in cattle is caused by an Aphthovirus (Picorna virus group) and is highly contagious.
193. Anthrax is caused by *Bacillus anthracis*, a spore-forming bacterium; opening the carcass is strictly prohibited.
194. First Agricultural University in India is GBPUAT, Pantnagar, established in 1960 and modeled after Land Grant Universities.
195. Seeds Act was passed by the Indian Parliament in 1966 and came into force in October 1969.
196. Operation Flood was launched in 1970 by the National Dairy Development Board (NDDB) led by Dr. Verghese Kurien.
197. Monopsony is a market structure characterized by a single buyer and multiple sellers (e.g. sugar factory in a region).
198. Double cross hybrid maize technique was developed by D.F. Jones in 1918 to produce hybrid seed commercially.
199. The term 'heterosis' (hybrid vigor) was coined by G.H. Shull in 1914.
200. Gottlieb Haberlandt is known as the Father of Plant Tissue Culture for introducing the concept of totipotency.
201. *Agrobacterium tumefaciens* is known as nature's genetic engineer because of its ability to transfer T-DNA via its Ti plasmid.
202. Anther or pollen culture is used to produce haploid plants, first demonstrated by Guha and Maheshwari in *Datura*.
203. Biolistics or Gene Gun method uses gold or tungsten microprojectiles coated with DNA to transform plant cells.
204. First KVK (Krishi Vigyan Kendra) was established in Pondicherry in 1974 based on the recommendation of Mohan Singh Mehta Committee.
205. ATMA (Agricultural Technology Management Agency) is a registered society operating at the district level for technology dissemination.
206. Subsoiler is a primary tillage implement used to break hard soil pans below the plow layer without turning the soil.
207. Rotavator performs both primary and secondary tillage simultaneously, pulverizing the soil in a single pass.
208. Draft of an agricultural implement is the horizontal component of pull parallel to the direction of motion.
209. Deoni cattle breed is dual-purpose, native to Bidar district in Karnataka, and famous for its black-and-white spotted coat.
210. Sirsi Supari is the first arecanut in India to receive a GI (Geographical Indication) tag, grown in Uttara Kannada, Karnataka.
211. International Year of Millets was celebrated globally in 2023 following a proposal by India.
212. Karnataka ranks first in India in total silk production, contributing nearly 45-50% of the country's mulberry silk.
213. Methane (CH₄) gas is released from anaerobic flooded rice fields due to the activity of methanogenic bacteria.
214. C₄ plants like Maize, Sorghum, and Sugarcane have higher water-use efficiency and exhibit Kranz anatomy.
215. Glyphosate is a non-selective, post-emergence systemic herbicide that targets the EPSP synthase pathway.
216. Biuret content in urea should not exceed 1.5% for soil application to prevent toxicity to germinating seeds.
217. Kaolinite is a 1:1 non-expanding type silicate clay mineral with very low Cation Exchange Capacity (3-15 meq/100g).
218. Bulk density of sandy soil (1.6 g/cm³) is higher than that of clay soil (1.1 g/cm³) due to larger sand particle spacing.
219. Particle density of mineral soils is generally taken as 2.65 g/cm³, representing the solid soil phase.
220. Organic carbon in soil organic matter is estimated by multiplying organic matter content by 0.58 (Bemmelen factor).
221. Phosphorus is most available in soils in the form of orthophosphate ions (H₂PO₄⁻) at pH values between 6.0 and 7.0.
222. Pure Line Theory was proposed by W. Johannsen in 1903 while working on the Princess variety of common bean.

223. Terms Gene, Genotype, and Phenotype were coined by W. Johannsen in 1903.
224. DNA double helix structure was solved by Watson and Crick in 1953 using X-ray diffraction data from Rosalind Franklin.
225. Cytoplasmic Male Sterility (CMS) is utilized for hybrid seed production in onion and pearl millet to avoid manual emasculation.
226. Emasculation is the manual removal of stamens/anthers from bisexual flowers before pollen shedding.
227. Law of Segregation is also called the Law of Purity of Gametes, stating that alleles do not blend during gamete formation.
228. Green Revolution in India was initiated in 1966 with the introduction of high-yielding semi-dwarf wheat varieties.
229. Mycorrhizae (specifically VAM) form symbiotic associations with roots, enhancing the uptake of immobile phosphorus.
230. Damping off of seedlings in nurseries is caused by the oomycete *Pythium debaryanum* under high moisture conditions.
231. Honeybees decode distance and direction of food sources through dances: round dance (<50m) and waggle dance (>50m).
232. Khapra beetle (*Trogoderma granarium*) is a major storage pest of wheat, capable of surviving in very dry conditions.
233. Bower system (Arbour system) of training grapes yields the highest production in South India due to maximum canopy exposure.
234. Mango variety Amrapali is a hybrid of Dashehari x Neelum ($D \times N = A$), characterized by dwarf habit and high density planting.
235. Mango variety Mallika is a hybrid of Neelum x Dashehari ($N \times D = M$), yielding large fruits with high beta-carotene content.
236. Mango variety Sindhu is a seedless hybrid developed from a cross of Ratna x Alphonso.
237. Isolation distance is the minimum distance required between seed crop and contaminant crop fields to prevent cross-pollination.
238. Adopter categories: Innovators (2.5%), Early Adopters (13.5%), Early Majority (34%), Late Majority (34%), Laggards (16%).
239. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
240. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
241. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
242. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
243. The edible seed portion of **Soybean** contains approximately **26%** protein content.
244. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
245. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
246. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
247. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
248. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
249. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
250. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
251. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
252. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
253. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
254. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
255. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
256. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
257. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.

258. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
259. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
260. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
261. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
262. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
263. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
264. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
265. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
266. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
267. The edible seed portion of **Soybean** contains approximately **26%** protein content.
268. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
269. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
270. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
271. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
272. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
273. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
274. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
275. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
276. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
277. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
278. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
279. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
280. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
281. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
282. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
283. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
284. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
285. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
286. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
287. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
288. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
289. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
290. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
291. The edible seed portion of **Soybean** contains approximately **26%** protein content.
292. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
293. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
294. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
295. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.

296. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
297. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
298. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
299. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
300. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
301. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
302. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
303. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
304. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
305. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
306. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
307. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
308. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
309. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
310. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
311. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
312. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
313. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
314. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
315. The edible seed portion of **Soybean** contains approximately **26%** protein content.
316. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
317. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
318. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
319. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
320. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
321. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
322. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
323. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
324. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
325. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
326. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
327. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
328. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
329. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
330. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
331. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
332. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
333. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.

334. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
335. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
336. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
337. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
338. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
339. The edible seed portion of **Soybean** contains approximately **26%** protein content.
340. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
341. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
342. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
343. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
344. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
345. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
346. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
347. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
348. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
349. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
350. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
351. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
352. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
353. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
354. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
355. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
356. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
357. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
358. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
359. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
360. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
361. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
362. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
363. The edible seed portion of **Soybean** contains approximately **26%** protein content.
364. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
365. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
366. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
367. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
368. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
369. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
370. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
371. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.

372. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
373. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
374. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
375. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
376. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
377. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
378. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
379. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
380. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
381. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
382. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
383. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
384. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
385. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
386. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
387. The edible seed portion of **Soybean** contains approximately **26%** protein content.
388. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
389. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
390. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
391. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
392. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
393. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
394. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
395. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
396. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
397. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
398. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
399. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
400. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
401. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
402. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
403. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
404. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
405. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
406. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
407. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
408. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
409. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.

410. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
411. The edible seed portion of **Soybean** contains approximately **26%** protein content.
412. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
413. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
414. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
415. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
416. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
417. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
418. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
419. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
420. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
421. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
422. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
423. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
424. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
425. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
426. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
427. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
428. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
429. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
430. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
431. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
432. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
433. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
434. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
435. The edible seed portion of **Soybean** contains approximately **26%** protein content.
436. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
437. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
438. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
439. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
440. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
441. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
442. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
443. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
444. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
445. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
446. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
447. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.

448. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
449. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
450. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
451. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
452. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
453. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
454. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
455. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
456. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
457. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
458. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
459. The edible seed portion of **Soybean** contains approximately **26%** protein content.
460. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
461. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
462. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
463. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
464. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
465. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
466. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
467. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
468. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
469. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
470. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
471. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
472. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
473. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
474. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
475. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
476. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
477. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
478. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
479. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
480. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
481. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
482. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
483. The edible seed portion of **Soybean** contains approximately **26%** protein content.
484. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
485. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.

486. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
487. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
488. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
489. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
490. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
491. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
492. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
493. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
494. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
495. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
496. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
497. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
498. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
499. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
500. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
501. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
502. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
503. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
504. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
505. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
506. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
507. The edible seed portion of **Soybean** contains approximately **26%** protein content.
508. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
509. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
510. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
511. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
512. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
513. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
514. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
515. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
516. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
517. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
518. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
519. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
520. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
521. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
522. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
523. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.

524. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
525. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
526. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
527. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
528. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
529. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
530. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
531. The edible seed portion of **Soybean** contains approximately **26%** protein content.
532. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
533. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
534. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
535. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
536. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
537. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
538. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
539. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
540. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
541. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
542. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
543. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
544. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
545. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
546. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
547. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
548. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
549. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
550. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
551. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
552. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
553. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
554. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
555. The edible seed portion of **Soybean** contains approximately **26%** protein content.
556. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
557. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
558. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
559. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
560. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
561. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.

562. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
563. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
564. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
565. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
566. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
567. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
568. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
569. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
570. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
571. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
572. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
573. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
574. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
575. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
576. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
577. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
578. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
579. The edible seed portion of **Soybean** contains approximately **26%** protein content.
580. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
581. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
582. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
583. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
584. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
585. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
586. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
587. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
588. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
589. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
590. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
591. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
592. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
593. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
594. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
595. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
596. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
597. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
598. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
599. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.

600. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
601. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
602. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
603. The edible seed portion of **Soybean** contains approximately **26%** protein content.
604. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
605. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
606. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
607. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
608. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
609. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
610. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
611. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
612. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
613. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
614. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
615. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
616. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
617. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
618. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
619. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
620. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
621. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
622. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
623. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
624. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
625. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
626. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
627. The edible seed portion of **Soybean** contains approximately **26%** protein content.
628. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
629. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
630. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
631. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
632. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
633. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
634. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
635. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
636. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
637. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.

638. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
639. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
640. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
641. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
642. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
643. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
644. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
645. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
646. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
647. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
648. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
649. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
650. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
651. The edible seed portion of **Soybean** contains approximately **26%** protein content.
652. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
653. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
654. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
655. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
656. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
657. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
658. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
659. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
660. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
661. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
662. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
663. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
664. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
665. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
666. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
667. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
668. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
669. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
670. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
671. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
672. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
673. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
674. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
675. The edible seed portion of **Soybean** contains approximately **26%** protein content.

676. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
677. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
678. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
679. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
680. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
681. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
682. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
683. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
684. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
685. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
686. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
687. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
688. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
689. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
690. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
691. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
692. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
693. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
694. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
695. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
696. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
697. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
698. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
699. The edible seed portion of **Soybean** contains approximately **26%** protein content.
700. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
701. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
702. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
703. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
704. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
705. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
706. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
707. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
708. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
709. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
710. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
711. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
712. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
713. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.

714. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
715. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
716. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
717. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
718. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
719. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
720. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
721. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
722. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
723. The edible seed portion of **Soybean** contains approximately **26%** protein content.
724. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
725. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
726. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
727. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
728. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
729. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
730. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
731. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
732. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
733. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
734. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
735. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
736. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
737. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
738. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
739. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
740. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
741. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
742. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
743. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
744. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
745. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
746. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
747. The edible seed portion of **Soybean** contains approximately **26%** protein content.
748. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
749. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
750. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
751. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.

752. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
753. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
754. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
755. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
756. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
757. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
758. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
759. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
760. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
761. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
762. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
763. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
764. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
765. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
766. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
767. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
768. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
769. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
770. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
771. The edible seed portion of **Soybean** contains approximately **26%** protein content.
772. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
773. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
774. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
775. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
776. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
777. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
778. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
779. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
780. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
781. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
782. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
783. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
784. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
785. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
786. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
787. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
788. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
789. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.

790. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
791. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
792. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
793. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
794. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
795. The edible seed portion of **Soybean** contains approximately **26%** protein content.
796. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
797. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
798. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
799. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
800. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
801. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
802. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
803. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
804. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
805. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
806. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
807. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
808. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
809. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
810. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
811. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
812. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
813. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
814. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
815. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
816. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
817. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
818. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
819. The edible seed portion of **Soybean** contains approximately **26%** protein content.
820. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
821. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
822. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
823. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
824. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
825. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
826. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
827. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.

828. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
829. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
830. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
831. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
832. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
833. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
834. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
835. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
836. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
837. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
838. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
839. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
840. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
841. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
842. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
843. The edible seed portion of **Soybean** contains approximately **26%** protein content.
844. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
845. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
846. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
847. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
848. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
849. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
850. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
851. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
852. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
853. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
854. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
855. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
856. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
857. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
858. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
859. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
860. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
861. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
862. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
863. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
864. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
865. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.

866. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
867. The edible seed portion of **Soybean** contains approximately **26%** protein content.
868. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
869. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
870. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
871. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
872. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
873. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
874. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
875. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
876. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
877. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
878. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
879. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
880. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
881. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
882. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
883. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
884. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
885. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
886. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
887. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
888. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
889. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
890. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
891. The edible seed portion of **Soybean** contains approximately **26%** protein content.
892. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
893. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
894. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
895. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
896. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
897. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
898. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
899. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
900. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
901. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
902. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
903. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.

904. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
905. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
906. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
907. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
908. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
909. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
910. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
911. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
912. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
913. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
914. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
915. The edible seed portion of **Soybean** contains approximately **26%** protein content.
916. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
917. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
918. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
919. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
920. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
921. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
922. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
923. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
924. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
925. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
926. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
927. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
928. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
929. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
930. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
931. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
932. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
933. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
934. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
935. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
936. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
937. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
938. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
939. The edible seed portion of **Soybean** contains approximately **26%** protein content.
940. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
941. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.

942. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
943. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
944. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
945. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
946. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
947. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
948. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
949. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
950. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
951. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
952. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
953. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
954. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
955. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.
956. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
957. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
958. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
959. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
960. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
961. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
962. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
963. The edible seed portion of **Soybean** contains approximately **26%** protein content.
964. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
965. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
966. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
967. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
968. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
969. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
970. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
971. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
972. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
973. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
974. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
975. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
976. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.
977. Average seed rate required for sowing one hectare of land with **Sorghum (Jowar)** is **80-100 kg/ha**.
978. The edible seed portion of **Maize** contains approximately **40-42%** protein content.
979. Recommended field planting spacing for **Soybean** under standard cultivation packages is **30 x 10 cm**.

980. Average seed rate required for sowing one hectare of land with **Groundnut** is **100 kg/ha**.
981. The edible seed portion of **Chickpea (Bengal Gram)** contains approximately **6-7%** protein content.
982. Recommended field planting spacing for **Rice** under standard cultivation packages is **30 x 10 cm**.
983. Average seed rate required for sowing one hectare of land with **Wheat** is **15-20 kg/ha**.
984. The edible seed portion of **Finger Millet (Ragi)** contains approximately **10-12%** protein content.
985. Recommended field planting spacing for **Sorghum (Jowar)** under standard cultivation packages is **45 x 15 cm**.
986. Average seed rate required for sowing one hectare of land with **Maize** is **75-80 kg/ha**.
987. The edible seed portion of **Soybean** contains approximately **26%** protein content.
988. Recommended field planting spacing for **Groundnut** under standard cultivation packages is **20 x 15 cm**.
989. Average seed rate required for sowing one hectare of land with **Chickpea (Bengal Gram)** is **10 kg/ha**.
990. The edible seed portion of **Rice** contains approximately **11-12% (gluten)** protein content.
991. Recommended field planting spacing for **Wheat** under standard cultivation packages is **45 x 15 cm**.
992. Average seed rate required for sowing one hectare of land with **Finger Millet (Ragi)** is **70-80 kg/ha**.
993. The edible seed portion of **Sorghum (Jowar)** contains approximately **10% (zein)** protein content.
994. Recommended field planting spacing for **Maize** under standard cultivation packages is **30 x 10 cm**.
995. Average seed rate required for sowing one hectare of land with **Soybean** is **30-40 kg/ha (drilling)**.
996. The edible seed portion of **Groundnut** contains approximately **21%** protein content.
997. Recommended field planting spacing for **Chickpea (Bengal Gram)** under standard cultivation packages is **22.5 x 10 cm**.
998. Average seed rate required for sowing one hectare of land with **Rice** is **10-12 kg/ha**.
999. The edible seed portion of **Wheat** contains approximately **7-8%** protein content.
1000. Recommended field planting spacing for **Finger Millet (Ragi)** under standard cultivation packages is **60 x 20 cm**.

--- Expected Questions ---

Expected Questions Section

Top 500 Expected MCQs

This section contains a compiled pool of expected MCQs covering the core subjects.

Q1. What is the botanical scientific name of Rice?

- A. Cicer rubra
- B. Oryza sativa
- C. Triticum vulgare
- D. Oryza gluma
- **Correct Answer: B**
- **Explanation:** The scientific name of Rice is Oryza sativa, belonging to the family Gramineae/Poaceae.

Q2. What is the botanical scientific name of Wheat?

- A. Oryza gluma
- B. Triticum vulgare
- C. Cicer rubra
- D. Triticum aestivum
- **Correct Answer: D**
- **Explanation:** The scientific name of Wheat is Triticum aestivum, belonging to the family Gramineae/Poaceae.

Q3. What is the botanical scientific name of Finger Millet (Ragi)?

- A. Cicer rubra
- B. Eleusine coracana
- C. Triticum vulgare
- D. Oryza gluma
- **Correct Answer: B**
- **Explanation:** The scientific name of Finger Millet (Ragi) is Eleusine coracana, belonging to the family Gramineae/Poaceae.

Q4. What is the botanical scientific name of Sorghum (Jowar)?

- A. Triticum vulgare
- B. Sorghum bicolor
- C. Cicer rubra
- D. Oryza gluma
- **Correct Answer: B**
- **Explanation:** The scientific name of Sorghum (Jowar) is Sorghum bicolor, belonging to the family Gramineae/Poaceae.

Q5. What is the botanical scientific name of Maize?

- A. Zea mays
- B. Oryza gluma
- C. Cicer rubra
- D. Triticum vulgare

• **Correct Answer: A**

• **Explanation:** The scientific name of Maize is Zea mays, belonging to the family Gramineae/Poaceae.

Q6. What is the botanical scientific name of Soybean?

- A. Cicer rubra
- B. Triticum vulgare
- C. Oryza gluma
- D. Glycine max

• **Correct Answer: D**

• **Explanation:** The scientific name of Soybean is Glycine max, belonging to the family Leguminosae/Fabaceae.

Q7. What is the botanical scientific name of Groundnut?

- A. Cicer rubra
- B. Oryza gluma
- C. Arachis hypogaea
- D. Triticum vulgare

• **Correct Answer: C**

• **Explanation:** The scientific name of Groundnut is Arachis hypogaea, belonging to the family Leguminosae/Fabaceae.

Q8. What is the botanical scientific name of Chickpea (Bengal Gram)?

- A. Cicer arietinum
- B. Triticum vulgare
- C. Cicer rubra
- D. Oryza gluma

• **Correct Answer: A**

• **Explanation:** The scientific name of Chickpea (Bengal Gram) is Cicer arietinum, belonging to the family Leguminosae/Fabaceae.

Q9. What color tag is attached to the bag of Breeder Seed?

- A. Golden Yellow
- B. Red
- C. Green
- D. Orange

• **Correct Answer: A**

• **Explanation:** Breeder Seed is labeled with a Golden Yellow colored tag, indicating a genetic purity of 100%.

Q10. What color tag is attached to the bag of Foundation Seed?

- A. Orange
- B. Green
- C. Red
- D. White

• **Correct Answer: D**

• **Explanation:** Foundation Seed is labeled with a White colored tag, indicating a genetic purity of 99.5%.

Q11. What color tag is attached to the bag of Certified Seed?

- A. Azure Blue
- B. Red
- C. Green
- D. Orange

• **Correct Answer: A**

• **Explanation:** Certified Seed is labeled with a Azure Blue colored tag, indicating a genetic purity of 99.0%.

Q12. What color tag is attached to the bag of Truthful Label Seed?

- A. Opal Green
- B. Green
- C. Red
- D. Orange

• **Correct Answer: A**

• **Explanation:** Truthful Label Seed is labeled with a Opal Green colored tag, indicating a genetic purity of 97.0%.

Q13. Deficiency of which symptom is characteristic of Nitrogen (N) deficiency in plants?

- A. Necrosis on margins
- B. V-shaped yellowing on older leaf tips
- C. Stunted growth with no chlorosis
- D. Rosetting of leaves
- E. Purple coloration on younger leaves

• **Correct Answer: B**

• **Explanation:** Deficiency of Nitrogen (N) leads to V-shaped yellowing on older leaf tips because it is a key Constituent of proteins and chlorophyll.

Q14. Deficiency of which symptom is characteristic of Phosphorus (P) deficiency in plants?

- A. Rosetting of leaves
- B. Purple coloration on younger leaves
- C. Stunted growth with no chlorosis
- D. Necrosis on margins
- E. Purple/bronze coloration on older leaves

• **Correct Answer: E**

- **Explanation:** Deficiency of Phosphorus (P) leads to Purple/bronze coloration on older leaves because it is a key Energy storage and transfer (ATP).

Q15. Deficiency of which symptom is characteristic of Potassium (K) deficiency in plants?

- A. Rosetting of leaves
- B. Stunted growth with no chlorosis
- C. Purple coloration on younger leaves
- D. Scorching and burning of leaf margins
- E. Necrosis on margins

- **Correct Answer: D**

- **Explanation:** Deficiency of Potassium (K) leads to Scorching and burning of leaf margins because it is a key Stomatal regulation and enzyme activation.

Q16. Deficiency of which symptom is characteristic of Calcium (Ca) deficiency in plants?

- A. Terminal bud death and hooking of leaf tips
- B. Rosetting of leaves
- C. Necrosis on margins
- D. Stunted growth with no chlorosis
- E. Purple coloration on younger leaves

- **Correct Answer: A**

- **Explanation:** Deficiency of Calcium (Ca) leads to Terminal bud death and hooking of leaf tips because it is a key Cell wall constituent (calcium pectate).

Q17. Deficiency of which symptom is characteristic of Magnesium (Mg) deficiency in plants?

- A. Interveinal chlorosis on older leaves
- B. Purple coloration on younger leaves
- C. Necrosis on margins
- D. Stunted growth with no chlorosis
- E. Rosetting of leaves

- **Correct Answer: A**

- **Explanation:** Deficiency of Magnesium (Mg) leads to Interveinal chlorosis on older leaves because it is a key Central atom of chlorophyll.

Q18. Deficiency of which symptom is characteristic of Sulfur (S) deficiency in plants?

- A. Stunted growth with no chlorosis
- B. Rosetting of leaves
- C. Purple coloration on younger leaves
- D. Yellowing of younger leaves and tea yellowing
- E. Necrosis on margins

- **Correct Answer: D**

- **Explanation:** Deficiency of Sulfur (S) leads to Yellowing of younger leaves and tea yellowing because it is a key Constituent of amino acids (methionine, cysteine).

Q19. Deficiency of which symptom is characteristic of Iron (Fe) deficiency in plants?

- A. Necrosis on margins
- B. Stunted growth with no chlorosis
- C. Purple coloration on younger leaves
- D. Rosetting of leaves
- E. Interveinal chlorosis on younger leaves

• **Correct Answer: E**

- **Explanation:** Deficiency of Iron (Fe) leads to Interveinal chlorosis on younger leaves because it is a key Chlorophyll synthesis and electron transfer.

Q20. Deficiency of which symptom is characteristic of Zinc (Zn) deficiency in plants?

- A. Rosetting of leaves
- B. Khaira disease in rice and white bud in maize
- C. Stunted growth with no chlorosis
- D. Purple coloration on younger leaves
- E. Necrosis on margins

• **Correct Answer: B**

- **Explanation:** Deficiency of Zinc (Zn) leads to Khaira disease in rice and white bud in maize because it is a key Auxin synthesis and enzyme activation.

Q21. Deficiency of which symptom is characteristic of Boron (B) deficiency in plants?

- A. Hollow heart in groundnut and fruit cracking
- B. Rosetting of leaves
- C. Purple coloration on younger leaves
- D. Stunted growth with no chlorosis
- E. Necrosis on margins

• **Correct Answer: A**

- **Explanation:** Deficiency of Boron (B) leads to Hollow heart in groundnut and fruit cracking because it is a key Pollen germination and sugar translocation.

Q22. Deficiency of which symptom is characteristic of Copper (Cu) deficiency in plants?

- A. Purple coloration on younger leaves
- B. Stunted growth with no chlorosis
- C. Citrus dieback and exanthema
- D. Rosetting of leaves
- E. Necrosis on margins

• **Correct Answer: C**

- **Explanation:** Deficiency of Copper (Cu) leads to Citrus dieback and exanthema because it is a key Photosynthesis and respiration enzyme constituent.

Q23. What is the causal pathogen of Late Blight of Potato?

- A. Puccinia graminis
- B. Meloidogyne incognita
- C. Phytophthora infestans
- D. Albugo candida
- **Correct Answer: C**
- **Explanation:** Late Blight of Potato is caused by the Oomycete Fungus pathogen Phytophthora infestans (Irish Famine of 1845).

Q24. What is the causal pathogen of Loose Smut of Wheat?

- A. Puccinia graminis
- B. Ustilago nuda tritici
- C. Albugo candida
- D. Meloidogyne incognita
- **Correct Answer: B**
- **Explanation:** Loose Smut of Wheat is caused by the Basidiomycete Fungus pathogen Ustilago nuda tritici (Internally seed-borne; controlled by Vitavax).

Q25. What is the causal pathogen of Karnal Bunt of Wheat?

- A. Puccinia graminis
- B. Albugo candida
- C. Tilletia indica
- D. Meloidogyne incognita
- **Correct Answer: C**
- **Explanation:** Karnal Bunt of Wheat is caused by the Basidiomycete Fungus pathogen Tilletia indica (Fishy smell due to trimethylamine; cancer of wheat).

Q26. What is the causal pathogen of Citrus Canker?

- A. Meloidogyne incognita
- B. Xanthomonas axonopodis pv. citri
- C. Albugo candida
- D. Puccinia graminis
- **Correct Answer: B**
- **Explanation:** Citrus Canker is caused by the Bacterium pathogen Xanthomonas axonopodis pv. citri (Acidic copper sprays or streptomycin control).

Q27. What is the causal pathogen of Tikka Disease of Groundnut?

- A. Cercospora personata
- B. Puccinia graminis
- C. Albugo candida
- D. Meloidogyne incognita
- **Correct Answer: A**
- **Explanation:** Tikka Disease of Groundnut is caused by the Ascomycete Fungus pathogen Cercospora personata

(Controlled by Mancozeb or Carbendazim).

Q28. What is the causal pathogen of Red Rot of Sugarcane?

- A. Puccinia graminis
- B. Meloidogyne incognita
- C. Albugo candida
- D. Colletotrichum falcatum
- **Correct Answer: D**
- **Explanation:** Red Rot of Sugarcane is caused by the Deuteromycete Fungus pathogen Colletotrichum falcatum (Red discoloration of pith with white bands).

Q29. What is the causal pathogen of Ergot of Bajra?

- A. Meloidogyne incognita
- B. Albugo candida
- C. Puccinia graminis
- D. Claviceps fusiformis
- **Correct Answer: D**
- **Explanation:** Ergot of Bajra is caused by the Ascomycete Fungus pathogen Claviceps fusiformis (Sclerotia produce toxic alkaloids causing ergotism).

Q30. What is the causal pathogen of Blast of Rice?

- A. Puccinia graminis
- B. Pyricularia oryzae
- C. Meloidogyne incognita
- D. Albugo candida
- **Correct Answer: B**
- **Explanation:** Blast of Rice is caused by the Ascomycete Fungus pathogen Pyricularia oryzae (Spindle-shaped lesions on leaves; neck blast stage).

Q31. What is the causal pathogen of Panama Wilt of Banana?

- A. Albugo candida
- B. Meloidogyne incognita
- C. Puccinia graminis
- D. Fusarium oxysporum f. sp. cubense
- **Correct Answer: D**
- **Explanation:** Panama Wilt of Banana is caused by the Soil Fungus pathogen Fusarium oxysporum f. sp. cubense (Vascular wilting; reclaimed by growing resistant varieties).

Q32. What is the causal pathogen of Black Arm of Cotton?

- A. Puccinia graminis
- B. Albugo candida
- C. Meloidogyne incognita
- D. Xanthomonas campestris pv. malvacearum

- **Correct Answer: D**

- **Explanation:** Black Arm of Cotton is caused by the Bacterium pathogen *Xanthomonas campestris* pv. *malvacearum* (Angular leaf spots and boll rotting).

Q33. Which insect pest is scientifically named *Pectinophora gossypiella* and attacks Cotton?

- A. *Sitophilus oryzae*
- B. *Pectinophora gossypiella*
- C. *Tribolium castaneum*
- D. *Aphis craccivora*

- **Correct Answer: B**

- **Explanation:** Pink Bollworm (*Pectinophora gossypiella*) belongs to the order Lepidoptera and damages Cotton crops.

Q34. Which insect pest is scientifically named *Nilaparvata lugens* and attacks Rice?

- A. *Aphis craccivora*
- B. *Nilaparvata lugens*
- C. *Sitophilus oryzae*
- D. *Tribolium castaneum*

- **Correct Answer: B**

- **Explanation:** Brown Plant Hopper (BPH) (*Nilaparvata lugens*) belongs to the order Hemiptera and damages Rice crops.

Q35. Which insect pest is scientifically named *Leptocorisa acuta* and attacks Rice?

- A. *Sitophilus oryzae*
- B. *Aphis craccivora*
- C. *Leptocorisa acuta*
- D. *Tribolium castaneum*

- **Correct Answer: C**

- **Explanation:** Rice Gundhi Bug (*Leptocorisa acuta*) belongs to the order Hemiptera and damages Rice crops.

Q36. Which insect pest is scientifically named *Spodoptera frugiperda* and attacks Maize?

- A. *Aphis craccivora*
- B. *Sitophilus oryzae*
- C. *Spodoptera frugiperda*
- D. *Tribolium castaneum*

- **Correct Answer: C**

- **Explanation:** Fall Armyworm (*Spodoptera frugiperda*) belongs to the order Lepidoptera and damages Maize crops.

Q37. Which insect pest is scientifically named *Atherigona soccata* and attacks Sorghum?

- A. *Aphis craccivora*
- B. *Sitophilus oryzae*
- C. *Tribolium castaneum*
- D. *Atherigona soccata*

- **Correct Answer: D**

- **Explanation:** Shoot Fly (*Atherigona soccata*) belongs to the order Diptera and damages Sorghum crops.

Q38. Which insect pest is scientifically named *Helicoverpa armigera* and attacks Chickpea?

- A. *Tribolium castaneum*
- B. *Helicoverpa armigera*
- C. *Aphis craccivora*
- D. *Sitophilus oryzae*

- **Correct Answer: B**

- **Explanation:** Gram Pod Borer (*Helicoverpa armigera*) belongs to the order Lepidoptera and damages Chickpea crops.

Q39. Which insect pest is scientifically named *Bemisia tabaci* and attacks Cotton/Tomato?

- A. *Bemisia tabaci*
- B. *Tribolium castaneum*
- C. *Sitophilus oryzae*
- D. *Aphis craccivora*

- **Correct Answer: A**

- **Explanation:** Whitefly (*Bemisia tabaci*) belongs to the order Hemiptera and damages Cotton/Tomato crops.

Q40. Which insect pest is scientifically named *Lipaphis erysimi* and attacks Mustard?

- A. *Aphis craccivora*
- B. *Lipaphis erysimi*
- C. *Tribolium castaneum*
- D. *Sitophilus oryzae*

- **Correct Answer: B**

- **Explanation:** Mustard Aphid (*Lipaphis erysimi*) belongs to the order Hemiptera and damages Mustard crops.

Q41. What is the dominant soil type in the North Eastern Transition Zone of Karnataka?

- A. Alluvial soil
- B. Red sandy loam
- C. Lateritic soil
- D. Desert sand

- **Correct Answer: B**

- **Explanation:** North Eastern Transition Zone (covering districts like Bidar) has dominantly Red sandy loam.

Q42. What is the dominant soil type in the North Eastern Dry Zone of Karnataka?

- A. Alluvial soil
- B. Lateritic soil
- C. Desert sand
- D. Deep black soil

- **Correct Answer: D**

- **Explanation:** North Eastern Dry Zone (covering districts like Raichur, Kalaburagi) has dominantly Deep black soil.

Q43. What is the dominant soil type in the Northern Dry Zone of Karnataka?

- A. Alluvial soil
- B. Desert sand
- C. Lateritic soil
- D. Medium to deep black soil

- **Correct Answer: D**

- **Explanation:** Northern Dry Zone (covering districts like Vijayapura, Bagalkot) has dominantly Medium to deep black soil.

Q44. What is the dominant soil type in the Central Dry Zone of Karnataka?

- A. Lateritic soil
- B. Red sandy loam
- C. Alluvial soil
- D. Desert sand

- **Correct Answer: B**

- **Explanation:** Central Dry Zone (covering districts like Chitradurga, Davanagere) has dominantly Red sandy loam.

Q45. What is the dominant soil type in the Eastern Dry Zone of Karnataka?

- A. Red loamy and sandy soil
- B. Desert sand
- C. Lateritic soil
- D. Alluvial soil

- **Correct Answer: A**

- **Explanation:** Eastern Dry Zone (covering districts like Bengaluru Rural/Urban, Kolar) has dominantly Red loamy and sandy soil.

Q46. What is the dominant soil type in the Southern Dry Zone of Karnataka?

- A. Alluvial soil
- B. Desert sand
- C. Lateritic soil
- D. Red sandy loam

- **Correct Answer: D**

- **Explanation:** Southern Dry Zone (covering districts like Mysuru, Mandya) has dominantly Red sandy loam.

Q47. What is the dominant soil type in the Southern Transition Zone of Karnataka?

- A. Desert sand
- B. Lateritic soil
- C. Red sandy to clayey loam
- D. Alluvial soil

- **Correct Answer: C**

- **Explanation:** Southern Transition Zone (covering districts like Hassan, Shivamogga) has dominantly Red sandy to clayey loam.

Q48. What is the dominant soil type in the Northern Transition Zone of Karnataka?

- A. Desert sand
- B. Lateritic soil
- C. Black clayey and red sandy soil
- D. Alluvial soil

- **Correct Answer: C**

- **Explanation:** Northern Transition Zone (covering districts like Dharwad, Belagavi) has dominantly Black clayey and red sandy soil.

Q49. What is the dominant soil type in the Hilly Zone of Karnataka?

- A. Desert sand
- B. Alluvial soil
- C. Lateritic soil
- D. Lateritic and forest soils

- **Correct Answer: D**

- **Explanation:** Hilly Zone (covering districts like Kodagu, Chikkamagaluru) has dominantly Lateritic and forest soils.

Q50. What is the dominant soil type in the Coastal Zone of Karnataka?

- A. Lateritic and alluvial soils
- B. Alluvial soil
- C. Lateritic soil
- D. Desert sand

- **Correct Answer: A**

- **Explanation:** Coastal Zone (covering districts like Dakshina Kannada, Udupi, Uttara K.) has dominantly Lateritic and alluvial soils.

Q51. In which year was the 'Krishi Bhagya' scheme launched by the Government of Karnataka?

- A. 2016
- B. 2014
- C. 2010
- D. 2005

- **Correct Answer: B**

- **Explanation:** 'Krishi Bhagya' was launched in 2014 to support Rainwater harvesting through farm ponds (Krishi Honda) and micro-irrigation in dry zones.

Q52. In which year was the 'Ganga Kalyana' scheme launched by the Government of Karnataka?

- A. 2016
- B. 2005

- C. 1997
- D. 2010
- **Correct Answer: C**
- **Explanation:** 'Ganga Kalyana' was launched in 1997 to support Subsidized drilling of borewells and installation of pump sets for SC/ST and minority farmers.

Q53. In which year was the 'Raitha Siri' scheme launched by the Government of Karnataka?

- A. 2019
- B. 2010
- C. 2016
- D. 2005
- **Correct Answer: A**
- **Explanation:** 'Raitha Siri' was launched in 2019 to support Financial incentive of Rs. 10,000 per hectare for growing minor millets (Siri Dhanya).

Q54. In which year was the 'Raitha Vidya Nidhi' scheme launched by the Government of Karnataka?

- A. 2021
- B. 2010
- C. 2016
- D. 2005
- **Correct Answer: A**
- **Explanation:** 'Raitha Vidya Nidhi' was launched in 2021 to support Scholarships for the children of farmers to pursue higher and technical education.

Q55. In which year was the 'Savayava Bhagya' scheme launched by the Government of Karnataka?

- A. 2005
- B. 2010
- C. 2016
- D. 2013
- **Correct Answer: D**
- **Explanation:** 'Savayava Bhagya' was launched in 2013 to support Promotion of organic farming through model organic villages in each taluk.

Q56. In which year was the 'Krushi Aranya Protsaha Yojane' scheme launched by the Government of Karnataka?

- A. 2016
- B. 2010
- C. 2005
- D. 2011
- **Correct Answer: D**
- **Explanation:** 'Krushi Aranya Protsaha Yojane' was launched in 2011 to support Agro-forestry survival incentive of Rs. 30 per plant for tree seedlings planted on farmlands.

Q57. Which indigenous Cattle breed is native to Karnataka and described as: Famous draft breed characterized

by long, vertical horns and alert disposition?

- A. Gir
- B. Ongole
- C. Sahiwal
- D. Hallikar
- **Correct Answer: D**

• **Explanation:** Hallikar is a native Cattle breed of Karnataka.

Q58. Which indigenous Cattle breed is native to Karnataka/Maharashtra and described as: Dual-purpose breed showing Gir-like forehead and black/white coat?

- A. Ongole
- B. Gir
- C. Deoni
- D. Sahiwal
- **Correct Answer: C**

• **Explanation:** Deoni is a native Cattle breed of Karnataka/Maharashtra.

Q59. Which indigenous Cattle breed is native to Karnataka and described as: Premium draft breed used historically for royal baggage transport?

- A. Gir
- B. Ongole
- C. Amritmahal
- D. Sahiwal
- **Correct Answer: C**

• **Explanation:** Amritmahal is a native Cattle breed of Karnataka.

Q60. Which indigenous Cattle breed is native to Karnataka/Maharashtra and described as: Draft breed native to the banks of the Krishna River?

- A. Ongole
- B. Krishna Valley
- C. Sahiwal
- D. Gir
- **Correct Answer: B**

• **Explanation:** Krishna Valley is a native Cattle breed of Karnataka/Maharashtra.

Q61. Which indigenous Buffalo breed is native to Haryana and described as: Jet black dairy buffalo with tightly curved horns and high milk fat yield?

- A. Sahiwal
- B. Gir
- C. Murrah

- D. Ongole
- **Correct Answer: C**
- **Explanation:** Murrah is a native Buffalo breed of Haryana.

Q62. Which indigenous Buffalo breed is native to Gujarat and described as: Dairy buffalo showing distinct sickle-shaped horns and white brow bands?

- A. Ongole
- B. Gir
- C. Surti
- D. Sahiwal
- **Correct Answer: C**
- **Explanation:** Surti is a native Buffalo breed of Gujarat.

Q63. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #63)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q64. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #64)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q65. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #65)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q66. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #66)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum

- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q67. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #67)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q68. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #68)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q69. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #69)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q70. Regarding Economics, NABARD was established in which year? (Question Variant #70)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q71. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #71)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize

- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q72. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #72)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q73. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #73)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q74. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #74)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q75. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #75)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q76. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #76)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum

- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q77. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #77)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q78. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #78)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q79. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #79)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q80. Regarding Economics, NABARD was established in which year? (Question Variant #80)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q81. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #81)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize

- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q82. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #82)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q83. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #83)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q84. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #84)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q85. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #85)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q86. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #86)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum

- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q87. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #87)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q88. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #88)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q89. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #89)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q90. Regarding Economics, NABARD was established in which year? (Question Variant #90)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q91. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #91)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize

- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q92. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #92)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q93. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #93)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q94. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #94)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q95. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #95)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q96. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #96)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum

- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q97. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #97)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q98. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #98)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q99. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #99)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q100. Regarding Economics, NABARD was established in which year? (Question Variant #100)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q101. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #101)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize

- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q102. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #102)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q103. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #103)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q104. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #104)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q105. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #105)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q106. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #106)

- A. Phytophthora infestans
- B. Alternaria solani

- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q107. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #107)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q108. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #108)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q109. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #109)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q110. Regarding Economics, NABARD was established in which year? (Question Variant #110)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q111. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #111)

- A. Finger Millet (Ragi)

- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q112. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #112)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q113. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #113)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q114. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #114)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q115. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #115)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q116. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #116)

- A. Phytophthora infestans

- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q117. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #117)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q118. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #118)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q119. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #119)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q120. Regarding Economics, NABARD was established in which year? (Question Variant #120)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q121. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #121)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q122. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #122)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q123. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #123)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q124. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #124)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q125. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #125)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q126. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #126)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q127. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #127)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q128. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #128)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q129. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #129)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q130. Regarding Economics, NABARD was established in which year? (Question Variant #130)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q131. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #131)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q132. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #132)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q133. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #133)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q134. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #134)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q135. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #135)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q136. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #136)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q137. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #137)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q138. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #138)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q139. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #139)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q140. Regarding Economics, NABARD was established in which year? (Question Variant #140)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q141. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #141)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q142. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #142)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q143. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #143)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q144. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #144)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q145. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #145)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q146. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #146)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q147. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #147)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q148. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #148)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q149. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #149)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q150. Regarding Economics, NABARD was established in which year? (Question Variant #150)

- A. 1982
- B. 1975
- C. 1990

- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q151. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #151)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q152. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #152)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q153. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #153)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q154. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #154)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q155. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #155)

- A. Azure Blue
- B. White
- C. Golden Yellow

- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q156. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #156)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**
- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q157. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #157)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**
- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q158. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #158)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q159. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #159)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q160. Regarding Economics, NABARD was established in which year? (Question Variant #160)

- A. 1982
- B. 1975

- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q161. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #161)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q162. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #162)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q163. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #163)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q164. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #164)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q165. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #165)

- A. Azure Blue

- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q166. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #166)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**
- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q167. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #167)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**
- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q168. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #168)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q169. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #169)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q170. Regarding Economics, NABARD was established in which year? (Question Variant #170)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q171. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #171)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q172. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #172)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q173. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #173)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q174. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #174)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q175. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #175)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q176. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #176)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q177. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #177)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q178. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #178)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q179. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #179)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q180. Regarding Economics, NABARD was established in which year? (Question Variant #180)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q181. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #181)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q182. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #182)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q183. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #183)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q184. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #184)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q185. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #185)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q186. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #186)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q187. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #187)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q188. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #188)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q189. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #189)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q190. Regarding Economics, NABARD was established in which year? (Question Variant #190)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q191. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #191)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q192. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #192)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q193. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #193)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q194. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #194)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

- **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q195. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #195)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q196. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #196)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q197. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #197)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q198. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #198)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q199. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #199)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer

- D. Spencer Hatch

• **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q200. Regarding Economics, NABARD was established in which year? (Question Variant #200)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q201. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #201)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q202. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #202)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q203. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #203)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q204. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #204)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel

- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q205. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #205)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q206. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #206)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q207. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #207)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q208. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #208)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q209. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #209)

- A. Daniel Benor
- B. F.L. Brayne

- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q210. Regarding Economics, NABARD was established in which year? (Question Variant #210)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q211. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #211)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q212. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #212)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q213. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #213)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q214. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #214)

- A. D.F. Jones

- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q215. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #215)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q216. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #216)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q217. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #217)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q218. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #218)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q219. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #219)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q220. Regarding Economics, NABARD was established in which year? (Question Variant #220)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q221. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #221)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q222. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #222)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q223. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #223)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q224. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #224)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q225. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #225)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q226. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #226)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q227. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #227)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q228. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #228)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q229. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #229)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q230. Regarding Economics, NABARD was established in which year? (Question Variant #230)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q231. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #231)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q232. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #232)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q233. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #233)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q234. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #234)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q235. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #235)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q236. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #236)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q237. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #237)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q238. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #238)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q239. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #239)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q240. Regarding Economics, NABARD was established in which year? (Question Variant #240)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q241. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #241)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q242. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #242)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q243. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #243)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q244. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #244)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q245. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #245)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q246. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #246)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q247. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #247)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q248. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #248)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla

- D. Nagpur

• **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q249. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #249)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q250. Regarding Economics, NABARD was established in which year? (Question Variant #250)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q251. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #251)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q252. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #252)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q253. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #253)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)

- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q254. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #254)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q255. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #255)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q256. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #256)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**
- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q257. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #257)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**
- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q258. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #258)

- A. Hesaraghatta, Bengaluru
- B. Lucknow

- C. Shimla
- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q259. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #259)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q260. Regarding Economics, NABARD was established in which year? (Question Variant #260)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q261. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #261)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q262. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #262)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q263. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #263)

- A. Magnesium (Mg)
- B. Iron (Fe)

- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q264. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #264)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q265. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #265)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q266. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #266)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q267. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #267)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q268. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #268)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q269. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #269)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q270. Regarding Economics, NABARD was established in which year? (Question Variant #270)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q271. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #271)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q272. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #272)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q273. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #273)

- A. Magnesium (Mg)

- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q274. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #274)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q275. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #275)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q276. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #276)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q277. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #277)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q278. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant

#278)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q279. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #279)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q280. Regarding Economics, NABARD was established in which year? (Question Variant #280)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q281. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #281)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q282. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #282)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q283. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #283)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q284. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #284)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q285. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #285)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q286. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #286)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q287. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #287)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q288. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #288)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q289. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #289)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q290. Regarding Economics, NABARD was established in which year? (Question Variant #290)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q291. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #291)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q292. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #292)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q293. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #293)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q294. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #294)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q295. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #295)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q296. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #296)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q297. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #297)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q298. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #298)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q299. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #299)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q300. Regarding Economics, NABARD was established in which year? (Question Variant #300)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q301. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #301)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q302. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #302)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting

- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q303. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #303)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q304. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #304)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q305. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #305)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q306. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #306)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q307. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #307)

- A. Water Hyacinth
- B. Parthenium

- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q308. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #308)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q309. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #309)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q310. Regarding Economics, NABARD was established in which year? (Question Variant #310)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q311. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #311)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q312. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #312)

- A. Crown Root Initiation (CRI)
- B. Tillering

- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q313. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #313)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q314. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #314)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q315. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #315)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q316. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #316)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q317. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #317)

- A. Water Hyacinth

- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q318. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #318)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q319. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #319)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q320. Regarding Economics, NABARD was established in which year? (Question Variant #320)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q321. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #321)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q322. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #322)

- A. Crown Root Initiation (CRI)

- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q323. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #323)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q324. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #324)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q325. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #325)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q326. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #326)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q327. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #327)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q328. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #328)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q329. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #329)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q330. Regarding Economics, NABARD was established in which year? (Question Variant #330)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q331. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #331)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q332. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #332)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q333. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #333)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q334. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #334)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q335. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #335)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q336. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #336)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q337. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #337)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q338. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #338)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q339. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #339)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q340. Regarding Economics, NABARD was established in which year? (Question Variant #340)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q341. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #341)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q342. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #342)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q343. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #343)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q344. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #344)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

- **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q345. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #345)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q346. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #346)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q347. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #347)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q348. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #348)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q349. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #349)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q350. Regarding Economics, NABARD was established in which year? (Question Variant #350)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q351. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #351)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize

- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q352. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #352)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q353. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #353)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q354. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #354)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q355. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #355)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q356. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #356)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum

- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q357. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #357)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q358. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #358)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q359. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #359)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q360. Regarding Economics, NABARD was established in which year? (Question Variant #360)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q361. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #361)

- A. Finger Millet (Ragi)
- B. Wheat

- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q362. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #362)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q363. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #363)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q364. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #364)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q365. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #365)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q366. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #366)

- A. Phytophthora infestans

- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q367. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #367)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q368. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #368)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q369. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #369)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q370. Regarding Economics, NABARD was established in which year? (Question Variant #370)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q371. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #371)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q372. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #372)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q373. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #373)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q374. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #374)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q375. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #375)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q376. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #376)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q377. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #377)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q378. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #378)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

• **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q379. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #379)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

• **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q380. Regarding Economics, NABARD was established in which year? (Question Variant #380)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q381. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #381)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q382. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #382)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q383. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #383)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q384. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #384)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**
- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q385. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #385)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q386. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #386)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q387. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #387)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q388. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #388)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q389. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #389)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q390. Regarding Economics, NABARD was established in which year? (Question Variant #390)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q391. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #391)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q392. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #392)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q393. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #393)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q394. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #394)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q395. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #395)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q396. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #396)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q397. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #397)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q398. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #398)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q399. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #399)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q400. Regarding Economics, NABARD was established in which year? (Question Variant #400)

- A. 1982
- B. 1975
- C. 1990

- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q401. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #401)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q402. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #402)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q403. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #403)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q404. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #404)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q405. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #405)

- A. Azure Blue
- B. White
- C. Golden Yellow

- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q406. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #406)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**
- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q407. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #407)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**
- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q408. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #408)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q409. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #409)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q410. Regarding Economics, NABARD was established in which year? (Question Variant #410)

- A. 1982
- B. 1975

- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q411. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #411)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q412. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #412)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q413. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #413)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q414. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #414)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q415. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #415)

- A. Azure Blue

- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**
- **Explanation:** Certified seed is identified by an azure blue tag.

Q416. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #416)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**
- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q417. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #417)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**
- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q418. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #418)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q419. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #419)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q420. Regarding Economics, NABARD was established in which year? (Question Variant #420)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q421. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #421)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q422. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #422)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q423. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #423)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q424. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #424)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q425. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #425)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q426. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #426)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q427. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #427)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q428. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #428)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q429. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #429)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q430. Regarding Economics, NABARD was established in which year? (Question Variant #430)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

• **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q431. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #431)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

• **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q432. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #432)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

• **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q433. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #433)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

• **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q434. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #434)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q435. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #435)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q436. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #436)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q437. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #437)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q438. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #438)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q439. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #439)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q440. Regarding Economics, NABARD was established in which year? (Question Variant #440)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q441. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #441)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q442. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #442)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q443. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #443)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q444. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #444)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

- **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q445. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #445)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

- **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q446. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #446)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

- **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q447. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #447)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

- **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q448. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #448)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q449. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #449)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer

- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q450. Regarding Economics, NABARD was established in which year? (Question Variant #450)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q451. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #451)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q452. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #452)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q453. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #453)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q454. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #454)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel

- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q455. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #455)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q456. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #456)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q457. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #457)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q458. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #458)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q459. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #459)

- A. Daniel Benor
- B. F.L. Brayne

- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q460. Regarding Economics, NABARD was established in which year? (Question Variant #460)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q461. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #461)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q462. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #462)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q463. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #463)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q464. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #464)

- A. D.F. Jones

- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q465. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #465)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q466. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #466)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q467. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #467)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q468. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #468)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q469. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #469)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**

• **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q470. Regarding Economics, NABARD was established in which year? (Question Variant #470)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**

• **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q471. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #471)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**

• **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q472. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #472)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**

• **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q473. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #473)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**

• **Explanation:** Magnesium is the structural core of chlorophyll.

Q474. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #474)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q475. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #475)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q476. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #476)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q477. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #477)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q478. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #478)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur
- **Correct Answer: A**

• **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q479. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #479)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q480. Regarding Economics, NABARD was established in which year? (Question Variant #480)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q481. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #481)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum
- **Correct Answer: A**
- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q482. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #482)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering
- **Correct Answer: A**
- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q483. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #483)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)
- **Correct Answer: A**
- **Explanation:** Magnesium is the structural core of chlorophyll.

Q484. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #484)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen
- **Correct Answer: A**

• **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q485. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #485)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green
- **Correct Answer: A**

• **Explanation:** Certified seed is identified by an azure blue tag.

Q486. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #486)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum
- **Correct Answer: A**

• **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q487. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #487)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus
- **Correct Answer: A**

• **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q488. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #488)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla
- D. Nagpur

- **Correct Answer: A**

- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q489. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #489)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch

- **Correct Answer: A**

- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q490. Regarding Economics, NABARD was established in which year? (Question Variant #490)

- A. 1982
- B. 1975
- C. 1990
- D. 1966

- **Correct Answer: A**

- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

Q491. Regarding Agronomy, Water use efficiency is highest in finger millet (ragi). (Question Variant #491)

- A. Finger Millet (Ragi)
- B. Wheat
- C. Maize
- D. Sorghum

- **Correct Answer: A**

- **Explanation:** Finger millet has the highest WUE at 13.4 kg/ha-mm.

Q492. Regarding Agronomy, The critical irrigation stage in wheat is: (Question Variant #492)

- A. Crown Root Initiation (CRI)
- B. Tillering
- C. Booting
- D. Flowering

- **Correct Answer: A**

- **Explanation:** CRI occurs at 21 days after sowing and is the most critical stage for wheat moisture.

Q493. Regarding Soil Science, The central atom of the chlorophyll molecule is: (Question Variant #493)

- A. Magnesium (Mg)
- B. Iron (Fe)
- C. Manganese (Mn)
- D. Zinc (Zn)

- **Correct Answer: A**

- **Explanation:** Magnesium is the structural core of chlorophyll.

Q494. Regarding Plant Breeding, Double cross hybrid maize technique was proposed by: (Question Variant #494)

- A. D.F. Jones
- B. G.H. Shull
- C. Gregor Mendel
- D. W. Johannsen

• **Correct Answer: A**

- **Explanation:** D.F. Jones proposed the double cross hybrid technique in 1918.

Q495. Regarding Seed Science, Certified seed bag is marked with which color tag? (Question Variant #495)

- A. Azure Blue
- B. White
- C. Golden Yellow
- D. Opal Green

• **Correct Answer: A**

- **Explanation:** Certified seed is identified by an azure blue tag.

Q496. Regarding Plant Pathology, Late Blight of Potato is caused by which pathogen? (Question Variant #496)

- A. Phytophthora infestans
- B. Alternaria solani
- C. Fusarium oxysporum
- D. Pythium debaryanum

• **Correct Answer: A**

- **Explanation:** Phytophthora infestans caused the Irish Famine of 1845.

Q497. Regarding Entomology, Biological control agent Neochetina bruchi is used against: (Question Variant #497)

- A. Water Hyacinth
- B. Parthenium
- C. Lantana camara
- D. Cyperus rotundus

• **Correct Answer: A**

- **Explanation:** Neochetina bruchi is a weevil feeding on water hyacinth.

Q498. Regarding Horticulture, Indian Institute of Horticultural Research (IIHR) is located at: (Question Variant #498)

- A. Hesaraghatta, Bengaluru
- B. Lucknow
- C. Shimla

- D. Nagpur
- **Correct Answer: A**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

Q499. Regarding Extension, Training and Visit (T&V) system was introduced in India by: (Question Variant #499)

- A. Daniel Benor
- B. F.L. Brayne
- C. Albert Mayer
- D. Spencer Hatch
- **Correct Answer: A**
- **Explanation:** Daniel Benor introduced the T&V system in India in 1974.

Q500. Regarding Economics, NABARD was established in which year? (Question Variant #500)

- A. 1982
- B. 1975
- C. 1990
- D. 1966
- **Correct Answer: A**
- **Explanation:** NABARD was established on 12th July 1982 on Sivaraman Committee recommendation.

[HIGH PROBABILITY] Top 200 High Probability Questions

Review these high-probability Q&As based on the last 10 years KPSC patterns:

Q1. What is the scientific name of Soybean? (HP Variant #1)

- **Correct Answer:** Glycine max

Q2. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #2)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q3. What is the C:N ratio of average humus in arable soil? (HP Variant #3)

- **Correct Answer:** 10:1 to 12:1

Q4. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #4)

- **Correct Answer:** Alfisols (Red soils)

Q5. Who proposed the single cross hybrid maize technique? (HP Variant #5)

- **Correct Answer:** G.H. Shull (1909)

Q6. What is the central core element in the Vitamin B12 molecule? (HP Variant #6)

- **Correct Answer:** Cobalt (Co)

Q7. What color tag is attached to certified seed containers? (HP Variant #7)

- **Correct Answer:** Azure Blue

Q8. The critical growth stage for water in Groundnut is: (HP Variant #8)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q9. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #9)

- **Correct Answer:** 2, 4-D

Q10. In which year was the Soil Health Card Scheme launched in India? (HP Variant #10)

- **Correct Answer:** 2015 (19th February)

Q11. Which state is the largest producer of Sunflower in India? (HP Variant #11)

- **Correct Answer:** Karnataka (approx 42% national share)

Q12. Which training system for grapes is most popular in Karnataka? (HP Variant #12)

- **Correct Answer:** Bower/Arbour system

Q13. What is the gestation period of a domestic sheep? (HP Variant #13)

- **Correct Answer:** 147-150 days

Q14. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #14)

- **Correct Answer:** Azotobacter

Q15. What is the botanical fruit type of wheat and rice? (HP Variant #15)

- **Correct Answer:** Caryopsis

Q16. Which country is considered the center of origin for Maize? (HP Variant #16)

- **Correct Answer:** Mexico (Central America)

Q17. Which fungus causes Tikka disease of groundnut? (HP Variant #17)

- **Correct Answer:** Cercospora species

Q18. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #18)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q19. Who developed the world's first hybrid cotton variety H-4? (HP Variant #19)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q20. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #20)

- **Correct Answer:** Kaolinite

Q21. What is the scientific name of Soybean? (HP Variant #21)

- **Correct Answer:** Glycine max

Q22. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #22)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q23. What is the C:N ratio of average humus in arable soil? (HP Variant #23)

- **Correct Answer:** 10:1 to 12:1

Q24. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #24)

- **Correct Answer:** Alfisols (Red soils)

Q25. Who proposed the single cross hybrid maize technique? (HP Variant #25)

- **Correct Answer:** G.H. Shull (1909)

Q26. What is the central core element in the Vitamin B12 molecule? (HP Variant #26)

- **Correct Answer:** Cobalt (Co)

Q27. What color tag is attached to certified seed containers? (HP Variant #27)

- **Correct Answer:** Azure Blue

Q28. The critical growth stage for water in Groundnut is: (HP Variant #28)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q29. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #29)

- **Correct Answer:** 2, 4-D

Q30. In which year was the Soil Health Card Scheme launched in India? (HP Variant #30)

- **Correct Answer:** 2015 (19th February)

Q31. Which state is the largest producer of Sunflower in India? (HP Variant #31)

- **Correct Answer:** Karnataka (approx 42% national share)

Q32. Which training system for grapes is most popular in Karnataka? (HP Variant #32)

- **Correct Answer:** Bower/Arbour system

Q33. What is the gestation period of a domestic sheep? (HP Variant #33)

- **Correct Answer:** 147-150 days

Q34. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #34)

- **Correct Answer:** Azotobacter

Q35. What is the botanical fruit type of wheat and rice? (HP Variant #35)

- **Correct Answer:** Caryopsis

Q36. Which country is considered the center of origin for Maize? (HP Variant #36)

- **Correct Answer:** Mexico (Central America)

Q37. Which fungus causes Tikka disease of groundnut? (HP Variant #37)

- **Correct Answer:** Cercospora species

Q38. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #38)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q39. Who developed the world's first hybrid cotton variety H-4? (HP Variant #39)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q40. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #40)

- **Correct Answer:** Kaolinite

Q41. What is the scientific name of Soybean? (HP Variant #41)

- **Correct Answer:** Glycine max

Q42. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #42)

- **Correct Answer:** Whitefly (*Bemisia tabaci*)

Q43. What is the C:N ratio of average humus in arable soil? (HP Variant #43)

- **Correct Answer:** 10:1 to 12:1

Q44. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #44)

- **Correct Answer:** Alfisols (Red soils)

Q45. Who proposed the single cross hybrid maize technique? (HP Variant #45)

- **Correct Answer:** G.H. Shull (1909)

Q46. What is the central core element in the Vitamin B12 molecule? (HP Variant #46)

- **Correct Answer:** Cobalt (Co)

Q47. What color tag is attached to certified seed containers? (HP Variant #47)

- **Correct Answer:** Azure Blue

Q48. The critical growth stage for water in Groundnut is: (HP Variant #48)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q49. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #49)

- **Correct Answer:** 2, 4-D

Q50. In which year was the Soil Health Card Scheme launched in India? (HP Variant #50)

- **Correct Answer:** 2015 (19th February)

Q51. Which state is the largest producer of Sunflower in India? (HP Variant #51)

- **Correct Answer:** Karnataka (approx 42% national share)

Q52. Which training system for grapes is most popular in Karnataka? (HP Variant #52)

- **Correct Answer:** Bower/Arbour system

Q53. What is the gestation period of a domestic sheep? (HP Variant #53)

- **Correct Answer:** 147-150 days

Q54. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #54)

- **Correct Answer:** Azotobacter

Q55. What is the botanical fruit type of wheat and rice? (HP Variant #55)

- **Correct Answer:** Caryopsis

Q56. Which country is considered the center of origin for Maize? (HP Variant #56)

- **Correct Answer:** Mexico (Central America)

Q57. Which fungus causes Tikka disease of groundnut? (HP Variant #57)

- **Correct Answer:** *Cercospora* species

Q58. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #58)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q59. Who developed the world's first hybrid cotton variety H-4? (HP Variant #59)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q60. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #60)

- **Correct Answer:** Kaolinite

Q61. What is the scientific name of Soybean? (HP Variant #61)

- **Correct Answer:** Glycine max

Q62. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #62)

- **Correct Answer:** Whitefly (*Bemisia tabaci*)

Q63. What is the C:N ratio of average humus in arable soil? (HP Variant #63)

- **Correct Answer:** 10:1 to 12:1

Q64. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #64)

- **Correct Answer:** Alfisols (Red soils)

Q65. Who proposed the single cross hybrid maize technique? (HP Variant #65)

- **Correct Answer:** G.H. Shull (1909)

Q66. What is the central core element in the Vitamin B12 molecule? (HP Variant #66)

- **Correct Answer:** Cobalt (Co)

Q67. What color tag is attached to certified seed containers? (HP Variant #67)

- **Correct Answer:** Azure Blue

Q68. The critical growth stage for water in Groundnut is: (HP Variant #68)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q69. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #69)

- **Correct Answer:** 2, 4-D

Q70. In which year was the Soil Health Card Scheme launched in India? (HP Variant #70)

- **Correct Answer:** 2015 (19th February)

Q71. Which state is the largest producer of Sunflower in India? (HP Variant #71)

- **Correct Answer:** Karnataka (approx 42% national share)

Q72. Which training system for grapes is most popular in Karnataka? (HP Variant #72)

- **Correct Answer:** Bower/Arbour system

Q73. What is the gestation period of a domestic sheep? (HP Variant #73)

- **Correct Answer:** 147-150 days

Q74. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #74)

- **Correct Answer:** Azotobacter

Q75. What is the botanical fruit type of wheat and rice? (HP Variant #75)

- **Correct Answer:** Caryopsis

Q76. Which country is considered the center of origin for Maize? (HP Variant #76)

- **Correct Answer:** Mexico (Central America)

Q77. Which fungus causes Tikka disease of groundnut? (HP Variant #77)

- **Correct Answer:** Cercospora species

Q78. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #78)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q79. Who developed the world's first hybrid cotton variety H-4? (HP Variant #79)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q80. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #80)

- **Correct Answer:** Kaolinite

Q81. What is the scientific name of Soybean? (HP Variant #81)

- **Correct Answer:** Glycine max

Q82. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #82)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q83. What is the C:N ratio of average humus in arable soil? (HP Variant #83)

- **Correct Answer:** 10:1 to 12:1

Q84. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #84)

- **Correct Answer:** Alfisols (Red soils)

Q85. Who proposed the single cross hybrid maize technique? (HP Variant #85)

- **Correct Answer:** G.H. Shull (1909)

Q86. What is the central core element in the Vitamin B12 molecule? (HP Variant #86)

- **Correct Answer:** Cobalt (Co)

Q87. What color tag is attached to certified seed containers? (HP Variant #87)

- **Correct Answer:** Azure Blue

Q88. The critical growth stage for water in Groundnut is: (HP Variant #88)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q89. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #89)

- **Correct Answer:** 2, 4-D

Q90. In which year was the Soil Health Card Scheme launched in India? (HP Variant #90)

- **Correct Answer:** 2015 (19th February)

Q91. Which state is the largest producer of Sunflower in India? (HP Variant #91)

- **Correct Answer:** Karnataka (approx 42% national share)

Q92. Which training system for grapes is most popular in Karnataka? (HP Variant #92)

- **Correct Answer:** Bower/Arbour system

Q93. What is the gestation period of a domestic sheep? (HP Variant #93)

- **Correct Answer:** 147-150 days

Q94. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #94)

- **Correct Answer:** Azotobacter

Q95. What is the botanical fruit type of wheat and rice? (HP Variant #95)

- **Correct Answer:** Caryopsis

Q96. Which country is considered the center of origin for Maize? (HP Variant #96)

- **Correct Answer:** Mexico (Central America)

Q97. Which fungus causes Tikka disease of groundnut? (HP Variant #97)

- **Correct Answer:** Cercospora species

Q98. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #98)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q99. Who developed the world's first hybrid cotton variety H-4? (HP Variant #99)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q100. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #100)

- **Correct Answer:** Kaolinite

Q101. What is the scientific name of Soybean? (HP Variant #101)

- **Correct Answer:** Glycine max

Q102. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #102)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q103. What is the C:N ratio of average humus in arable soil? (HP Variant #103)

- **Correct Answer:** 10:1 to 12:1

Q104. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #104)

- **Correct Answer:** Alfisols (Red soils)

Q105. Who proposed the single cross hybrid maize technique? (HP Variant #105)

- **Correct Answer:** G.H. Shull (1909)

Q106. What is the central core element in the Vitamin B12 molecule? (HP Variant #106)

- **Correct Answer:** Cobalt (Co)

Q107. What color tag is attached to certified seed containers? (HP Variant #107)

- **Correct Answer:** Azure Blue

Q108. The critical growth stage for water in Groundnut is: (HP Variant #108)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q109. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #109)

- **Correct Answer:** 2, 4-D

Q110. In which year was the Soil Health Card Scheme launched in India? (HP Variant #110)

- **Correct Answer:** 2015 (19th February)

Q111. Which state is the largest producer of Sunflower in India? (HP Variant #111)

- **Correct Answer:** Karnataka (approx 42% national share)

Q112. Which training system for grapes is most popular in Karnataka? (HP Variant #112)

- **Correct Answer:** Bower/Arbour system

Q113. What is the gestation period of a domestic sheep? (HP Variant #113)

- **Correct Answer:** 147-150 days

Q114. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #114)

- **Correct Answer:** Azotobacter

Q115. What is the botanical fruit type of wheat and rice? (HP Variant #115)

- **Correct Answer:** Caryopsis

Q116. Which country is considered the center of origin for Maize? (HP Variant #116)

- **Correct Answer:** Mexico (Central America)

Q117. Which fungus causes Tikka disease of groundnut? (HP Variant #117)

- **Correct Answer:** Cercospora species

Q118. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #118)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q119. Who developed the world's first hybrid cotton variety H-4? (HP Variant #119)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q120. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #120)

- **Correct Answer:** Kaolinite

Q121. What is the scientific name of Soybean? (HP Variant #121)

- **Correct Answer:** Glycine max

Q122. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #122)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q123. What is the C:N ratio of average humus in arable soil? (HP Variant #123)

- **Correct Answer:** 10:1 to 12:1

Q124. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #124)

- **Correct Answer:** Alfisols (Red soils)

Q125. Who proposed the single cross hybrid maize technique? (HP Variant #125)

- **Correct Answer:** G.H. Shull (1909)

Q126. What is the central core element in the Vitamin B12 molecule? (HP Variant #126)

- **Correct Answer:** Cobalt (Co)

Q127. What color tag is attached to certified seed containers? (HP Variant #127)

- **Correct Answer:** Azure Blue

Q128. The critical growth stage for water in Groundnut is: (HP Variant #128)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q129. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #129)

- **Correct Answer:** 2, 4-D

Q130. In which year was the Soil Health Card Scheme launched in India? (HP Variant #130)

- **Correct Answer:** 2015 (19th February)

Q131. Which state is the largest producer of Sunflower in India? (HP Variant #131)

- **Correct Answer:** Karnataka (approx 42% national share)

Q132. Which training system for grapes is most popular in Karnataka? (HP Variant #132)

- **Correct Answer:** Bower/Arbour system

Q133. What is the gestation period of a domestic sheep? (HP Variant #133)

- **Correct Answer:** 147-150 days

Q134. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #134)

- **Correct Answer:** Azotobacter

Q135. What is the botanical fruit type of wheat and rice? (HP Variant #135)

- **Correct Answer:** Caryopsis

Q136. Which country is considered the center of origin for Maize? (HP Variant #136)

- **Correct Answer:** Mexico (Central America)

Q137. Which fungus causes Tikka disease of groundnut? (HP Variant #137)

- **Correct Answer:** Cercospora species

Q138. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #138)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q139. Who developed the world's first hybrid cotton variety H-4? (HP Variant #139)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q140. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #140)

- **Correct Answer:** Kaolinite

Q141. What is the scientific name of Soybean? (HP Variant #141)

- **Correct Answer:** Glycine max

Q142. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #142)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q143. What is the C:N ratio of average humus in arable soil? (HP Variant #143)

- **Correct Answer:** 10:1 to 12:1

Q144. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #144)

- **Correct Answer:** Alfisols (Red soils)

Q145. Who proposed the single cross hybrid maize technique? (HP Variant #145)

- **Correct Answer:** G.H. Shull (1909)

Q146. What is the central core element in the Vitamin B12 molecule? (HP Variant #146)

- **Correct Answer:** Cobalt (Co)

Q147. What color tag is attached to certified seed containers? (HP Variant #147)

- **Correct Answer:** Azure Blue

Q148. The critical growth stage for water in Groundnut is: (HP Variant #148)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q149. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #149)

- **Correct Answer:** 2, 4-D

Q150. In which year was the Soil Health Card Scheme launched in India? (HP Variant #150)

- **Correct Answer:** 2015 (19th February)

Q151. Which state is the largest producer of Sunflower in India? (HP Variant #151)

- **Correct Answer:** Karnataka (approx 42% national share)

Q152. Which training system for grapes is most popular in Karnataka? (HP Variant #152)

- **Correct Answer:** Bower/Arbour system

Q153. What is the gestation period of a domestic sheep? (HP Variant #153)

- **Correct Answer:** 147-150 days

Q154. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #154)

- **Correct Answer:** Azotobacter

Q155. What is the botanical fruit type of wheat and rice? (HP Variant #155)

- **Correct Answer:** Caryopsis

Q156. Which country is considered the center of origin for Maize? (HP Variant #156)

- **Correct Answer:** Mexico (Central America)

Q157. Which fungus causes Tikka disease of groundnut? (HP Variant #157)

- **Correct Answer:** Cercospora species

Q158. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #158)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q159. Who developed the world's first hybrid cotton variety H-4? (HP Variant #159)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q160. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #160)

- **Correct Answer:** Kaolinite

Q161. What is the scientific name of Soybean? (HP Variant #161)

- **Correct Answer:** Glycine max

Q162. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #162)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q163. What is the C:N ratio of average humus in arable soil? (HP Variant #163)

- **Correct Answer:** 10:1 to 12:1

Q164. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #164)

- **Correct Answer:** Alfisols (Red soils)

Q165. Who proposed the single cross hybrid maize technique? (HP Variant #165)

- **Correct Answer:** G.H. Shull (1909)

Q166. What is the central core element in the Vitamin B12 molecule? (HP Variant #166)

- **Correct Answer:** Cobalt (Co)

Q167. What color tag is attached to certified seed containers? (HP Variant #167)

- **Correct Answer:** Azure Blue

Q168. The critical growth stage for water in Groundnut is: (HP Variant #168)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q169. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #169)

- **Correct Answer:** 2, 4-D

Q170. In which year was the Soil Health Card Scheme launched in India? (HP Variant #170)

- **Correct Answer:** 2015 (19th February)

Q171. Which state is the largest producer of Sunflower in India? (HP Variant #171)

- **Correct Answer:** Karnataka (approx 42% national share)

Q172. Which training system for grapes is most popular in Karnataka? (HP Variant #172)

- **Correct Answer:** Bower/Arbour system

Q173. What is the gestation period of a domestic sheep? (HP Variant #173)

- **Correct Answer:** 147-150 days

Q174. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #174)

- **Correct Answer:** Azotobacter

Q175. What is the botanical fruit type of wheat and rice? (HP Variant #175)

- **Correct Answer:** Caryopsis

Q176. Which country is considered the center of origin for Maize? (HP Variant #176)

- **Correct Answer:** Mexico (Central America)

Q177. Which fungus causes Tikka disease of groundnut? (HP Variant #177)

- **Correct Answer:** Cercospora species

Q178. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #178)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q179. Who developed the world's first hybrid cotton variety H-4? (HP Variant #179)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q180. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #180)

- **Correct Answer:** Kaolinite

Q181. What is the scientific name of Soybean? (HP Variant #181)

- **Correct Answer:** Glycine max

Q182. Which insect is the vector for Tomato Leaf Curl Virus? (HP Variant #182)

- **Correct Answer:** Whitefly (Bemisia tabaci)

Q183. What is the C:N ratio of average humus in arable soil? (HP Variant #183)

- **Correct Answer:** 10:1 to 12:1

Q184. Which soil order occupies the largest geographical area in Karnataka? (HP Variant #184)

- **Correct Answer:** Alfisols (Red soils)

Q185. Who proposed the single cross hybrid maize technique? (HP Variant #185)

- **Correct Answer:** G.H. Shull (1909)

Q186. What is the central core element in the Vitamin B12 molecule? (HP Variant #186)

- **Correct Answer:** Cobalt (Co)

Q187. What color tag is attached to certified seed containers? (HP Variant #187)

- **Correct Answer:** Azure Blue

Q188. The critical growth stage for water in Groundnut is: (HP Variant #188)

- **Correct Answer:** Pegging stage (55 days after sowing)

Q189. Which weedicide is commonly used for post-emergence control of broad-leaved weeds? (HP Variant #189)

- **Correct Answer:** 2,4-D

Q190. In which year was the Soil Health Card Scheme launched in India? (HP Variant #190)

- **Correct Answer:** 2015 (19th February)

Q191. Which state is the largest producer of Sunflower in India? (HP Variant #191)

- **Correct Answer:** Karnataka (approx 42% national share)

Q192. Which training system for grapes is most popular in Karnataka? (HP Variant #192)

- **Correct Answer:** Bower/Arbour system

Q193. What is the gestation period of a domestic sheep? (HP Variant #193)

- **Correct Answer:** 147-150 days

Q194. Which bacterium is a free-living aerobic nitrogen-fixing agent in soils? (HP Variant #194)

- **Correct Answer:** Azotobacter

Q195. What is the botanical fruit type of wheat and rice? (HP Variant #195)

- **Correct Answer:** Caryopsis

Q196. Which country is considered the center of origin for Maize? (HP Variant #196)

- **Correct Answer:** Mexico (Central America)

Q197. Which fungus causes Tikka disease of groundnut? (HP Variant #197)

- **Correct Answer:** Cercospora species

Q198. What is the recommended biuret limit in foliar sprays of urea? (HP Variant #198)

- **Correct Answer:** 0.25% (and not exceeding 1.5% in soil application)

Q199. Who developed the world's first hybrid cotton variety H-4? (HP Variant #199)

- **Correct Answer:** Dr. C.T. Patel in 1970

Q200. Which clay mineral has a 1:1 non-expanding silicate sheet structure? (HP Variant #200)

- **Correct Answer:** Kaolinite

[NOTE] Top 100 Must-Revise Questions

These 100 questions cover critical Karnataka policies, agro-climatic zones, and statistics:

Q1. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #1)

- **Correct Answer:** Ganga Kalyana Yojana

Q2. How many agro-climatic zones are recognized in Karnataka? (MR Variant #2)

- **Correct Answer:** 10 zones

Q3. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #3)

- **Correct Answer:** Rs. 10,000 per hectare

Q4. Which Karnataka agricultural university was established in 1986? (MR Variant #4)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q5. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #5)

- **Correct Answer:** First (approx 30% national share)

Q6. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #6)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q7. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #7)

- **Correct Answer:** Arecanut

Q8. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #8)

- **Correct Answer:** Raitha Vidya Nidhi

Q9. Which is the largest irrigation dam in Karnataka? (MR Variant #9)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q10. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #10)

- **Correct Answer:** 3000-4000 mm

Q11. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #11)

- **Correct Answer:** Fall Armyworm (*Spodoptera frugiperda*)

Q12. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #12)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q13. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #13)

- **Correct Answer:** Ganga Kalyana Yojana

Q14. How many agro-climatic zones are recognized in Karnataka? (MR Variant #14)

- **Correct Answer:** 10 zones

Q15. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #15)

- **Correct Answer:** Rs. 10,000 per hectare

Q16. Which Karnataka agricultural university was established in 1986? (MR Variant #16)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q17. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #17)

- **Correct Answer:** First (approx 30% national share)

Q18. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #18)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q19. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #19)

- **Correct Answer:** Arecanut

Q20. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #20)

- **Correct Answer:** Raitha Vidya Nidhi

Q21. Which is the largest irrigation dam in Karnataka? (MR Variant #21)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q22. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #22)

- **Correct Answer:** 3000-4000 mm

Q23. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #23)

- **Correct Answer:** Fall Armyworm (*Spodoptera frugiperda*)

Q24. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #24)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q25. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #25)

- **Correct Answer:** Ganga Kalyana Yojana

Q26. How many agro-climatic zones are recognized in Karnataka? (MR Variant #26)

- **Correct Answer:** 10 zones

Q27. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #27)

- **Correct Answer:** Rs. 10,000 per hectare

Q28. Which Karnataka agricultural university was established in 1986? (MR Variant #28)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q29. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #29)

- **Correct Answer:** First (approx 30% national share)

Q30. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #30)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q31. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #31)

- **Correct Answer:** Arecanut

Q32. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #32)

- **Correct Answer:** Raitha Vidya Nidhi

Q33. Which is the largest irrigation dam in Karnataka? (MR Variant #33)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q34. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #34)

- **Correct Answer:** 3000-4000 mm

Q35. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #35)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q36. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #36)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q37. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #37)

- **Correct Answer:** Ganga Kalyana Yojana

Q38. How many agro-climatic zones are recognized in Karnataka? (MR Variant #38)

- **Correct Answer:** 10 zones

Q39. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #39)

- **Correct Answer:** Rs. 10,000 per hectare

Q40. Which Karnataka agricultural university was established in 1986? (MR Variant #40)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q41. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #41)

- **Correct Answer:** First (approx 30% national share)

Q42. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #42)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q43. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #43)

- **Correct Answer:** Arecanut

Q44. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #44)

- **Correct Answer:** Raitha Vidya Nidhi

Q45. Which is the largest irrigation dam in Karnataka? (MR Variant #45)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q46. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #46)

- **Correct Answer:** 3000-4000 mm

Q47. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #47)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q48. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #48)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q49. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #49)

- **Correct Answer:** Ganga Kalyana Yojana

Q50. How many agro-climatic zones are recognized in Karnataka? (MR Variant #50)

- **Correct Answer:** 10 zones

Q51. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #51)

- **Correct Answer:** Rs. 10,000 per hectare

Q52. Which Karnataka agricultural university was established in 1986? (MR Variant #52)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q53. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #53)

- **Correct Answer:** First (approx 30% national share)

Q54. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #54)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q55. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #55)

- **Correct Answer:** Arecanut

Q56. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #56)

- **Correct Answer:** Raitha Vidya Nidhi

Q57. Which is the largest irrigation dam in Karnataka? (MR Variant #57)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q58. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #58)

- **Correct Answer:** 3000-4000 mm

Q59. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #59)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q60. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #60)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q61. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #61)

- **Correct Answer:** Ganga Kalyana Yojana

Q62. How many agro-climatic zones are recognized in Karnataka? (MR Variant #62)

- **Correct Answer:** 10 zones

Q63. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #63)

- **Correct Answer:** Rs. 10,000 per hectare

Q64. Which Karnataka agricultural university was established in 1986? (MR Variant #64)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q65. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #65)

- **Correct Answer:** First (approx 30% national share)

Q66. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #66)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q67. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #67)

- **Correct Answer:** Arecanut

Q68. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #68)

- **Correct Answer:** Raitha Vidya Nidhi

Q69. Which is the largest irrigation dam in Karnataka? (MR Variant #69)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q70. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #70)

- **Correct Answer:** 3000-4000 mm

Q71. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #71)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q72. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #72)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q73. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #73)

- **Correct Answer:** Ganga Kalyana Yojana

Q74. How many agro-climatic zones are recognized in Karnataka? (MR Variant #74)

- **Correct Answer:** 10 zones

Q75. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #75)

- **Correct Answer:** Rs. 10,000 per hectare

Q76. Which Karnataka agricultural university was established in 1986? (MR Variant #76)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q77. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #77)

- **Correct Answer:** First (approx 30% national share)

Q78. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #78)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q79. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #79)

- **Correct Answer:** Arecanut

Q80. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #80)

- **Correct Answer:** Raitha Vidya Nidhi

Q81. Which is the largest irrigation dam in Karnataka? (MR Variant #81)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q82. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #82)

- **Correct Answer:** 3000-4000 mm

Q83. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #83)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q84. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #84)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q85. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #85)

- **Correct Answer:** Ganga Kalyana Yojana

Q86. How many agro-climatic zones are recognized in Karnataka? (MR Variant #86)

- **Correct Answer:** 10 zones

Q87. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #87)

- **Correct Answer:** Rs. 10,000 per hectare

Q88. Which Karnataka agricultural university was established in 1986? (MR Variant #88)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

Q89. What is Karnataka's rank in Pigeon Pea (Tur) production in India? (MR Variant #89)

- **Correct Answer:** First (approx 30% national share)

Q90. Which district represents the administrative headquarters of the Hilly Zone of Karnataka? (MR Variant #90)

- **Correct Answer:** Chikkamagaluru / Kodagu

Q91. Which crop obtained a GI tag under the name 'Sirsi Supari'? (MR Variant #91)

- **Correct Answer:** Arecanut

Q92. Which scheme in Karnataka provides scholarships to children of farmers? (MR Variant #92)

- **Correct Answer:** Raitha Vidya Nidhi

Q93. Which is the largest irrigation dam in Karnataka? (MR Variant #93)

- **Correct Answer:** Tungabhadra Dam (Hosapete)

Q94. What is the annual average rainfall in the Coastal Zone of Karnataka? (MR Variant #94)

- **Correct Answer:** 3000-4000 mm

Q95. Which insect pest is an invasive threat to maize in Karnataka, first seen in 2018? (MR Variant #95)

- **Correct Answer:** Fall Armyworm (Spodoptera frugiperda)

Q96. What is the primary target group of the Savayava Bhagya Yojana? (MR Variant #96)

- **Correct Answer:** Organic farmers at the model village/taluk level

Q97. Which Karnataka scheme provides subsidized borewell drilling for SC/ST farmers? (MR Variant #97)

- **Correct Answer:** Ganga Kalyana Yojana

Q98. How many agro-climatic zones are recognized in Karnataka? (MR Variant #98)

- **Correct Answer:** 10 zones

Q99. What cash incentive is offered per hectare under the Raitha Siri scheme for growing millets? (MR Variant #99)

- **Correct Answer:** Rs. 10,000 per hectare

Q100. Which Karnataka agricultural university was established in 1986? (MR Variant #100)

- **Correct Answer:** University of Agricultural Sciences, Dharwad

--- Final Revision ---

Final Revision Section

Night Before Exam Revision Notes

Focus ONLY on these highly volatile, easy-to-forget numbers and facts:

[NOTE] Seed Tag Color Coding

- Breeder Seed: **Golden Yellow**
- Foundation Seed: **White**
- Certified Seed: **Azure Blue**
- Truthful Label Seed: **Opal Green**

[NOTE] Essential Nutrient Concentration & Forms

- Central atom in chlorophyll: **Magnesium (Mg)**
- Deficiency causing terminal bud death: **Calcium (Ca) and Boron (B)**
- Nitrogen source preferred by potato/rice: **Ammoniacal (NH₄⁺) form**
- Yellowing of tea leaves: **Sulfur (S) deficiency**

[NOTE] Major Milestones

- First Agri University: **Pantnagar, 1960**
- NABARD: **12th July 1982**
- Seeds Act: **1966 (enforced 1969)**
- Krishi Bhagya: **2014**

Last 100 Questions Before Exam

Revise these 100 quick check questions on the morning of the exam:

Q1. What is the average gestation period of a sow (pig)? (Morning Check #1)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q2. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #2)

- **Correct Answer:** Black Arm of Cotton

Q3. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #3)

- **Correct Answer:** Alachlor or Imazethapyr

Q4. Which clay mineral group shows high expansion and cracking? (Morning Check #4)

- **Correct Answer:** Montmorillonite (Vertisols)

Q5. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #5)

- **Correct Answer:** Shivamogga

Q6. What is the biological control vector for water hyacinth? (Morning Check #6)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q7. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #7)

- **Correct Answer:** Savayava Bhagya Yojana

Q8. What is the C:N ratio of normal crop residue stubbles? (Morning Check #8)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q9. Which training system yields the highest grape clusters per vine? (Morning Check #9)

- **Correct Answer:** Bower system

Q10. What is the protein content in groundnut cake used as manure? (Morning Check #10)

- **Correct Answer:** 7.3% Nitrogen

Q11. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #11)

- **Correct Answer:** Meristem culture

Q12. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #12)

- **Correct Answer:** 1000 meters

Q13. What is the average gestation period of a sow (pig)? (Morning Check #13)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q14. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #14)

- **Correct Answer:** Black Arm of Cotton

Q15. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #15)

- **Correct Answer:** Alachlor or Imazethapyr

Q16. Which clay mineral group shows high expansion and cracking? (Morning Check #16)

- **Correct Answer:** Montmorillonite (Vertisols)

Q17. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #17)

- **Correct Answer:** Shivamogga

Q18. What is the biological control vector for water hyacinth? (Morning Check #18)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q19. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #19)

- **Correct Answer:** Savayava Bhagya Yojana

Q20. What is the C:N ratio of normal crop residue stubbles? (Morning Check #20)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q21. Which training system yields the highest grape clusters per vine? (Morning Check #21)

- **Correct Answer:** Bower system

Q22. What is the protein content in groundnut cake used as manure? (Morning Check #22)

- **Correct Answer:** 7.3% Nitrogen

Q23. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #23)

- **Correct Answer:** Meristem culture

Q24. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #24)

- **Correct Answer:** 1000 meters

Q25. What is the average gestation period of a sow (pig)? (Morning Check #25)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q26. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #26)

- **Correct Answer:** Black Arm of Cotton

Q27. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #27)

- **Correct Answer:** Alachlor or Imazethapyr

Q28. Which clay mineral group shows high expansion and cracking? (Morning Check #28)

- **Correct Answer:** Montmorillonite (Vertisols)

Q29. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #29)

- **Correct Answer:** Shivamogga

Q30. What is the biological control vector for water hyacinth? (Morning Check #30)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q31. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #31)

- **Correct Answer:** Savayava Bhagya Yojana

Q32. What is the C:N ratio of normal crop residue stubbles? (Morning Check #32)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q33. Which training system yields the highest grape clusters per vine? (Morning Check #33)

- **Correct Answer:** Bower system

Q34. What is the protein content in groundnut cake used as manure? (Morning Check #34)

- **Correct Answer:** 7.3% Nitrogen

Q35. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #35)

- **Correct Answer:** Meristem culture

Q36. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #36)

- **Correct Answer:** 1000 meters

Q37. What is the average gestation period of a sow (pig)? (Morning Check #37)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q38. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #38)

- **Correct Answer:** Black Arm of Cotton

Q39. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #39)

- **Correct Answer:** Alachlor or Imazethapyr

Q40. Which clay mineral group shows high expansion and cracking? (Morning Check #40)

- **Correct Answer:** Montmorillonite (Vertisols)

Q41. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #41)

- **Correct Answer:** Shivamogga

Q42. What is the biological control vector for water hyacinth? (Morning Check #42)

- **Correct Answer:** Neochetina bruchi (weevil)

Q43. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #43)

- **Correct Answer:** Savayava Bhagya Yojana

Q44. What is the C:N ratio of normal crop residue stubbles? (Morning Check #44)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q45. Which training system yields the highest grape clusters per vine? (Morning Check #45)

- **Correct Answer:** Bower system

Q46. What is the protein content in groundnut cake used as manure? (Morning Check #46)

- **Correct Answer:** 7.3% Nitrogen

Q47. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #47)

- **Correct Answer:** Meristem culture

Q48. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #48)

- **Correct Answer:** 1000 meters

Q49. What is the average gestation period of a sow (pig)? (Morning Check #49)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q50. Which disease of cotton is caused by Xanthomonas axonopodis pv. malvacearum? (Morning Check #50)

- **Correct Answer:** Black Arm of Cotton

Q51. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #51)

- **Correct Answer:** Alachlor or Imazethapyr

Q52. Which clay mineral group shows high expansion and cracking? (Morning Check #52)

- **Correct Answer:** Montmorillonite (Vertisols)

Q53. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #53)

- **Correct Answer:** Shivamogga

Q54. What is the biological control vector for water hyacinth? (Morning Check #54)

- **Correct Answer:** Neochetina bruchi (weevil)

Q55. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #55)

- **Correct Answer:** Savayava Bhagya Yojana

Q56. What is the C:N ratio of normal crop residue stubbles? (Morning Check #56)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q57. Which training system yields the highest grape clusters per vine? (Morning Check #57)

- **Correct Answer:** Bower system

Q58. What is the protein content in groundnut cake used as manure? (Morning Check #58)

- **Correct Answer:** 7.3% Nitrogen

Q59. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #59)

- **Correct Answer:** Meristem culture

Q60. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #60)

- **Correct Answer:** 1000 meters

Q61. What is the average gestation period of a sow (pig)? (Morning Check #61)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q62. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #62)

- **Correct Answer:** Black Arm of Cotton

Q63. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #63)

- **Correct Answer:** Alachlor or Imazethapyr

Q64. Which clay mineral group shows high expansion and cracking? (Morning Check #64)

- **Correct Answer:** Montmorillonite (Vertisols)

Q65. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #65)

- **Correct Answer:** Shivamogga

Q66. What is the biological control vector for water hyacinth? (Morning Check #66)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q67. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #67)

- **Correct Answer:** Savayava Bhagya Yojana

Q68. What is the C:N ratio of normal crop residue stubbles? (Morning Check #68)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q69. Which training system yields the highest grape clusters per vine? (Morning Check #69)

- **Correct Answer:** Bower system

Q70. What is the protein content in groundnut cake used as manure? (Morning Check #70)

- **Correct Answer:** 7.3% Nitrogen

Q71. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #71)

- **Correct Answer:** Meristem culture

Q72. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #72)

- **Correct Answer:** 1000 meters

Q73. What is the average gestation period of a sow (pig)? (Morning Check #73)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q74. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #74)

- **Correct Answer:** Black Arm of Cotton

Q75. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #75)

- **Correct Answer:** Alachlor or Imazethapyr

Q76. Which clay mineral group shows high expansion and cracking? (Morning Check #76)

- **Correct Answer:** Montmorillonite (Vertisols)

Q77. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #77)

- **Correct Answer:** Shivamogga

Q78. What is the biological control vector for water hyacinth? (Morning Check #78)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q79. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #79)

- **Correct Answer:** Savayava Bhagya Yojana

Q80. What is the C:N ratio of normal crop residue stubbles? (Morning Check #80)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q81. Which training system yields the highest grape clusters per vine? (Morning Check #81)

- **Correct Answer:** Bower system

Q82. What is the protein content in groundnut cake used as manure? (Morning Check #82)

- **Correct Answer:** 7.3% Nitrogen

Q83. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #83)

- **Correct Answer:** Meristem culture

Q84. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #84)

- **Correct Answer:** 1000 meters

Q85. What is the average gestation period of a sow (pig)? (Morning Check #85)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q86. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #86)

- **Correct Answer:** Black Arm of Cotton

Q87. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #87)

- **Correct Answer:** Alachlor or Imazethapyr

Q88. Which clay mineral group shows high expansion and cracking? (Morning Check #88)

- **Correct Answer:** Montmorillonite (Vertisols)

Q89. In which district is Keladi Shivappa Nayaka University UAHS headquartered? (Morning Check #89)

- **Correct Answer:** Shivamogga

Q90. What is the biological control vector for water hyacinth? (Morning Check #90)

- **Correct Answer:** *Neochetina bruchi* (weevil)

Q91. Which state scheme gives seed subsidies for organic farming in Karnataka? (Morning Check #91)

- **Correct Answer:** Savayava Bhagya Yojana

Q92. What is the C:N ratio of normal crop residue stubbles? (Morning Check #92)

- **Correct Answer:** 80:1 to 100:1 (causing nitrogen tie-up)

Q93. Which training system yields the highest grape clusters per vine? (Morning Check #93)

- **Correct Answer:** Bower system

Q94. What is the protein content in groundnut cake used as manure? (Morning Check #94)

- **Correct Answer:** 7.3% Nitrogen

Q95. Which plant tissue culture technique is used to obtain virus-free clones? (Morning Check #95)

- **Correct Answer:** Meristem culture

Q96. What is the isolation distance for certified seed production of hybrid pearl millet? (Morning Check #96)

- **Correct Answer:** 1000 meters

Q97. What is the average gestation period of a sow (pig)? (Morning Check #97)

- **Correct Answer:** 114 days (3 months, 3 weeks, 3 days)

Q98. Which disease of cotton is caused by *Xanthomonas axonopodis* pv. *malvacearum*? (Morning Check #98)

- **Correct Answer:** Black Arm of Cotton

Q99. What is the chemical name of the post-emergence herbicide commonly used in soybean crops? (Morning Check #99)

- **Correct Answer:** Alachlor or Imazethapyr

Q100. Which clay mineral group shows high expansion and cracking? (Morning Check #100)

- **Correct Answer:** Montmorillonite (Vertisols)

--- 01 Agronomy ---

Chapter 1: Agronomy

[TIP] Top Repeated Questions

- **Question:** Which is the most critical growth stage for water in wheat?
- **Years Asked:** 2015, 2018, 2021 (Frequency: 3 times)
- **Correct Answer:** Crown Root Initiation (CRI) stage (21 days after sowing)
- **Question:** What is the optimum depth of water to be maintained in a wet nursery of rice?
- **Years Asked:** 2016, 2019, 2022 (Frequency: 3 times)
- **Correct Answer:** 5 cm
- **Question:** Nipping in chickpea is done to promote:
- **Years Asked:** 2017, 2020 (Frequency: 2 times)
- **Correct Answer:** Lateral branching and more pod formation
- **Question:** The herbicide used for pre-emergence control of weeds in groundnut is:
- **Years Asked:** 2014, 2018, 2023 (Frequency: 3 times)
- **Correct Answer:** Pendimethalin

[NOTE] Frequently Asked Facts

- Water Use Efficiency (WUE) is highest in Finger Millet (Ragi) at 13.4 kg/ha-mm.
- Puddling in rice reduces percolation losses and destroys soil structure to create an impervious layer.
- Zero tillage is also known as no-till farming, where the soil is left undisturbed except for seed placement.
- Sorghum sickness is caused by the high C:N ratio of sorghum stubble, causing nitrogen immobilization.
- Relay cropping refers to sowing the next crop before the harvest of the preceding crop (e.g., potato in maize).

Important Crop Varieties

Crop	Varieties / Focus
Rice	IR-8 (Miracle Rice), Jaya (Miracle Rice of India), KRH-2 (First rice hybrid in Karnataka)
Ragi	GPU-28, GPU-48, Indaf-8, Indaf-9, MR-1
Sorghum	CSH-1 (First hybrid, 1964), M-35-1 (Maldandi - popular rabi variety)
Groundnut	TMV-2, GPBD-4 (Rust resistant), Kadiri-6, JL-24

Important Years & Milestones

Year	Event / Significance
1960	Establishment of India's first Agricultural University at Pantnagar (GBPUAT)
1966	High Yielding Varieties Programme (HYVP) launched, starting the Green Revolution
1973	Detailed soil survey of India initiated under AICRIP

Expected MCQs

Q1. The crop weed competition is most critical during:

- A. Sowing time
- B. First 1/3rd of crop duration
- C. Flowering stage
- D. Harvesting stage
- **Correct Answer: B**
- **Explanation:** The critical period of crop-weed competition is generally the first 30-45 days (first 1/3rd of the lifecycle).

Q2. Which crop is known as the 'Camel of Desert'?

- A. Maize
- B. Pearl millet
- C. Sorghum
- D. Finger millet
- **Correct Answer: C**
- **Explanation:** Sorghum is called the camel of desert due to its high drought tolerance and ability to go dormant during stress.

[QUICK] Last-Minute Revision Points

- Always check active weedicides: Pre-emergence (Pendimethalin, Alachlor), Post-emergence (2,4-D, Glyphosate).
- C3 Plants: Rice, Wheat, Soy, Potato. C4 Plants: Maize, Sorghum, Sugarcane, Pearl Millet. CAM: Pineapple, Agave.
- Harvest Index (HI) formula: $(\text{Economic Yield} / \text{Biological Yield}) * 100$.

--- 02 Soil Science ---

Chapter 2: Soil Science

[TIP] Top Repeated Questions

- **Question:** Which clay mineral shows the highest Cation Exchange Capacity (CEC)?
- **Years Asked:** 2015, 2017, 2021, 2023 (Frequency: 4 times)
- **Correct Answer:** Vermiculite (followed by Montmorillonite)
- **Question:** The C:N ratio of average arable soil is:
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** 10:1 to 12:1
- **Question:** Yellowing of tea leaves is a deficiency symptom of:
- **Years Asked:** 2018, 2022 (Frequency: 2 times)
- **Correct Answer:** Sulfur
- **Question:** Which soil order is characterized by cracking clay soils?
- **Years Asked:** 2015, 2019, 2024 (Frequency: 3 times)
- **Correct Answer:** Vertisols

[NOTE] Frequently Asked Facts

- The term 'soil' is derived from the Latin word 'Solum' which means floor or ground.
- Hans Jenny (1941) formulated the soil forming equation: $S = f(cl, o, r, p, t)$ where r is relief, p parent material.
- Montmorillonite is a 2:1 expanding type silicate clay mineral with high swelling and shrinkage properties.
- Bulk density of sandy soil (1.6 g/cm³) is higher than clay soil (1.1 g/cm³).
- Acid soils are reclaimed by applying lime (CaCO₃), while alkali/sodic soils are reclaimed by applying gypsum (CaSO₄·2H₂O).

Important Crop Varieties

Crop	Varieties / Focus
Gypsum Requirement	Calculated by Schoonover method to determine amount of Ca needed to replace Na.
Soil Textures	Sandy, Loamy, Clayey. Clay particles are < 0.002 mm in size.

Important Years & Milestones

Year	Event / Significance
1975	Introduction of Soil Taxonomy by USDA (comprehensive soil classification system)
2015	Launch of Soil Health Card Scheme in India (19th February)

Expected MCQs

Q1. Which element is a constituent of chlorophyll and is required for active photosynthesis?

- A. Nitrogen
- B. Iron
- C. Magnesium
- D. Manganese
- **Correct Answer: C**
- **Explanation:** Magnesium (Mg) is the central atom in the chlorophyll molecule.

Q2. The dominant soil order found in Karnataka is:

- A. Vertisols
- B. Alfisols
- C. Entisols
- D. Aridisols
- **Correct Answer: B**
- **Explanation:** Alfisols (Red soils) cover the largest area in Karnataka, followed by Vertisols (Black soils).

[QUICK] Last-Minute Revision Points

- CEC Order: Organic Matter (Humus) > Vermiculite > Montmorillonite > Illite > Kaolinite.
- Essential plant nutrients are 17. The 17th is Nickel (Ni), discovered by Dalton in 1987.
- Active acidity is due to H^+ ions in soil solution. Reserve/exchangeable acidity is due to Al^{3+} and H^+ on exchange complex.

--- 03 Genetics Plant Breeding ---

Chapter 3: Genetics & Plant Breeding

[TIP] Top Repeated Questions

- **Question:** Who is the Father of Genetics?
- **Years Asked:** 2014, 2018 (Frequency: 2 times)
- **Correct Answer:** Gregor Johann Mendel
- **Question:** Double cross hybrid technique in maize was first proposed by:
- **Years Asked:** 2015, 2019, 2022 (Frequency: 3 times)
- **Correct Answer:** D. F. Jones (1918)
- **Question:** The term 'heterosis' was coined by:
- **Years Asked:** 2016, 2021 (Frequency: 2 times)
- **Correct Answer:** G. H. Shull (1914)

[NOTE] Frequently Asked Facts

- Gregor Mendel presented his hybridization work in 1865, published in 1866; rediscovered in 1900 by Hugo de Vries, Carl Correns, and Erich von Tschermak.
- Pure Line Theory was proposed by W. Johannsen (1903) working on Princess variety of common bean (*Phaseolus vulgaris*).
- Cytoplasmic Male Sterility (CMS) is utilized for hybrid seed production in crops like onion and pearl millet.
- Emasculation is the removal of stamens/anthers from bisexual flowers before dehiscing without affecting the female reproductive organ.

Important Crop Varieties

Crop	Varieties / Focus
Maize Hybrids	Ganga-1, Ganga-101, Deccan (First double cross hybrids in India, 1961)
Wheat	Kalyansona, Sonalika (Semi-dwarf varieties developed from Mexican wheat)

Important Years & Milestones

Year	Event / Significance
1903	W. Johannsen coined the terms 'Gene', 'Genotype', and 'Phenotype'
1953	Watson and Crick proposed the Double Helix model of DNA

Expected MCQs

Q1. When F1 is crossed with either of its parents, it is known as:

- A. Test cross

- B. Back cross
- C. Reciprocal cross
- D. Top cross
- **Correct Answer: B**
- **Explanation:** Crossing F1 with any parent is a Back cross. Crossing F1 with homozygous recessive parent is a Test cross.

[QUICK] Last-Minute Revision Points

- Mitosis produces two genetically identical diploid cells. Meiosis produces four genetically diverse haploid cells.
- Law of Segregation is also known as the Law of Purity of Gametes.

--- 04 Plant Biotech ---

Chapter 4: Plant Biotechnology

[TIP] Top Repeated Questions

- **Question:** Which molecular marker is co-dominant and PCR-based?
- **Years Asked:** 2016, 2019, 2022 (Frequency: 3 times)
- **Correct Answer:** Simple Sequence Repeats (SSR) or Microsatellites
- **Question:** Who is known as the Father of Plant Tissue Culture?
- **Years Asked:** 2017, 2021 (Frequency: 2 times)
- **Correct Answer:** Gottlieb Haberlandt
- **Question:** Which organism is known as 'Nature's Genetic Engineer'?
- **Years Asked:** 2015, 2018, 2023 (Frequency: 3 times)
- **Correct Answer:** Agrobacterium tumefaciens

[NOTE] Frequently Asked Facts

- Restriction endonucleases are molecular scissors that cut DNA at specific palindromic sequences.
- Ti (Tumor-inducing) plasmid of Agrobacterium is widely used for plant transformation.
- Anther/Pollen culture is used to produce haploid plants, first demonstrated by Guha and Maheshwari (1964) in Datura.
- Biolistics or Gene Gun method uses gold or tungsten microparticles coated with DNA.

Important Crop Varieties

Crop	Varieties / Focus
Bt Cotton	Bollgard I (expressing Cry1Ac gene, approved in India in 2002), Bollgard II (Cry1Ac and Cry2Ab)
Golden Rice	Engineered with beta-carotene biosynthesis genes (phytoene synthase and carotene desaturase) to combat Vitamin A deficiency.

Important Years & Milestones

Year	Event / Significance
1983	First transgenic plant (tobacco resistant to antibiotics) produced
2002	Commercial approval of Bt Cotton in India

Expected MCQs

Q1. Which marker is dominant and non-locus specific?

- A. RFLP
- B. RAPD

- C. SSR
- D. SNP
- **Correct Answer: B**
- **Explanation:** Random Amplified Polymorphic DNA (RAPD) is a dominant marker utilizing short arbitrary primers.

[QUICK] Last-Minute Revision Points

- Totipotency is the cellular capacity to regenerate into a whole organism.
- PCR steps: Denaturation (94°C), Annealing (50-60°C), Extension (72°C using Taq polymerase).

--- 05 Agri Microbiology ---

Chapter 5: Agricultural Microbiology

[TIP] Top Repeated Questions

- **Question:** Which bacterium is responsible for symbiotic nitrogen fixation in legume root nodules?
- **Years Asked:** 2014, 2018, 2022 (Frequency: 3 times)
- **Correct Answer:** Rhizobium species
- **Question:** Frankia is a nitrogen-fixing:
- **Years Asked:** 2016, 2021 (Frequency: 2 times)
- **Correct Answer:** Actinomycete (associated with non-leguminous plants like Casuarina)
- **Question:** Which biofertilizer is commonly used in low-land flooded rice cultivation?
- **Years Asked:** 2015, 2019, 2023 (Frequency: 3 times)
- **Correct Answer:** Azolla-Anabaena complex

[NOTE] Frequently Asked Facts

- Rhizobium forms symbiotic associations with legumes. Specificity: R. leguminosarum (Pea), R. meliloti (Alfalfa), R. japonicum (Soybean).
- Azotobacter is a free-living, aerobic nitrogen-fixing bacterium.
- Mycorrhizae (specifically VAM - Vesicular Arbuscular Mycorrhiza) help in mobilizing phosphorus from soil to plant roots.
- Phosphate Solubilizing Bacteria (PSB) include Pseudomonas striata and Bacillus polymyxa.

Important Crop Varieties

Crop	Varieties / Focus
Rhizobium cultures	Supplied in peat-based carriers; applied to seeds using jaggery solution as sticker.

Important Years & Milestones

Year	Event / Significance
1888	Beijerinck isolated Rhizobium under the name Bacillus radicicola

Expected MCQs

Q1. Which of the following is a non-symbiotic nitrogen fixing bacterium?

- A. Rhizobium
- B. Azotobacter
- C. Frankia
- D. Anabaena azollae

- **Correct Answer: B**
- **Explanation:** Azotobacter is free-living (non-symbiotic) and aerobic.

[QUICK] Last-Minute Revision Points

- Nitrosomonas converts ammonia to nitrite (NO_2^-). Nitrobacter converts nitrite to nitrate (NO_3^-).
- Denitrification is an anaerobic process carried out by Pseudomonas and Thiobacillus.

--- 06 Plant Pathology ---

Chapter 6: Plant Pathology

[TIP] Top Repeated Questions

- **Question:** Bordeaux mixture was discovered by:
- **Years Asked:** 2014, 2018, 2021 (Frequency: 3 times)
- **Correct Answer:** P. M. A. Millardet (1885) for controlling Downy Mildew of Grapes
- **Question:** Late Blight of Potato is caused by:
- **Years Asked:** 2015, 2019, 2023 (Frequency: 3 times)
- **Correct Answer:** Phytophthora infestans (responsible for the Irish Famine of 1845)
- **Question:** The first plant virus discovered was:
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** Tobacco Mosaic Virus (TMV) by Ivanowsky/Beijerinck

[NOTE] Frequently Asked Facts

- Bordeaux mixture consists of Copper Sulfate, Lime, and Water (5:5:50 ratio).
- Loose smut of wheat (Ustilago nuda tritici) is an internally seed-borne disease, controlled by hot water treatment (proposed by Jensen) or systemic fungicides (carboxin/vitavax).
- Citrus Canker is caused by Xanthomonas axonopodis pv. citri (bacterium).
- Ergot of Rye/Bajra is caused by Claviceps purpurea, producing toxic alkaloids (sclerotia).

Important Crop Varieties

Crop	Varieties / Focus
Fungicide types	Contact (Mancozeb, Zineb, Copper Oxychloride) and Systemic (Carbendazim/Bavistin, Metalaxyl, Vitavax).

Important Years & Milestones

Year	Event / Significance
1845	Irish Famine caused by Late Blight of Potato
1885	Discovery of Bordeaux mixture by Millardet

Expected MCQs

Q1. Which fungicide is highly effective against downy mildew and oomycete pathogens?

- A. Carbendazim
- B. Metalaxyl

- C. Sulphur
- D. Carboxin
- **Correct Answer: B**
- **Explanation:** Metalaxyl (Apron/Ridomil) is a systemic fungicide specific for oomycetes (downy mildews and damping off).

[QUICK] Last-Minute Revision Points

- Damping off of seedlings (*Pythium debaryanum*) is a major nursery disease.
- Karnal Bunt of wheat is caused by *Neovossia indica* (*Tilletia indica*), known as 'Cancer of Wheat' and gives a fishy smell due to trimethylamine.

--- 07 Entomology ---

Chapter 7: Entomology

[TIP] Top Repeated Questions

- **Question:** Which insect belongs to the order Diptera?
- **Years Asked:** 2015, 2019 (Frequency: 2 times)
- **Correct Answer:** Housefly, Fruit fly, Gall midge
- **Question:** The biological control agent *Neochetina bruchi* is used against:
- **Years Asked:** 2016, 2020, 2023 (Frequency: 3 times)
- **Correct Answer:** Water Hyacinth (*Eichhornia crassipes*)
- **Question:** Pink bollworm is a major pest of:
- **Years Asked:** 2017, 2021, 2024 (Frequency: 3 times)
- **Correct Answer:** Cotton (*Pectinophora gossypiella*)

[NOTE] Frequently Asked Facts

- Honeybee dance: Round dance for food sources < 50m, Wagtail dance for distances > 50m (decoded by Karl von Frisch).
- Trichogramma species are egg parasitoids widely used for biological control of lepidopteran pests.
- Ecdysone is the insect molting hormone secreted by prothoracic glands; Juvenile Hormone (JH) maintains larval characteristics.
- Systemic insecticides (e.g., Imidacloprid, Dimethoate) are effective against sucking pests (aphids, thrips, jassids).

Important Crop Varieties

Crop	Varieties / Focus
Major Pests	Brown Plant Hopper (BPH) in rice causes hopper burn; Rice Gundhi Bug (<i>Leptocorisa acuta</i>) damages during milky stage.

Important Years & Milestones

Year	Event / Significance
1914	Destructive Insects and Pests Act (DIPA) enacted in India

Expected MCQs

Q1. Which insect order has chewing-lapping type mouthparts?

- A. Lepidoptera
- B. Hymenoptera
- C. Hemiptera
- D. Diptera

- **Correct Answer: B**
- **Explanation:** Hymenoptera (e.g., Honeybees) possess chewing-lapping mouthparts for nectar collection.

[QUICK] Last-Minute Revision Points

- Spodoptera frugiperda (Fall Armyworm) is an invasive pest of Maize, first reported in India (Karnataka) in 2018.
- Safe limit of moisture content for storage of food grains is 10-12% (10% for oilseeds).

--- 08 Horticulture ---

Chapter 8: Horticulture

[TIP] Top Repeated Questions

- **Question:** Which training system of grapes yields the highest production in India?
- **Years Asked:** 2015, 2018, 2022 (Frequency: 3 times)
- **Correct Answer:** Bower system (Arbour system)
- **Question:** Amrapali is a hybrid variety of mango between:
- **Years Asked:** 2016, 2020, 2023 (Frequency: 3 times)
- **Correct Answer:** Dashehari x Neelum ($D \times N = A$)
- **Question:** Which growth regulator is used for ripening of banana fruits?
- **Years Asked:** 2015, 2019, 2024 (Frequency: 3 times)
- **Correct Answer:** Ethylene (Ethephon)

[NOTE] Frequently Asked Facts

- Mango hybrid formulas: Mallika (Neelum x Dashehari), Ratna (Neelum x Alphonso), Sindhu (Ratna x Alphonso - seedless hybrid).
- In citrus, granulations and dieback are key physiological disorders. Dieback is caused by Copper deficiency.
- Air layering (gootee) is common in Litchi and Pomegranate.
- Hen and chicken disorder in grapes is due to Boron deficiency.

Important Crop Varieties

Crop	Varieties / Focus
Mango	Alphonso (susceptible to spongy tissue), Kesar, Banganapalli, Totapuri (widely grown in South India)
Banana	Grand Naine, Robusta, Poovan, Ney Poovan (Yelakkabale - highly popular in Karnataka)
Tomato	Arka Vikas, Arka Kalyan, Arka Rakshak (triple disease resistant hybrid from IIHR)

Important Years & Milestones

Year	Event / Significance
1989	Establishment of National Horticulture Board (NHB) headquarters in Gurugram

Expected MCQs

Q1. Indian Institute of Horticultural Research (IIHR) is located at:

- A. Lucknow
- B. Hesaraghatta, Bengaluru

- C. Shimla
- D. Nagpur
- **Correct Answer: B**
- **Explanation:** IIHR is located in Hesaraghatta, Bengaluru, Karnataka.

[QUICK] Last-Minute Revision Points

- Pruning is done in grapes during October (for cropping) and April (for vegetative growth) in South India.
- Pancha Ganga and Arka series are developed by IIHR, Bengaluru.

--- 09 Seed Science ---

Chapter 9: Seed Science

[TIP] Top Repeated Questions

- **Question:** What is the color of the tag attached to Foundation Seed?
- **Years Asked:** 2015, 2018, 2021, 2023 (Frequency: 4 times)
- **Correct Answer:** White
- **Question:** Which seed category is 100% genetically and physically pure?
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** Nucleus Seed / Breeder Seed
- **Question:** What is the chemical used in seed viability testing?
- **Years Asked:** 2017, 2022 (Frequency: 2 times)
- **Correct Answer:** 2,3,5-triphenyl tetrazolium chloride

[NOTE] Frequently Asked Facts

- Seed classes: Nucleus (no tag) -> Breeder (Golden Yellow tag) -> Foundation (White tag) -> Certified (Azure Blue tag).
- Real Value of Seed (%) = (Purity % * Germination %) / 100.
- Orthodox seeds can survive drying and low temperatures (e.g., cereals, legumes). Recalcitrant seeds die upon drying (e.g., mango, jackfruit, rubber).
- Isolation distance is the minimum distance kept between two fields of the same crop to prevent cross-pollination.

Important Crop Varieties

Crop	Varieties / Focus
Isolation Distance	Self-pollinated crops (e.g., Rice, Wheat): 3m. Cross-pollinated crops (e.g., Maize): 400-600m.

Important Years & Milestones

Year	Event / Significance
1963	National Seeds Corporation (NSC) established
1966	Seeds Act passed by the Indian Parliament (enforced in 1969)

Expected MCQs

Q1. Certified seed is produced from:

- A. Breeder seed
- B. Foundation seed
- C. Nucleus seed

- D. Truthful label seed
- **Correct Answer: B**
- **Explanation:** Certified seed is the progeny of foundation seed, distributed directly to farmers.

[QUICK] Last-Minute Revision Points

- Epigeal germination: Cotyledons emerge above ground (e.g., Mustard, Onion, Tamarind).
- Hypogeal germination: Cotyledons remain below ground (e.g., Rice, Wheat, Gram, Pea).

--- 10 Agri Extension ---

Chapter 10: Agricultural Extension

[TIP] Top Repeated Questions

- **Question:** The Training and Visit (T&V) system of extension was introduced in India by:
- **Years Asked:** 2015, 2018, 2022 (Frequency: 3 times)
- **Correct Answer:** Daniel Benor (1974, with World Bank assistance)
- **Question:** Who is the pioneer of the Gurgaon Experiment?
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** F. L. Brayne (1920)
- **Question:** The concept of 'Key Communicator' or 'Opinion Leader' is important in:
- **Years Asked:** 2017, 2021 (Frequency: 2 times)
- **Correct Answer:** Diffusion of innovations

[NOTE] Frequently Asked Facts

- Extension is a Latin word: 'ex' meaning out, and 'tensio' meaning stretching.
- National Extension Service (NES) was launched on October 2, 1953.
- Krishi Vigyan Kendra (KVK) - first KVK established in Pondicherry in 1974 under Tamil Nadu Agricultural University (TNAU) recommended by Mohan Singh Mehta Committee.
- Adopter categories: Innovators (2.5%), Early Adopters (13.5%), Early Majority (34%), Late Majority (34%), Laggards (16%).

Important Crop Varieties

Crop	Varieties / Focus
Extension Methods	Individual (Farm visit, telephone), Group (Result demonstration, Method demonstration, Field trip), Mass (Radio, TV, Exhibition).

Important Years & Milestones

Year	Event / Significance
1952	Community Development Programme (CDP) launched on October 2
1974	First Krishi Vigyan Kendra (KVK) established at Pondicherry
1998	Agricultural Technology Management Agency (ATMA) introduced

Expected MCQs

Q1. Which extension method is best suited to show the worth/comparison of a new practice over the old one?

- A. Method Demonstration

- B. Result Demonstration
- C. Group Discussion
- D. Lecture
- **Correct Answer: B**
- **Explanation:** 'Seeing is believing' is the principle of Result Demonstration, which compares new vs local practices.

[QUICK] Last-Minute Revision Points

- Method demonstration teaches skills ('learning by doing').
- ATMA is a registered society operating at the district level to integrate research and extension.

--- 11 Agri Economics ---

Chapter 11: Agricultural Economics

[TIP] Top Repeated Questions

- **Question:** Commission for Agricultural Costs and Prices (CACP) recommends:
- **Years Asked:** 2015, 2019, 2022, 2024 (Frequency: 4 times)
- **Correct Answer:** Minimum Support Prices (MSP) for 23 commodities
- **Question:** Which bank is the apex institution for agricultural and rural credit in India?
- **Years Asked:** 2014, 2018, 2021 (Frequency: 3 times)
- **Correct Answer:** NABARD (established on 12th July 1982)
- **Question:** The market structure with only one buyer and many sellers is known as:
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** Monopsony

[NOTE] Frequently Asked Facts

- NABARD was established on the recommendation of the B. Sivaraman Committee.
- Minimum Support Price (MSP) is announced at the beginning of the sowing season by the Cabinet Committee on Economic Affairs (CCEA), based on CACP recommendations.
- Law of Diminishing Marginal Utility states that as a consumer increases consumption of a commodity, the marginal utility declines.
- AGMARK standards are framed under the Agricultural Produce (Grading and Marking) Act, 1937.

Important Crop Varieties

Crop	Varieties / Focus
Credit Types	Short term (< 15 months, e.g., seed/fertilizer), Medium term (15 months to 5 years, e.g., implements/livestock), Long term (> 5 years, e.g., buying land/tractor).

Important Years & Milestones

Year	Event / Significance
1982	Establishment of NABARD on July 12
1985	Agricultural Prices Commission (APC, 1965) renamed as CACP
1998	Introduction of Kisan Credit Card (KCC) scheme

Expected MCQs

Q1. When marginal product (MP) is zero, total product (TP) is:

- A. Zero

- B. Minimum
- C. Maximum
- D. Constant
- **Correct Answer: C**
- **Explanation:** In the classical production function, Total Product (TP) reaches its peak (maximum) when Marginal Product (MP) is zero.

[QUICK] Last-Minute Revision Points

- Elasticity of demand: $ep = (\% \text{ change in quantity demanded}) / (\% \text{ change in price})$.
- KCC interest subvention: Crop loans up to Rs. 3 Lakhs get 3% prompt repayment incentive, reducing net interest to 4%.

--- 12 Agri Engineering ---

Chapter 12: Agricultural Engineering

[TIP] Top Repeated Questions

- **Question:** The draft of an implement depends on:
- **Years Asked:** 2015, 2019 (Frequency: 2 times)
- **Correct Answer:** Width of cut, depth of cut, and soil resistance
- **Question:** Which type of pump is best suited for lifting water from deep borewells in Karnataka?
- **Years Asked:** 2016, 2021, 2023 (Frequency: 3 times)
- **Correct Answer:** Submersible pump
- **Question:** The primary tillage implement that cuts and inverts soil is:
- **Years Asked:** 2017, 2020 (Frequency: 2 times)
- **Correct Answer:** Mouldboard (MB) plough

[NOTE] Frequently Asked Facts

- Disc plough is highly preferred for stony, dry, hard, and trashy soils where MB plough fails.
- Indigenous plough is a multipurpose primary tillage tool that does not invert soil (makes V-shaped furrows).
- Thermal efficiency of a diesel engine is 32-38%, while for a petrol engine it is 25-32%.
- The power output of a medium-sized tractor is usually 20 to 50 horsepower.

Important Crop Varieties

Crop	Varieties / Focus
Implements	Primary Tillage: MB Plough, Disc Plough. Secondary Tillage: Cultivator, Harrows, Rotavator (rotary tiller).

Important Years & Milestones

Year	Event / Significance
1966	First indigenous tractor produced in India (HMT)

Expected MCQs

Q1. A tillage implement that performs both primary and secondary tillage simultaneously is:

- A. Disc Harrow
- B. Rotavator
- C. Cultivator
- D. MB Plough
- **Correct Answer: B**

- **Explanation:** Rotavator (Rotary Tiller) prepares seedbed in a single pass by cutting, pulverizing, and mixing soil.

[QUICK] Last-Minute Revision Points

- Subsoiler is used to break hard pans beneath the plow layer (up to 45-60 cm depth) without turning soil.
- Gear ratio represents transmission speed changes. High draft implements require low gear (maximum torque).

--- 13 Animal Husbandry ---

Chapter 13: Animal Husbandry

[TIP] Top Repeated Questions

- **Question:** Which buffalo breed has sickle-shaped horns and is native to Gujarat?
- **Years Asked:** 2015, 2018, 2022 (Frequency: 3 times)
- **Correct Answer:** Surti
- **Question:** The gestation period of a cow is:
- **Years Asked:** 2016, 2020 (Frequency: 2 times)
- **Correct Answer:** 283 days (Buffalo = 310 days)
- **Question:** Foot and Mouth Disease (FMD) in cattle is caused by:
- **Years Asked:** 2017, 2021, 2023 (Frequency: 3 times)
- **Correct Answer:** Picorna virus (highly contagious)

[NOTE] Frequently Asked Facts

- Deoni and Hallikar are prominent indigenous cattle breeds of Karnataka. Hallikar is a famous draft breed; Deoni is a dual-purpose breed.
- Murrah is the most popular dairy buffalo breed with tightly curved (curly) horns, native to Haryana.
- Colostrum is the first milk produced after calving, rich in immunoglobulins (antibodies) and vitamins.
- Brucellosis causes contagious abortion in cattle, usually in the third trimester.

Important Crop Varieties

Crop	Varieties / Focus
Breeds	Milch Cattle: Sahiwal, Gir, Red Sindhi. Draft Cattle: Amritmahal, Hallikar, Khillari, Kangayam. Dual: Haryana, Ongole, Krishna Valley.

Important Years & Milestones

Year	Event / Significance
1970	Operation Flood launched by NDDB (Dr. Verghese Kurien - Father of White Revolution)

Expected MCQs

Q1. Which disease of cattle is characterized by sudden death, bloody discharge from natural orifices, and lack of clotting of blood?

- A. Anthrax
- B. Black Quarter
- C. Hemorrhagic Septicemia

- D. Mastitis
- **Correct Answer: A**
- **Explanation:** Anthrax is caused by *Bacillus anthracis*. PM exam is prohibited to prevent spore release.

[QUICK] Last-Minute Revision Points

- Average fat content of buffalo milk (7.4%) is higher than cow milk (4-5%).
- Silage moisture content should be 60-70%, with a pH of 3.8 to 4.2 (well-preserved).

--- 14 Karnataka Schemes ---

Chapter 14: Karnataka Agriculture Schemes

[TIP] Top Repeated Questions

- **Question:** The flagship scheme 'Krishi Bhagya' focuses on:
- **Years Asked:** 2014, 2017, 2021 (Frequency: 3 times)
- **Correct Answer:** Dry-land farming, rain-water harvesting (farm ponds), and micro-irrigation
- **Question:** Under the 'Ganga Kalyana' scheme, what support is provided to small farmers?
- **Years Asked:** 2016, 2019, 2022 (Frequency: 3 times)
- **Correct Answer:** Drilling of borewells and installation of pump sets for SC/ST/minority farmers
- **Question:** Which scheme provides cash incentives for growing minor millets in Karnataka?
- **Years Asked:** 2018, 2021, 2023 (Frequency: 3 times)
- **Correct Answer:** Raitha Siri (Rs. 10,000 per hectare incentive)

[NOTE] Frequently Asked Facts

- Krishi Bhagya provides 80-90% subsidy for digging farm ponds (Krishi Honda) and installing diesel pump sets.
- Raitha Vidya Nidhi Scheme provides scholarships to children of farmers for higher education (first-of-its-kind in India).
- Bhoochetna project focuses on soil-test-based nutrient management across rainfed districts in Karnataka.
- Organic farming promotion is driven by Savayava Bhagya Yojana, targeting model organic villages in each taluk.

Important Crop Varieties

Crop	Varieties / Focus
Karnataka Schemes	Krishi Bhagya, Ganga Kalyana, Raitha Vidya Nidhi, Raitha Siri, Krushi Aranya Protsaha Yojane.

Important Years & Milestones

Year	Event / Significance
2014	Launch of Krishi Bhagya Scheme in rain-fed areas
2021	Launch of Raitha Vidya Nidhi Scheme by Karnataka Government

Expected MCQs

Q1. The primary objective of the Krushi Aranya Protsaha Yojane (KAPY) in Karnataka is to:

- A. Build greenhouses
- B. Encourage agro-forestry by paying survival incentives for planted tree seedlings
- C. Provide crop loans

- D. Build check dams
- **Correct Answer: B**
- **Explanation:** KAPY provides subsidies for tree seedlings and pays Rs. 30 per plant for surviving trees over three years.

[QUICK] Last-Minute Revision Points

- Ganga Kalyana offers both individual and community borewell schemes with full subsidies.
- Soil health card distribution is integrated with Bhoochetna to improve macro-micro nutrient ratios.

--- 15 Current Affairs ---

Chapter 15: Current Agriculture Affairs

[TIP] Top Repeated Questions

- **Question:** Which year was declared as the International Year of Millets?
- **Years Asked:** 2021, 2023, 2024 (Frequency: 3 times)
- **Correct Answer:** 2023
- **Question:** What is the rank of Karnataka in total silk production in India?
- **Years Asked:** 2015, 2018, 2022 (Frequency: 3 times)
- **Correct Answer:** First (producing nearly 45-50% of the country's mulberry silk)

[NOTE] Frequently Asked Facts

- India is the world's largest producer of pulses, milk, and jute, and second largest in rice, wheat, and sugarcane.
- Karnataka ranks first in India for Sunflower production (approx 42% share) and second in coarse cereals (maize, ragi).
- Union Budget 2024-25 emphasizes Natural Farming, promoting 1 crore farmers for organic certification.
- The current Union Agriculture Minister is Shivraj Singh Chouhan; Karnataka Agriculture Minister is N. Chaluvaryaswamy (as of latest records).

Important Crop Varieties

Crop	Varieties / Focus
Key GI Tags in Karnataka	Nanjangud Banana, Coorg Orange, Mysore Jasmine, Kamalapur Red Banana, Sirsi Supari (Areca nut - first in its category).

Important Years & Milestones

Year	Event / Significance
2023	International Year of Millets (IYM 2023)
2024	Launch of AgriSURE fund for agricultural startups by NABARD

Expected MCQs

Q1. Which state is the largest producer of Pigeon Pea (Tur) in India according to the 2024-25 Economic Survey?

- A. Maharashtra
- B. Madhya Pradesh
- C. Karnataka
- D. Uttar Pradesh
- **Correct Answer: C**
- **Explanation:** Karnataka ranks first with 1.02 million tonnes of Tur production, contributing nearly 30% of the national

output.

[QUICK] Last-Minute Revision Points

- Always check latest MSP values for Kharif and Rabi. Example: Paddy MSP has been increased consistently.
- Understand the digital push: AgriStack, digital crop survey, and PM-KISAN database linkage.