REF. BY

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

© 022 - 3090 0000 / 4125 2525 ■ wellness@thyrocare.com ⊕ www.thyrocare.com

# **REPORT**

NAME : MANJU GOLAS (55Y/F)

: WELLNESS - DSA

TEST ASKED: AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	
ANTI CCP (ACCP)	E.L.I.S.A	0.62	OD Ratio	
Reference Range :				

Negative: < 0.80 Equivocal: 0.80 - 1.20 Positive: > 1.20

Clinical Significance :

Anti-Cyclic-Citrullinated-Peptide (Anti-CCP) Antibodies hold promise for early and more accurate detection of Rheumatoid Arthritis before the disease proceeds into an irreversible damage.

Analytical Specifications:

Anti-Cyclic-Citrullinated-Peptide (Anti-CCP) antibodies are detected using a solid phase enzyme immuno assay having an analytical sensitivity of 1.0 U/ml. No cross reactivity to other auto antigen is found. Sensitivity of the method is 68% and specificity is 92%.

**Method:** SOLID PHASE CAPTURE ENZYME IMMUNOASSAY

ANTI NUCLEAR ANTIBODIES (ANA)

E.L.I.S.A

0.35

**OD Ratio** 

Reference Range:

Negative < 0.80Equivocal 0.8 - 1.20Positive > 1.20

Method: SOLID PHASE ENZYME IMMUNOASSAY

Please correlate with clinical conditions.

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type

**Labcode** : 210602373/DSA00

: SERUM

**Barcode** : 71358513

SPA.

Dr.Durgaprasad N Agrawal MD

· ·

Dr.Caesar Sengupta MD

Page : 1 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS
AMYLASE	PHOTOMETRY	55	U/L

Reference Range :-

Adults: 30-118 U/L

### Interpretation:

Lipemic Sera (Hypertriglyceridemia) may contain inhibitors, Which falsely depress results. About 20% of patients with Acute Pancreatitis have abnormal lipids. Normal serum amylase may occur in Pancreatitis, Especially relapsing and chronic pancreatitis. Moderate increases may be reported in normal pregnancy.

### Clinical Significance:

Causes of high Serum Amylase include Acute Pancreatitis, Pancreatic Pseudocyst, Pancreatic Ascites, Pancreatic Abscess, Neoplasm in or adjacent to Pancreas, Trauma to Pancreas, and common Duct Stones. Nonpancreatic Causes include inflammatory salivary lesions (Eg, Mumps), Perforated Peptic Ulcer, Intestinal Obstruction, Biliary Tract Disease, Peritonitis, Acute Appendicitis, Diabetic Ketoacidosis, and Extrapancreatic Carcinomas. Amylase levels more than 25-fold the upper limit of normal are often found when metastatic tumors produce Ectopic Amylase.

### Specifications:

Sample Type

**Barcode** 

Precision: Within run %CV has been recorded 1% and between run %CV of 1.5%. Analytical sensitivity (Lower Detection Limit) 3 U/I

### Kit Validation References:

Tietz Nw, Huang WY, Rauh DF ET Al. Laboratory tests in the differential diagnosis of Hyperamylasemia. Clin Chem 1986;32: 301-307

### Please correlate with clinical conditions.

ENZYMATIC PHOTOMETRIC TEST. Method:-

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

Labcode : 210602373/DSA00

: SERUM

: 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 2 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♀ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

NAME : MANJU GOLAS (55Y/F)

REF. BY: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	
APOLIPOPROTEIN - A1 (APO-A1)	NEPHELOMETRY	122	mg/dl	
Reference Range :				
Male : 110 - 205				
Female : 125 - 215				
Method: FULLY AUTOMATED NEPHELOMET	TRY-BN-II			
APOLIPOPROTEIN - B (APO-B)	NEPHELOMETRY	102	mg/dl	
Reference Range :				
Male : 55 - 140				
Female : 55 - 125				
<b>Method:</b> FULLY AUTOMATED NEPHELOMET	TRY-BN-II			
APO B / APO A1 RATIO (APO B/A1)	CALCULATED	0.8	Ratio	
Reference Range :				

Male : 0.35 - 1.0 Female : 0.30 - 0.9

Method: DERIVED FROM SERUM APO A1 AND APO B VALUES

Please correlate with clinical conditions.

Sample Collected on (SCT) : 20 Jun 2016 08:00
Sample Received on (SRT) : 22 Jun 2016 03:43
Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type : SERUM

**Labcode** : 210602373/DSA00

**Barcode** : 71358513

SPA.

Dr.Durgaprasad N Agrawal MD [

Dr.Caesar Sengupta MD

Page : 3 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe,

Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

TEST ASKED: AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS
HIGH SENSITIVITY C-REACTIVE PROTEIN (hs-CRP)	NEPHELOMETRY	2.5	mg/L

Reference Range :-

Risk of coronary Heart Disease.

< 1.00 -- Low Risk 1.00 - 3.00 -- Average Risk > 3.00 -- High Risk

#### Interpretation:

High sensitivity C-reactive protein, when used in conjunction with other clinical laboratory evaluation of acute coronary syndromes, may be useful as an independent marker of prognosis for recurrent events, in patients with stable coronary disease or acute coronary syndromes. hsCRP levels should not be substituted for assessment of traditional cardiovascular risk factors. Patients with persistently unexplained, marked evaluation of hsCRP after repeated testing should be evaluated for non - cardiovascular etiologies

### Clinical significance:

hsCRP measurements may be used as an independent risk marker for the identification of individuals at risk for future cardiovascular disease. Elevated CRP values may be indicative of prognosis of individuals with acute coronary syndromes, and may be useful in the management of such individuals.

Specifications: Precision: Within run %CV has been recorded 3.1% and between run %CV of 2.5%.

### References:

Sample Type

- 1. Chenillot O, Henny J, Steinmez J, et al. High sensitivity C-reactive protein: biological variations and reference limits. Clin Chem Lab Med 2000;38:1003-11.
- 2. Hind CRH, Pepys MB. The role of serum C-reactive protein measurements in clinical practice. Int Med 1984;5:112-51.

Please correlate with clinical conditions.

FULLY AUTOMATED NEPHELOMETRY-BN-II Method:-

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

Labcode : 210602373/DSA00

: SERUM

**Barcode** : 71358513 Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 4 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

TEST ASKED: AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

TEST NAME	TECHNOLOGY	VALUE	UNITS
HOMOCYSTEINE	C.L.I.A	45.8	µmol/L

Reference Range :-

Adult : < 30

### Clinical Significance:

Homocysteine is linked to increased risk of premature coronary artery disease, stroke and thromboembolism. Moreover, alzheimer's disease, osteoporosis, venous thrombosis, schizophrenia, cognitive deficiency and pregnancy complications also elevates Homocysteine levels.

### High Values:

Elevated homocysteine levels might be due to increasing age, genetic traits, drugs, renal dysfunction and dietary deficiency of vitamins or smoking. To lower your homocysteine, eat more green vegetables, stop smoking, alcohol. Folic acid helps lowering elevated levels.

### Caution While Interpretation:

To get most accurate results, it is mandatory to separate serum immediately. In separated serum, homocysteine remains stable for at least 48 hours at room temperature.

### Please correlate with clinical conditions.

Method:-COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

: 22 Jun 2016 10:15

: SERUM

Report Released on (RRT) Sample Type

: 210602373/DSA00 Labcode

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 5 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

: MANJU GOLAS (55Y/F) NAME

REF. BY : WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

ng/ml

- - 474001

TEST NAME		TECHNOLOGY	VALUE	UNITS
TRON		PHOTOMETRY	76.6	ug/dl
IRON	_	PHOTOMETRI	70.0	μg/dl
Reference Range	•			
Male: 70 - 180				
Female : 60 - 180				
Method : FE	ERROZINE METHOD WITHOUT D	EPROTEINIZATION		
<b>TOTAL IRON BIN</b>	DING CAPACITY (TIBC)	PHOTOMETRY	352	μg/dl
Reference Range	:			
Male: 225 - 535 uc	ı/dl Female: 215 - 535 μg/dl			
	PECTROPHOTOMETRIC ASSAY			
% TRANSFERRIN	SATURATION	CALCULATED	21.76	%
Reference Range	:			
13 - 45				
Method: D	ERIVED FROM IRON AND TIBC \	/ALUES		
FERRITIN		C.L.I.A	77.9	ng/ml
Reference Range	:			
Men: 22-322 ng/m	I			
5,				

FULLY AUTOMATED BIDIRECTIONALLY INTERFACED CHEMI LUMINESCENT IMMUNO ASSAY

C.L.I.A

Reference Range: > 5.38 ng/ml

Method:

Women: 10-291 ng/ml

Method:

**FOLIC ACID** 

FULLY AUTOMATED CHEMI LUMINESCENT IMMUNO ASSAY

: 22 Jun 2016 10:15

Please correlate with clinical conditions.

: 20 Jun 2016 08:00 Sample Collected on (SCT) Sample Received on (SRT) : 22 Jun 2016 03:43

: SERUM **Sample Type** 

Report Released on (RRT)

Labcode : 210602373/DSA00

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 6 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♀ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

NAME : MANJU GOLAS (55Y/F)

**REF. BY**: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

### **SAMPLE COLLECTED AT:**

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	NORMAL RANGE
CALCIUM	PHOTOMETRY	8.16	mg/dl	8.8 - 10.6
URIC ACID	PHOTOMETRY	4	mg/dl	Male: 3.5 - 7.2 Female: 2.6 - 6.0
BLOOD UREA NITROGEN (BUN)	PHOTOMETRY	16.34	mg/dl	7.9 - 20
CREATININE - SERUM	PHOTOMETRY	0.51	mg/dl	Male: 0.6 - 1.1 Female: 0.5 - 0.8
<b>BUN / SR.CREATININE RATIO</b>	CALCULATED	32.04	Ratio	9:1 - 23:1
SODIUM	I.S.E	140	mmol/l	136 - 146
CHLORIDE	I.S.E	102.4	mmol/l	98 - 106

#### Please correlate with clinical conditions.

### Method:

CALC - ARSENAZO III METHOD, END POINT.

URIC - Uricase / Peroxidase Method

BUN - KINETIC UV ASSAY.

SCRE - CREATININE ENZYMATIC METHOD

B/CR - DERIVED FROM SERUM BUN AND CREATININE VALUES

SOD - ION SELECTIVE ELECTRODE IN OLYMPUS AU2700

CHL - Ion selective electrode

Sample Collected on (SCT) : 20 Jun 2016 08:00
Sample Received on (SRT) : 22 Jun 2016 03:43
Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type : SERUM

**Labcode** : 210602373/DSA00

**Barcode** : 71358513

SPA.

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 7 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS
LIPASE	PHOTOMETRY	26	U/L

Reference Range :-

Adults: 13 - 60 U/L

Interpretation:

For diagnostic purposes, the results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings like serum amylase. Serum Lipase is usually normal in patients with elevated serum amylase, having peptic ulcer, salivary adenitis, inflammatory bowel disease, intestinal obstruction, and macroamylasemia. Lipemic sera may interfere with results.

Clinical Significance:

High serum Lipase is a specific marker for pancreatitis; after acute pancreatitis the Lipase activity increases within 4–8 hours, reaches a peak after 24 hours and decreases after 8 to 14 days. However, there is no correlation between the Lipase activity determined in serum and the extent of damage to the pancreas.

Specifications:

Precision: Within run %CV has been recorded 1.16 % and between run %CV of 0.65 %. Analytical sensitivity (lower detection limit) 3 U/I

Kit Validation References:

Tietz Nw Et Al. Lipase In Serum - The Elusive Enzyme: An Overview. Clin Chem 1993; 39:746-756.

: 22 Jun 2016 10:15

: SERUM

Please correlate with clinical conditions.

Method:-**ENZYMATIC COLORIMETRIC ASSAY** 

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) Sample Type

Labcode : 210602373/DSA00

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 8 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703





© 022 - 3090 0000 / 4125 2525 

wellness@thyrocare.com 

www.thyrocare.com

# **REPORT**

NAME : MANJU GOLAS (55Y/F)

**REF. BY**: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	NORMAL RANGE
TOTAL CHOLESTEROL	PHOTOMETRY	194	mg/dl	125 - 200
HDL CHOLESTEROL - DIRECT	PHOTOMETRY	43	mg/dl	35-80
LDL CHOLESTEROL - DIRECT	PHOTOMETRY	128	mg/dl	85 - 130
TRIGLYCERIDES	PHOTOMETRY	138	mg/dl	25 - 200
TC/ HDL CHOLESTEROL RATIO	CALCULATED	4.5	Ratio	3.0 - 5.0
LDL / HDL RATIO	CALCULATED	3	Ratio	1.5 - 3.5
VLDL CHOLESTEROL	CALCULATED	27.6	mg/dl	5 - 40
NON-HDL CHOLESTEROL	CALCULATED	151	mg/dl	< 160

Please correlate with clinical conditions.

### Method:

CHOL - CHOD POD METHOD

HCHO - ENZYME SELECTIVE PROTECTION METHOD

LDL - HOMOGENOUS ENZYMATIC COLORIMETRIC ASSAY

TRIG - ENZYMATIC COLORIMETRIC METHOD (GPO) [HIGHLY INFLUENCED BY LEVEL OF FASTING]

TC/H - DERIVED FROM SERUM CHOLESTEROL AND HDL VALUES

LDL/ - Derived from serum HDL and LDL Values

VLDL - DERIVED FROM SERUM TRIGLYCERIDE VALUES

NHDL - Derived from serum Cholesterol and HDL values

### \*REFERENCE RANGES AS PER NCEP ATP III GUIDELINES:

(mg/dl)	HDL	(mg/dl)	LDL	(mg/dl)	TRIGLYCERIDES	(mg/dl)
<200	LOW	<40	OPTIMAL	<100	NORMAL	<150
200-239	HIGH	>60	NEAR OPTIMAL	100-129	BORDERLINE HIGH	150-199
>240			BORDERLINE HIGH	130-159	HIGH	200-499
			HIGH	160-189	VERY HIGH	>500
			VERY HIGH	>190		
	<200 200-239	<200 LOW 200-239 HIGH	<200 LOW <40 200-239 HIGH >60	<200 LOW <40 OPTIMAL  200-239 HIGH >60 NEAR OPTIMAL  >240  BORDERLINE HIGH  HIGH	<200	<200

Alert !!! 10-12 hours fasting is mandatory for lipid parameters. If not, values might fluctuate.

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type : SERUM

**Labcode** : 210602373/DSA00

**Barcode** : 71358513

382

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 9 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

NAME : MANJU GOLAS (55Y/F)

**REF. BY**: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	NORMAL RANGE
ALKALINE PHOSPHATASE	PHOTOMETRY	83.2	U/I	M:53 to 128 - F:42 to 98
BILIRUBIN - TOTAL	PHOTOMETRY	0.34	mg/dl	0.30 - 1.20
BILIRUBIN -DIRECT	PHOTOMETRY	0.11	mg/dl	0 - 0.30
BILIRUBIN (INDIRECT)	CALCULATED	0.24	mg/dl	0 - 0.9
GAMMA GLUTAMYL TRANSFERASE (GGT)	PHOTOMETRY	20.6	U/I	M: 0 to 55 - F:0 to 38
ASPARTATE AMINOTRANSFERASE (SGOT )	PHOTOMETRY	22.1	U/I	M: 0 to 37 - F: 0 to 31
ALANINE TRANSAMINASE (SGPT)	PHOTOMETRY	17.3	U/I	M: 13 to 40 - F: 10 to 28
PROTEIN - TOTAL	PHOTOMETRY	7.5	gm/dl	5.7 - 8.2
ALBUMIN - SERUM	PHOTOMETRY	4.4	gm/dl	3.2 - 4.8
SERUM GLOBULIN	PHOTOMETRY	3.1	gm/dL	2.50-3.40
SERUM ALBUMIN/GLOBULIN RATIO	CALCULATED	1.42	Ratio	0.9 - 2.0

Please correlate with clinical conditions.

### Method:

ALKP - Modified IFCC method

BILT - Vanadate Oxidation

BILD - Vanadate Oxidation

BILI - DERIVED FROM SERUM TOTAL AND DIRECT BILIRUBIN VALUES

GGT - Modified IFCC method

SGOT - IFCC\* WITHOUT PYRIDOXAL PHOSPHATE ACTIVATION

SGPT - IFCC\* WITHOUT PYRIDOXAL PHOSPHATE ACTIVATION

PROT - BIURET METHOD

SALB - ALBUMIN BCG<sup>1</sup>METHOD (COLORIMETRIC ASSAY ENDPOINT)

SEGB - DERIVED FROM SERUM ALBUMIN AND PROTEIN VALUES

A/GR - DERIVED FROM SERUM ALBUMIN AND PROTEIN VALUES

Sample Collected on (SCT) : 20 Jun 2016 08:00

Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type : SERUM

**Labcode** : 210602373/DSA00

**Barcode** : 71358513

3827.

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 10 of 16

# **Thyrocare** D-37/1,TTC MIDC,Turbhe,

Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

TEST ASKED: AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS
LIPOPROTEIN (A) [LP(A)]	NEPHELOMETRY	56.3	mg/dl

Reference Range :-

Adults: < 30.0 mg/dl

Interpretation:

Determination of LPA may be useful to guide management of individuals with a family history of CHD or with existing disease. The levels of LPA in the blood depends on genetic factors; The range of variation in a population is relatively large and hence for diagnostic purpose, results should always be assessed in conjunction with the patient's medical history, clinical examination and other findings.

Specifications:

Precision: Intra Assay (%CV): 3.4 %, Inter Assay (%CV): 2.0 %; Sensitivity: 0.002 gm/l

External Quality Control Program Participation:

College of American Pathologists: General Chemistry and TDM; CAP Number: 7193855-01

: 22 Jun 2016 10:15

Kit Validation References:

Koschinsky ML, Marcovina SM. Lipoprotein A: Structural Implication for Pathophysiology. Int J Clin Lab Res, 1997; 27: 14-23.

Please correlate with clinical conditions.

FULLY AUTOMATED NEPHELOMETRY-BN-II Method:-

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43

Sample Type : SERUM

Report Released on (RRT)

: 210602373/DSA00 Labcode

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 11 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### REPORT

NAME : MANJU GOLAS (55Y/F) REF. BY

: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS
TESTOSTERONE	C.L.I.A	37.3	ng/dl

### Reference Range :-

Adult male 241 - 827 Adult female 14 - 76

Boys

< 1 year 12 - 21 1 - 6 years 3 - 32 3 - 68 7 - 12 years 28 - 1110 13 - 17 years

### Clinical Significance:

Clinical evaluation of serum testosterone, along with serum LH, assists in evaluation of Hypogonadal males. Major causes of lowered testosterone in males include Hypogonadotropic hypogonadism, testicular failure Hyperprolactinema, Hypopituitarism some types of liver and kidney diseases and critical illness.

### Specifications:

Precision: Intra assay (%CV): 2.6 %, Inter assay (%CV): 4.3%; Sensitivity: 10 ng/dl

External quality control program participation:

College of American pathologists: Ligand assay (special) survey; cap number: 7193855-01

# Please correlate with clinical conditions.

Method:-FULLY AUTOMATED BIDIRECTIONALLY INTERFACED CHEMI LUMINESCENT IMMUNO ASSAY

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43 Report Released on (RRT) : 22 Jun 2016 10:15

Sample Type : SERUM

Labcode : 210602373/DSA00

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 12 of 16

REF. BY

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703





Corporate Office: Thyrocare Technologies Limited ♀ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

: MANJU GOLAS (55Y/F) **NAME** 

: WELLNESS - DSA

**TEST ASKED:** AAROGYAM 1.6

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	REFERENCE RANGE
TOTAL TRIIODOTHYRONINE (T3)	C.L.I.A	105	ng/dl	60 - 200
TOTAL THYROXINE (T4)	C.L.I.A	7.8	μg/dl	4.5 - 12.0
THYROID STIMULATING HORMONE (TSH)	C.L.I.A	2.29	μIU/ml	0.30 - 5.5

**Comments:** SUGGESTING THYRONORMALCY Please correlate with clinical conditions.

### Method:

T3 - COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

T4 - COMPETITIVE CHEMI LUMINESCENT IMMUNO ASSAY

TSH - ULTRA SENSITIVE SANDWICH CHEMI LUMINESCENT IMMUNO ASSAY

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:43 Report Released on (RRT) : 22 Jun 2016 10:15

**Sample Type** : SERUM

Labcode : 210602373/DSA00

**Barcode** : 71358513

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page: 13 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited 9 D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

### **REPORT**

NAME : MANJU GOLAS (55Y/F)

REF. BY: WELLNESS - DSA

TEST ASKED: AAROGYAM 1.6

SAMPLE COLLECTED AT :

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

. 87 ng/n	nl
	.87 ng/n

Reference Range:

DEFICIENCY : <20 ng/ml
INSUFFICIENCY : 20-30 ng/ml
SUFFICIENCY : 30-100 ng/ml
TOXICITY : >100 ng/ml

Vitamin D Total test is analyzed on Siemens ADVIA Centaur, standardized against ID-LC/MS/MS, as per Vitamin D Standardization Program (VDSP).

Method: FULLY AUTOMATED CHEMI LUMINESCENT IMMUNO ASSAY

VITAMIN B-12 C.L.I.A 161 pg/ml

Reference Range:

Normal: 211 - 911 pg/ml

### Clinical significance:

Vitamin B12 or cyanocobalamin, is a complex corrinoid compound found exclusively from animal dietary sources, such as meat, eggs and milk. It is critical in normal DNA synthesis, which in turn affects erythrocyte maturation and in the formation of myelin sheath. Vitamin-B12 is used to find out neurological abnormalities and impaired DNA synthesis associated with macrocytic anemias. For diagnostic purpose, results should always be assessed in conjunction with the patients medical history, clinical examination and other findings.

Specifications: Intra assay (%CV):4.0%, Inter assay (%CV):4.4 %; Sensitivity:45 pg/ml

External quality control program participation:

College of American pathologists: ligand assay (general) survey; CAP number: 7193855-01

Kit validation references:

Chen IW, Sperling MI, Heminger IA. Vitamin B12.In: Pesce AJ, Kalpan LA, editors. Methods in clinical chemistry.

St.Louis:CV Mosby,1987.P.569-73.

Method: FULLY AUTOMATED BIDIRECTIONALLY INTERFACED CHEMI LUMINESCENT IMMUNO ASSAY

Please correlate with clinical conditions.

Sample Collected on (SCT) : 20 Jun 2016 08:00

Sample Received on (SRT) : 22 Jun 2016 03:43

Report Released on (RRT) : 22 Jun 2016 10:15

: SERUM

Report Released on (RRT)
Sample Type

**Labcode** : 210602373/DSA00

**Barcode** : 71358513

SPA.

Dr.Durgaprasad N Agrawal MD

(Qu

Dr.Caesar Sengupta MD

Page: 14 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

# **REPORT**

: MANJU GOLAS (55Y/F) NAME REF. BY

: WELLNESS - DSA

**TEST ASKED:** HBA, HEMOGRAM - 6 PART (DIFF)

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

TEST NAME	TECHNOLOGY	VALUE	UNITS	
DIABETES SCREEN (BLOOD)				
HbA1c	H.P.L.C	5.8	%	
Reference Range :				
Below 6.0% - Normal Value				
6.0% - 7.0% - Good Control				
7.0% - 8.0% - Fair Control				
8.0% - 10% - Unsatisfactory Control				
Above 10% - Poor Control				
<b>Method:</b> Fully Automated H.P.L.C. using	Biorad Variant II Turbo			
AVERAGE BLOOD GLUCOSE (ABG)	CALCULATED	120	mg/dl	

**Reference Range:** 

90 - 120 mg/dl : Excellent Control 121 - 150 mg/dl : Good Control 151 - 180 mg/dl : Average Control 181 - 210 mg/dl : Action Suggested > 211 mg/dl : Panic Value

(Note: Average Blood Glucose value is calculated from HBA1c value and it indicates Average Blood Sugar level

over past three months.)

Method: Derived from HBA1c values

Please correlate with clinical conditions.

: 20 Jun 2016 08:00 Sample Collected on (SCT) Sample Received on (SRT) : 22 Jun 2016 03:09 Report Released on (RRT) : 22 Jun 2016 07:40

**Sample Type** 

: EDTA

Labcode : 210644136/DSA00

**Barcode** : 71220882

Dr.Durgaprasad N Agrawal MD Dr.Caesar Sengupta MD

Page: 15 of 16

D-37/1,TTC MIDC,Turbhe, Navi Mumbai-400 703



Corporate Office: Thyrocare Technologies Limited ♥ D-37/3, TTC MIDC, Turbhe, Navi Mumbai - 400703

© 022 - 3090 0000 / 4125 2525 

wellness@thyrocare.com 

www.thyrocare.com

### **REPORT**

NAME : MANJU GOLAS (55Y/F)

**SAMPLE COLLECTED AT:** 

F21, A GOVT QUATER, PARK HOTEL, PADAV, GWALIOR, M.P.

- - 474001

REF. BY: WELLNESS - DSA

**TEST ASKED:** HBA, HEMOGRAM - 6 PART (DIFF)

TEST NAME	VALUE	UNITS	REFERENCE RANGE
TOTAL LEUCOCYTES COUNT	5.55	$X~10^3$ / $\mu L$	4 - 10
NEUTROPHILS	70.3	%	40-80
LYMPHOCYTE PERCENTAGE	25.4	%	M:20-40; F:20-40
MONOCYTES	1.6	%	0-10
EOSINOPHILS	2.2	%	0-6
BASOPHILS	0.2	%	< 2
IMMATURE GRANULOCYTE PERCENTAGE(IG%)	0.3	%	Male: 0-0.5 Female: 0-0.4
NEUTROPHILS - ABSOLUTE COUNT	3.9	X 10 <sup>3</sup> / μL	2.0 - 7.0
LYMPHOCYTES - ABSOLUTE COUNT	1.41	X 10 <sup>3</sup> / μL	1.00 - 3.00
MONOCYTES - ABSOLUTE COUNT	0.09	X 10 <sup>3</sup> / μ	0.20 - 1.00
BASOPHILS - ABSOLUTE COUNT	0.01	X 10 <sup>3</sup> / μ	0.02 - 0.10
EOSINOPHILS - ABSOLUTE COUNT	0.12	$X 10^{3} / \mu L$	0.02 - 0.50
IMMATURE GRANULOCYTES(IG)	0.02	X 10 <sup>3</sup> / μL	0.03
TOTAL RBC	4.71	X 10^6/µl	Male: 4.5-5.5 Female: 3.9-4.8
NUCLEATED RED BLOOD CELLS	Nil	$X 10^{3} / \mu L$	Nil in adults
NUCLEATED RED BLOOD CELLS %	Nil	%	Nil in adults
HEMOGLOBIN	12.3	g/dL	Male : 13-17 Female : 12-15
HEMATOCRIT(PCV)	44.1	%	Male : 40-50 Female : 36-46
MEAN CORPUSCULAR VOLUME(MCV)	93.6	fL	83-101
MEAN CORPUSCULAR HEMOGLOBIN(MCH)	26.1	pq	27-32
MEAN CORP.HEMO.CONC(MCHC)	27.9	g/dL	31.5-34.5
<b>RED CELL DISTRIBUTION WIDTH - SD(RDW-SD)</b>	50.9	fL	39 - 46
RED CELL DISTRIBUTION WIDTH (RDW-CV)	14.6	%	11.6-14
PLATELET DISTRIBUTION WIDTH(PDW)	17.2	fL	9.6-15.2
MEAN PLATELET VOLUME(MPV)	12.7	fL	6.5-12.0
PLATELET COUNT	254	X 10 <sup>3</sup> / μL	150-400
PLATELET TO LARGE CELL RATIO(PLCR)	46.6	%	19.7 - 42.4
PLATELETCRIT(PCT)	0.32	%	0.19 - 0.39

### Please Correlate with clinical conditions.

**Method:** Fully automated bidirectional analyser (6 Part Differential SYSMEX XN-1000)

(This device performs hematology analyses according to the Hydrodynamic Focussing (DC method), Flow Cytometry Method (using a semiconductor laser), and SLS- hemoglobin method)

~~ End of report ~~

Sample Collected on (SCT) : 20 Jun 2016 08:00 Sample Received on (SRT) : 22 Jun 2016 03:09

Report Released on (RRT) : 22 Jun 2016 07:40

Sample Type : EDTA

**Labcode** : 210644136/DSA00

**Barcode** : 71220882

382).

Dr.Durgaprasad N Agrawal MD

Dr.Caesar Sengupta MD

Page : 16 of 16