

**Question 1.** Compute the following quantities.

- 1) Compute  $P'_1(0), P'_2(0), P'_3(0), P'_4(0)$  by differentiating Legendre polynomials.
- 2) Compute  $P'_{2n+1}(0)$  and  $P'_{2n}(0) (n = 0, 1, 2, \dots)$  using Rodrigues' formula.
- 3) Compute  $P_{2n}(0)$  using Rodrigues' formula.

**Solution.** 1)  $P'_1(0) = 1, P'_2(0) = 0, P'_3(0) = -3/2, P'_4(0) = 0$ . 2)  $P'_{2n+1}(0) = (-1)^n \frac{(2n+2)!}{n!(n+1)!2^{2n+1}}$  and  $P'_{2n}(0) = 0$ . 3)  $P_{2n}(0) = (-1)^n \frac{(2n)!}{2^{2n}(n!)^2}$ .

Sol (1)  $P_1(x) = x \Rightarrow P'_1(0) = 1$

$$P_2(x) = \frac{1}{2}(3x^2 - 1) \Rightarrow P'_2(0) = [3x]_{x=0} = 0$$

$$P_3(x) = \frac{1}{2}(5x^3 - 3x) \Rightarrow P'_3(0) = \left[ \frac{1}{2}(15x^2 - 3) \right]_{x=0} = -\frac{3}{2}$$

$$P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3) \Rightarrow P'_4(0) = \left[ \frac{1}{8}(140x^3 - 60x) \right]_{x=0} = 0$$

(2)  $P_{2n}$  is an even function  $\Rightarrow P'_{2n}$  is odd  $\Rightarrow \underline{P'_{2n}(0) = 0}$

$$P'_{2n+1}(x) = \frac{1}{2^{2n+1}(2n+1)!} \frac{d}{dx} \left[ \frac{d^{2n+1}}{dx^{2n+1}} (x^2 - 1)^{2n+1} \right]$$

$$= \frac{1}{2^{2n+1}(2n+1)!} \frac{d^{2n+2}}{dx^{2n+2}} \left( \sum_{k=0}^{2n+1} \binom{2n+1}{k} (-1)^{2n+1-k} x^{2k} \right)$$

evaluated at  $x=0 \Rightarrow$  all term of deg  $> 2n+2$ ,  
and all terms of deg  $< 2n+2$  vanishing

$$\text{So } P'_{2n+1}(0) = \frac{1}{2^{2n+1}(2n+1)!} \frac{d^{2n+2}}{dx^{2n+2}} \binom{2n+1}{n+1} (-1)^n \underline{x^{2n+2}}$$

$$= \frac{1}{2^{2n+1}(2n+1)!} \binom{2n+1}{n+1} (-1)^n (2n+2)!$$

$$= \frac{(-1)^n (2n+2)!}{2^{2n+1}(n+1)!n!}$$

(3) (Same as (2))

$$P_{2n}(0) = \frac{1}{2^{2n}(2n)!} \frac{d^{2n}}{dx^{2n}} \binom{2n}{n} (-1)^{2n-n} x^{2n}$$

$$= \frac{1}{2^{2n}(2n)!} \binom{2n}{n} (-1)^n (2n)! = \underline{\underline{\frac{(-1)^n (2n)!}{2^{2n}(n!)^2}}}$$

**Question 2.** Prove that different form of Legendre equations are equivalent.

- 1)  $(\sin \theta \Theta'(\theta))' + k(k+1) \sin \theta \Theta(\theta) = 0.$
- 2)  $\Theta''(\theta) + \cot \theta \Theta'(\theta) + k(k+1)\Theta(\theta) = 0.$
- 3)  $((1-s^2)y')' + k(k+1)y = 0.$
- 4)  $(1-s^2)y'' - 2sy' + k(k+1)y = 0.$

**Solution.** 1)  $\Leftrightarrow$  2) and 3)  $\Leftrightarrow$  4) can be proved using straightforward computation. 2)  $\Leftrightarrow$  3) can be proved using the change of variable  $s = \cos \theta$ .

1)  $\Leftrightarrow$  2):

$$\text{pf } (\sin(\theta) \varphi'(\theta))' + k(k+1) \sin \theta \varphi(\theta) = 0$$

$$\Leftrightarrow \cos \theta \varphi'(\theta) + \sin \theta \varphi''(\theta) + k(k+1) \sin \theta \varphi(\theta) = 0$$

$$\Leftrightarrow \frac{\cos \theta}{\sin \theta} \varphi'(\theta) + \varphi''(\theta) + k(k+1) \varphi(\theta) = 0$$

$$\Leftrightarrow \varphi''(\theta) + \cot \theta \varphi'(\theta) + k(k+1) \varphi(\theta) = 0$$

3)  $\Leftrightarrow$  4):

$$\text{pf } (1-s^2)y' + k(k+1)y = 0$$

$$\text{Since } ((1-s^2)y')' = \frac{d}{ds}[(1-s^2)y'] = -2sy' + (1-s^2)y''$$

$$\Leftrightarrow (1-s^2)y'' - 2sy' + k(k+1)y = 0$$

2)  $\Leftrightarrow$  3):

$$\text{pf Let } s = \cos \theta \Rightarrow \theta = \arccos s, \quad y(s) = \varphi(\theta) \quad = y'(s)(-\sin \theta)$$

$$\frac{dy}{d\theta} = y'(s)(-\sin \theta), \quad \frac{d^2y}{d\theta^2} = -\cos \theta y'(s) - \sin \theta \frac{dy'(s)}{ds}$$

$$= -\cos \theta y'(s) + \sin^2(\theta) y''(s)$$

$$\text{So eq (2) } \Leftrightarrow y''(s) \sin^2 \theta - y'(s) \cos \theta + \cot \theta (-\sin \theta y'(s)) + k(k+1) y(s) = 0$$

$$\Leftrightarrow y''(s) \sin^2(\theta) - 2y'(s) \cos \theta + k(k+1) y(s) = 0$$

$$\Leftrightarrow (1-s^2)y''(s) - 2sy'(s) + k(k+1)y(s) = 0$$

$$\text{note } ((1-s^2)y')' = \frac{d}{ds}[(1-s^2)y'] = -2sy' + (1-s^2)y''$$

$$\Leftrightarrow (1-s^2)y'' - 2sy' + k(k+1)y = 0$$

**Question 3.** Find the expansion of the following functions in a series of Legendre polynomials.

1) Let  $f(s) = 0$  for  $-1 < s < 0$  and  $f(s) = 1$  for  $0 < s < 1$ .

2) Let  $f(s) = 3s^2 + 4s + 1$ .

**Solution.** 1) Using Rodrigue's formula and integration by parts, we get  $f(s) = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{2k+1}{2k(k+1)} P'_k(0) P_k(s)$ ,  $-1 < s < 1$ . 2)  $f(s) = 2P_0(s) + 4P_1(s) + 2P_2(s)$ . You get this using Rodrigue's formula, but a simpler method is to apply method of undetermined coefficients, set  $f(s) = \sum_{i=0}^2 a_i P_i(s) = a_0 + a_1 s + a_2 \cdot \frac{1}{2}(3s^2 - 1)$  and compare the coefficient of  $s^i$ .

$$(1) f(s) = \begin{cases} 0, & -1 < s < 0 \\ 1, & 0 < s < 1 \end{cases}$$

$$f(s) = \frac{1}{2} + \sum_{k=1}^{\infty} c_k P_k(s)$$

$$c_k = \frac{2k+1}{2} \int_{-1}^1 f(s) P'_k(s) ds = \frac{2k+1}{2} \int_0^1 P'_k(s) ds$$

Claim  $P'_k(0) = k(k+1) \int_0^1 P_k(s) ds$

pf  $\int_0^1 \frac{d}{ds} \left[ (1-s^2) \frac{dP_k(s)}{ds} \right] ds + k(k+1) \int_0^1 P_k(s) ds = 0$

$$\Rightarrow \left[ (1-s^2) \frac{dP_k(s)}{ds} \right]_0^1 + k(k+1) \int_0^1 P_k(s) ds = 0$$

eval at 0  $\Rightarrow P'_k(0) = k(k+1) \int_0^1 P_k(s) ds$

Therefore  $c_k = \frac{2k+1}{2k(k+1)} \cdot \frac{2k+1}{2} \int_0^1 P_k(s) ds = \frac{2k+1}{2k(k+1)} P'_k(0)$

So  $f(s) = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{2k+1}{2k(k+1)} P'_k(0) P_k(s)$ ,  $-1 < s < 1$

(2) set  $f(s) = \sum_{i=0}^2 a_i P_i(s) \Rightarrow a_0 + a_1 s + a_2 \frac{3s^2-1}{2} = 3s^2 + 4s + 1$

$$\Rightarrow \left(\frac{3a_2}{2}\right)s^2 + a_1 s + \left(a_0 - \frac{a_2}{2}\right) = 3s^2 + 4s + 1$$

$$\Rightarrow a_0 = 2, a_1 = 4, a_2 = 2$$

So  $f(s) = 2P_0(s) + 4P_1(s) + 2P_2(s)$

**Question 4.** Find the solution of Laplace's equation  $\nabla^2 u = 0$  in the sphere  $0 \leq r < a$  satisfying the boundary condition  $u(a, \theta) = 1$  if  $0 < \theta < \pi/2$  and  $u(a, \theta) = 0$  otherwise.

**Solution.**  $u(r, \theta) = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{(2k+1)P'_k(0)}{2k(k+1)} \left(\frac{r}{a}\right)^k P_k(\cos \theta)$ .

Sol  $\frac{1}{r^2} (r^2 \frac{\partial u}{\partial r})_r + \frac{1}{r^2 \sin \theta} (\sin \theta \frac{\partial u}{\partial \theta})_\theta = 0$

Suppose  $u$  separable,  $u(r, \theta) = R(r)\varphi(\theta)$

get  $\frac{(r^2 R')'}{R} = -\frac{(\sin \theta \varphi')'}{\sin \theta \varphi}$

$$\begin{cases} (\sin \theta \varphi')' + \lambda \sin \theta \varphi = 0 \\ (r^2 R')' - \lambda R = 0 \end{cases}$$

The radical eq:  $(r^2 R')' - n(n+1)R = 0 \Rightarrow r^2 R'' + 2r R' - n(n+1)R = 0$

sol:  $R_n = A_n r^n + B_n r^{-n-1}$

Since  $r^{-n-1} \rightarrow \infty$  as  $r \rightarrow 0$  for  $n \geq 0 \Rightarrow B_n$  must be 0.

$\Rightarrow$  the general sol is  $u(r, \theta) = \sum_{n=0}^{\infty} A_n r^n P_n(\cos \theta)$

Apply BV:  $f(\theta) = \sum_{n=0}^{\infty} A_n a^n P_n(\cos \theta) = \sum_{n=0}^{\infty} a_n P_n(\cos \theta) \Rightarrow A_n = \frac{a_n}{a^n}$

$a_n = \frac{2n+1}{2} \int_0^\pi f(\theta) P_n(\cos \theta) \sin \theta d\theta$

Since  $f(\theta)$  is 0 from 0 to  $\frac{\pi}{2} \Rightarrow a_n = \frac{2n+1}{2} \int_0^{\frac{\pi}{2}} P_n(\cos \theta) \sin \theta d\theta$

So  $A_n = \frac{(2n+1)P'_n(0)}{2n(n+1)a^n}, n \geq 2$

$= \frac{2n+1}{2} \int_0^1 P_n(x) dx = \frac{2n+1}{2} \frac{P'_n(0)}{n(n+1)}$   
by problem 3

$A_0 = \frac{a_0}{a^0} = \frac{1}{2}$

Therefore  $u(r, \theta) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{(2n+1)P'_n(0)}{2n(n+1)} \left(\frac{r}{a}\right)^n P_n(\cos \theta)$

**Question 5.** Find the solution of Laplace's equation  $\nabla^2 u = 0$  in the sphere  $0 \leq r < 1$  satisfying the boundary condition  $u = 3z^2 + 4z + 1$  if  $r = 1$ .

**Solution.**  $u(r, \theta) = 2 + 4r \cos \theta + r^2(3 \cos^2 \theta - 1)$  in spherical coordinates and  $u(x, y, z) = 2 + 4z + 2z^2 - x^2 - y^2$  in Euclidean coordinates.

Sol Same as problem 4 we get:

$$\begin{cases} (\sin \theta \varphi')' + \lambda \sin \theta \varphi = 0 \\ (r^2 R')' - \lambda R = 0 \end{cases}$$

The radial eq:  $(r^2 R')' - n(n+1)R = 0 \Rightarrow r^2 R'' + 2rR' - n(n+1)R = 0$

$$\text{sol: } R_n = A_n r^n + B_n r^{-(n+1)}$$

Since  $r^{-(n+1)} \rightarrow \infty$  as  $r \rightarrow 0$  for  $n \geq 0 \Rightarrow B_n$  must be 0.

$\Rightarrow$  the general sol is  $u(r, \theta) = \sum_{n=0}^{\infty} A_n r^n P_n(\cos \theta)$

$$\text{Apply BV: } u(1, \theta) = \sum_{n=0}^{\infty} A_n P_n(\cos \theta)$$

$$= f(\theta) = 3 \cos^2 \theta + 4 \cos \theta + 1 = 2P_2(\cos \theta) + 4P_1(\cos \theta) + 1P_0(\cos \theta)$$

by problem 3

$$= 2 + 4r \cos \theta + r^2(3 \cos^2 \theta - 1)$$

$$\text{So } A_0 = 2, A_1 = 4, A_2 = 2,$$

$$A_n = 0 \text{ for } n \geq 3$$

Therefore the general sol is  $u(r, \theta) = 2 + 4r \cos \theta + r^2(3 \cos^2 \theta - 1)$

in Euclidean coord it is:  $u(x, y, z) = 2 + 4z + 3z^2 - z^2 - r^2$

$$= 2 + 4z + 2z^2 - x^2 - y^2$$