

# TOWER OF FULLY COMMUTATIVE ELEMENTS OF TYPE $\tilde{A}$ AND APPLICATIONS

SADEK AL HARBAT

ABSTRACT. Let  $W^c(\tilde{A}_n)$  be the set of fully commutative elements in the affine Coxeter group  $W(\tilde{A}_n)$  of type  $\tilde{A}$ . We classify the elements of  $W^c(\tilde{A}_n)$  and give a normal form for them. We give a first application of this normal form to fully commutative affine braids. We then use this normal form to define two injections from  $W^c(\tilde{A}_{n-1})$  into  $W^c(\tilde{A}_n)$  and examine their properties. We finally consider the tower of affine Temperley-Lieb algebras of type  $\tilde{A}$  and use the injections above to prove the injectivity of this tower.

Braid groups; affine Coxeter groups; affine Temperley-Lieb algebra; fully commutative elements.

## 1. INTRODUCTION

Let  $(W, S)$  be a Coxeter system. We say that  $w$  in  $W$  is *fully commutative* if any reduced expression for  $w$  can be obtained from any other using only commutation relations among the members of the set  $S$ . If  $W$  is simply laced then the fully commutative elements of  $W$  are those with no  $sts$  factor in any reduced expression, where  $t$  and  $s$  are any non-commuting generators.

In this paper we are interested with the affine Coxeter group of type  $\tilde{A}$  which has an infinite set of fully commutative elements, as proved in [19] where Stembridge assigns to each fully commutative element  $w$  a unique labeled partial order, called the heap of  $w$ , whose linear extensions encode the reduced expressions for  $w$ . The notion of heap was used frequently as a way to approach affine fully commutative elements, while other notions, for example abacus diagrams, were used in [16]. In this work we only use algebraic methods to deal with them, such as the affine length (see Definition 2.6).

In a given Coxeter group the subset of fully commutative elements is indeed an interesting set with many remarkable properties, in particular relating to Kazhdan-Lusztig polynomials (see for example [3]) hence relating to  $\mu$ -coefficients. Moreover, they play the most important role in the  $M$ -coefficients notion, see [14]. Under certain conditions they are compatible with the classical Kazhdan-Lusztig cells, in the

sense that the set of fully commutative elements is a union of cells [13, 18]. There is also an intrinsic notion of cell coming from the structure of Temperley-Lieb algebra, those cells are classified in [8].

This paper is divided into two parts. The first part establishes a classification of affine fully commutative elements in type  $\tilde{A}$ : they are depicted by a normal form given in Theorem 2.11. This form is similar to Stembridge's description of fully commutative elements in the Coxeter groups of finite type  $A, B, D$  [20], although a classification of fully commutative elements of type  $A$  was given by Jones in [17] before even the official definition of fully commutative elements in the 90's, see for example [6, 10].

Classification is interesting in itself, nevertheless, since affine fully commutative elements in type  $\tilde{A}$  index a basis of the affine Temperley-Lieb algebra [7], it is to have consequences on the structure of the affine Temperley-Lieb algebra, on the tower of affine Temperley-Lieb algebras defined in [1] and on the traces on this algebra. This is precisely the point of the second part, which is divided into two applications.

The first application is to give a general form for “fully commutative braids” as follows: we lift the fully commutative elements to elements having the same expression in the  $\tilde{A}$ -type braid group  $B(\tilde{A}_n)$ , or: *fully commutative braids* (in this work we use the same symbols for the generators of the braid group and their images in the corresponding Coxeter group). Regarding  $B(\tilde{A}_{n-1})$  as a subgroup of  $B(\tilde{A}_n)$  by means of an injective homomorphism  $R_n$ , we give in Theorem 3.5 a general form for these fully commutative braids in terms of elements of  $B(\tilde{A}_{n-1})$  and the lift  $c_n$  of a certain Coxeter element to  $B(\tilde{A}_n)$ . The tower of affine braid groups:

$$\{1\} \longrightarrow B(\tilde{A}_1) \xrightarrow{R_1} \cdots \xrightarrow{R_{n-1}} B(\tilde{A}_{n-1}) \xrightarrow{R_n} B(\tilde{A}_n) \xrightarrow{R_{n+1}} \cdots$$

gives rise to an analogous injective tower of the group algebras  $K[B(\tilde{A}_n)]$  over an integral domain  $K$  of characteristic 0. Let  $q$  be an invertible element in  $K$ . The affine Temperley-Lieb algebra  $\widehat{TL}_{n+1}(q)$  is a quotient of the braid group algebra  $K[B(\tilde{A}_n)]$  and we get (see Section 5) a tower of affine Temperley-Lieb algebras:

$$\widehat{TL}_1(q) \xrightarrow{R_1} \widehat{TL}_2(q) \longrightarrow \cdots \xrightarrow{R_{n-1}} \widehat{TL}_n(q) \xrightarrow{R_n} \widehat{TL}_{n+1}(q) \xrightarrow{R_{n+1}} \cdots$$

The images by the quotient map of the fully commutative braids in  $K[B(\tilde{A}_n)]$  make up a basis of  $\widehat{TL}_{n+1}(q)$  and the form for fully commutative braids given in Theorem 3.5 is the key to the definition of *Markov elements* in  $\widehat{TL}_{n+1}(q)$ , and the key to proving that any trace on the affine Temperley-Lieb algebra is uniquely defined by its values on Markov elements [2, Theorem 4.6]. This in turn leads to the existence

and uniqueness of the affine Markov trace [2] and on the other hand is a step towards Green's conjectures (Property  $B$ ) [14].

The second application is to prove the faithfulness of the arrows of the tower of affine Temperley-Lieb algebras (Theorem 5.4). This was one of the most interesting questions since defining this tower in [1]. The faithfulness has consequences on the affine knot invariant defined in [1], and on the parabolic-like presentation defined in [1] on the level of affine Hecke algebra and recently for the affine Temperley-Lieb algebra.

The paper is organized as follows:

In Section 2, we give some general definitions, then we state and prove our main result, Theorem 2.11: a normal form for affine fully commutative elements in type  $\tilde{A}$ . This is the affine version of Theorem 2.3.

In Section 3, we define the tower of affine braid groups and establish its faithfulness. We then define fully commutative braids and, using our normal form and the fact that the lift  $c_n$  of a certain Coxeter element to  $B(\tilde{A}_n)$  acts as a Dynkin automorphism on  $B(\tilde{A}_{n-1})$  (Lemma 3.2), we find a general form for fully commutative braids in Theorem 3.5.

In Section 4, we show that the set  $W^c(\tilde{A}_{n-1})$  of fully commutative elements in the Coxeter group with  $n$  generators of type  $\tilde{A}$  injects into  $W^c(\tilde{A}_n)$  in two different ways (Theorem 4.2). The existence of these two injections  $I$  and  $J$  depends totally on the normal form of Theorem 2.11. The intersection of their images is the image of  $W^c(A_{n-1})$  on which they coincide.

In Section 5, we define the tower of affine Temperley-Lieb algebras coming from the tower of affine braid groups, then we prove in Theorem 5.4 the faithfulness of the arrows of this tower, using in a crucial way the injections  $I$  and  $J$  of the previous section.

## 2. A NORMAL FORM FOR AFFINE FULLY COMMUTATIVE ELEMENTS

Let  $(W, S)$  be a Coxeter system with associated Dynkin Diagram  $\Gamma$ . For  $s, t$  in  $S$  we let  $m_{st}$  be the order of  $st$  in  $W$ . Let  $w \in W$ . We denote by  $l(w)$  the length of a (any) reduced expression of  $w$ . We call *support of  $w$*  and denote by  $Supp(w)$  the subset of  $S$  consisting of all generators appearing in a (any) reduced expression of  $w$ . We define  $\mathcal{L}(w)$  to be the set of  $s \in S$  such that  $l(sw) < l(w)$ , in other words  $s$  appears at the left edge of some reduced expression of  $w$ . We define  $\mathcal{R}(w)$  similarly.

We know that from a given reduced expression of  $w$  we can arrive to any other reduced expression only by applying braid relations [4, §1.5 Proposition 5]. Among


these relations there are commutation relations: those that correspond to generators  $t$  and  $s$  with  $m_{st} = 2$ .

**Definition 2.1.** *Elements for which one can pass from any reduced expression to any other one only by applying commutation relations are called fully commutative elements. We denote by  $W^c$  the set of fully commutative elements in  $W$ .*

The center of our interest in this work is fully commutative elements in  $\tilde{A}$ -type Coxeter groups. In this case fully commutative elements have some additional elegant properties, in particular:

**Proposition 2.2.** [15, Lemma 3.1] *Let  $(W, S)$  be a Coxeter system such that  $m_{st}$  is odd or 2 for any  $s, t$  in  $S$ . Let  $w \in W$ . Then  $w$  is fully commutative if and only if every  $s$  in  $\text{Supp}(w)$  occurs the same number of times in any reduced expression of  $w$ .*

Consider the  $A$ -type Coxeter group with  $n$  generators  $W(A_n)$ , with the following Dynkin diagram:



We let:

$$\begin{aligned} [i, j] &= \sigma_i \sigma_{i-1} \dots \sigma_j \quad \text{for } n \geq i \geq j \geq 1 \quad \text{and} \quad [0, 1] = 1, \\ [i, j] &= \sigma_i \sigma_{i+1} \dots \sigma_j \quad \text{for } 1 \leq i \leq j \leq n \quad \text{and} \quad [n+1, n] = 1, \\ h(i, r) &= [i, 1][r, n] \quad \text{for } 0 \leq i < r \leq n+1 \quad \text{and} \quad (i, r) \neq (0, 1), \end{aligned}$$

hence

$$\begin{aligned} h(i, r) &= \sigma_i \sigma_{i-1} \dots \sigma_1 \sigma_r \sigma_{r+1} \dots \sigma_n \quad \text{for } 1 \leq i < r \leq n, \\ h(0, r) &= [r, n] \quad \text{for } 2 \leq r \leq n, \\ h(i, n+1) &= [i, 1] \quad \text{for } 1 \leq i \leq n, \\ h(0, n+1) &= 1. \end{aligned}$$

Considering right classes of  $W(A_{n-1})$  in  $W(A_n)$ , Stembridge has described canonical reduced words for elements of  $W(A_n)$ , namely:

$$[m_1, n_1][m_2, n_2] \dots [m_r, n_r]$$

where  $n \geq n_1 > \dots > n_r \geq 1$  and  $n_i \geq m_i \geq 1$  [20, p.1288]. He also proved [20, Corollary 5.8] that fully commutative elements are those for which the canonical reduced word satisfies  $m_1 > \dots > m_r$ . The set of fully commutative elements is stable under the inverse map; taking inverses we get:

**Theorem 2.3.** [20, Corollary 5.8]  $W^c(A_n)$  is the set of elements of the form:

$$(1) \quad [l_1, g_1][l_2, g_2] \cdots [l_s, g_s], \text{ with } \begin{cases} 1 \leq l_1 < \cdots < l_s \leq n, \\ 1 \leq g_1 < \cdots < g_s \leq n, \\ l_t \geq g_t \text{ for } 1 \leq t \leq s. \end{cases}$$

Inspecting the inequalities above, we see that the only term in expression (1) in which  $\sigma_n$  can occur is the  $s$ -th term. If  $\sigma_n$  does occur, then  $l_s$  must be equal to  $n$  and, whether or not  $g_s$  is equal to  $n$ ,  $\sigma_n$  occurs only once. Similarly, if  $\sigma_1$  does occur in expression (1), then  $g_1 = 1$  and  $\sigma_1 = \sigma_{g_1}$  will appear only once.

**Definition 2.4.** An element  $u$  in  $W^c(A_n)$  is called *extremal* if both  $\sigma_n$  and  $\sigma_1$  belong to  $\text{Supp}(u)$ .

**Lemma 2.5.** An extremal element different from  $[n, 1]$  can be written as

$$h(i, r) x \text{ with } 1 \leq i < r \leq n \text{ and } \text{Supp}(x) \subseteq \{\sigma_2, \dots, \sigma_{n-1}\}.$$


*Proof.* An extremal element  $u$  has a reduced expression of the form (1) above with  $g_1 = 1$  and  $l_s = n$ . If  $s = 1$  we have  $u = [n, 1] = h(n, n+1)$ , the only extremal element for which the leftmost term in the reduced expression (1) is  $\sigma_n$ .

Assume  $u \neq [n, 1]$ . For  $n = 1$  we have  $\sigma_1 = [1, 1]$ . For  $n = 2$  the element  $\sigma_1\sigma_2$  is the only extremal element different from  $[2, 1]$  and it is equal to  $h(1, 2)$ . Assume now  $n \geq 3$ .

The rightmost term in (1) is  $[n, g_s]$  with  $g_s > 1$ , so the generators on the right of  $\sigma_n$ , if any, belong to  $\{\sigma_2, \dots, \sigma_{n-1}\}$ . The generator  $\sigma_n$  commutes with any element in  $\{\sigma_2, \dots, \sigma_{n-2}\}$ , so, using the commutation relation  $\sigma_i\sigma_n = \sigma_n\sigma_i$  for  $2 \leq i \leq n-2$ , we can repeatedly push  $\sigma_n$  to the left in expression (1) above until the element on the left of  $\sigma_n$  is either  $\sigma_{n-1}$  or  $\sigma_1$ . In this process all generators  $\sigma_i$  that are pushed to the right of  $\sigma_n$  belong to  $\{\sigma_2, \dots, \sigma_{n-1}\}$ .

If we arrive at a subexpression  $a = \sigma_{n-1}\sigma_n$ , which happens if and only if  $l_{s-1} = n-1$ , then again it commutes with any element in  $\{\sigma_2, \dots, \sigma_{n-3}\}$  so we can push  $a$  to the left until the element on the left of  $a$  is either  $\sigma_{n-2}$  or  $\sigma_1$ . We continue in this way as long as  $l_{s-t} = n-t$  until we reach  $\sigma_1$ , and obtain the final expression  $[l_1, 1][n-k, n]x$ , with  $k = \max\{t \mid 0 \leq t < n-1 \text{ and } l_{s-t} = n-t\}$ , as announced.  $\square$

Now let  $W(\tilde{A}_n)$  be the affine Coxeter group of  $\tilde{A}$ -type with  $n+1$  generators, with the following Dynkin diagram:



Our notation encapsulates the fact that we view  $W(A_n)$  as the parabolic subgroup of  $W(\tilde{A}_n)$  generated by  $\sigma_1, \dots, \sigma_n$ . Recalling Proposition 2.2 we make the following definition.

**Definition 2.6.** We define the affine length of  $u$  in  $W^c(\tilde{A}_n)$  to be the number of times  $a_{n+1}$  occurs in a (any) reduced expression of  $u$ . We denote it by  $L(u)$ .

**Lemma 2.7.** Let  $w$  be a fully commutative element in  $W(\tilde{A}_n)$  with  $L(w) = m \geq 2$ . Fix a reduced expression of  $w$  as follows:

$$w = u_1 a_{n+1} u_2 a_{n+1} \dots u_m a_{n+1} u_{m+1}$$

with  $u_i$ , for  $1 \leq i \leq m+1$ , a reduced expression of a fully commutative element in  $W^c(A_n)$ . Then  $u_2, \dots, u_m$  are extremal elements and there is a reduced expression of  $w$  of the form:

$$(2) \quad w = h(i_1, r_1) a_{n+1} h(i_2, r_2) a_{n+1} \dots h(i_m, r_m) a_{n+1} v_{m+1}$$

where  $v_{m+1} \in W^c(A_n)$ ,  $0 \leq i_1 < r_1 \leq n+1$ ,  $(i_1, r_1) \neq (0, 1)$  and, for  $2 \leq t \leq m$ , we have either  $1 \leq i_t < r_t \leq n$  or  $(i_t, r_t) = (n, n+1)$ .

*Proof.* We first remark that in the same manner as in the previous lemma, we can write any fully commutative element in  $W(A_n)$  as  $h(i, r) x$  with  $0 \leq i < r \leq n+1$ ,  $(i, r) \neq (0, 1)$  and  $\text{Supp}(x) \subseteq \{\sigma_2, \dots, \sigma_{n-1}\}$ , in particular  $x$  commutes with  $a_{n+1}$ . Writing in this way  $u_1 = h(i_1, r_1) x_1$ , we can push  $x_1$  to the right of  $a_{n+1}$ , obtaining a new term  $u_2$  that we in turn write  $h(i_2, r_2) x_2$  with  $x_2$  commuting to  $a_{n+1}$ . Proceeding from left to right, we obtain formally form (2).

It remains to show that the elements  $u_i$ ,  $2 \leq i \leq m$ , are extremal. Indeed, if the support of some such  $u_i$  was contained in  $\{\sigma_2, \dots, \sigma_{n-1}\}$ , this  $u_i$  would commute with  $a_{n+1}$  and we would get a reduced expression containing  $a_{n+1} a_{n+1}$ , a contradiction. Now if some such  $u_i$  contained only one of  $\sigma_1, \sigma_n$ , then, using the commutation relations,  $a_{n+1} u_i a_{n+1}$  could be written  $\dots a_{n+1} \sigma_1 a_{n+1} \dots$  or  $\dots a_{n+1} \sigma_n a_{n+1} \dots$ , hence would contain a braid, which is impossible in a reduced expression for a fully commutative element.  $\square$

**Lemma 2.8.** *Let  $1 \leq l \leq n$  and  $0 \leq i < r \leq n+1$ ,  $(i, r) \neq (0, 1)$ . Then  $w = h(i, r) a_{n+1} \sigma_l$  is a reduced fully commutative element if and only if one of the following holds:*

- (1)  $l = r - 1 = i$ ;
- (2)  $i < l < r$ .

*Proof.* Assume first that  $r \leq n$  and write  $w = [i, 1][r, n]a_{n+1} \sigma_l$ . Using commutation relations we push  $\sigma_l$  to the left as long as it commutes with its left neighbour.

- If  $l \geq r$  we will arrive at the braid  $\sigma_l \sigma_{l+1} \sigma_l$  if  $l < n$ ,  $\sigma_n a_{n+1} \sigma_n$  if  $l = n$ :  $w$  is not fully commutative.
- Assume  $l < r$ . If  $i = 0$ , indeed  $w$  is reduced fully commutative. We proceed with  $i \geq 1$ .
  - If  $l < r - 1$  and  $l \leq i$ , again we get a braid  $\sigma_l \sigma_{l-1} \sigma_l$  by pushing  $\sigma_l$  leftmost hence  $w$  is not fully commutative.
  - If  $i < l$  indeed  $w$  is reduced fully commutative.
  - If  $r - 1 > 1$ , then  $\sigma_{r-1}$  commutes with  $a_{n+1}$  so for  $l = r - 1$  we get  $\sigma_i \dots \sigma_1 \sigma_r \sigma_{r-1} \sigma_{r+1} \dots \sigma_n a_{n+1}$  which is reduced fully commutative.
  - Finally if  $r - 1 = 1 = l$ , then  $\sigma_1$  cannot get past  $a_{n+1}$  on the left and again we have a reduced fully commutative element.

If  $r = n + 1$  and either  $l < i$  or  $l = i < n$ , the same process produces a braid  $\sigma_l \sigma_{l-1} \sigma_l$  or (if  $l = 1$ )  $\sigma_1 a_{n+1} \sigma_1$ , hence  $w$  is not fully commutative, while, for  $i < l$  or  $l = i = n$ ,  $w$  is reduced fully commutative.  $\square$

**Lemma 2.9.** *Let  $h(i, r)$  and  $h(i', r')$  be extremal elements different from  $[n, 1]$ . Then*

$$w = h(i, r) a_{n+1} h(i', r')$$

*is a reduced fully commutative element if and only if one of the following holds:*

- (1)  $i < i' < r' < r$ ;
- (2)  $i \leq i'$  and  $r' = r = i' + 1$ .

*Proof.* We have by assumption  $1 \leq i < r \leq n$  and  $1 \leq i' < r' \leq n$ . We write

$$w = [i, 1][r, n] a_{n+1} [i', 1][r', n].$$

Assume  $w$  is reduced fully commutative. From the previous lemma we must have  $i' = r - 1 = i$  or  $i < i' < r$ . We now examine  $r'$ , after noticing that if  $r' > i' + 1$ , then  $[i', 1]\sigma_{r'} = \sigma_{r'}[i', 1]$  so Lemma 2.8 imposes  $r' = r - 1 = i$  or  $i < r' < r$ .

If  $i' = r - 1 = i$ , then  $r' > i' + 1 = r$  is impossible by the previous remark, while  $r' = i' + 1 = r$  produces a reduced fully commutative  $w$ .

If  $i < i' < r$  then

- if  $r' > i' + 1$ , the previous remark gives  $r' = r - 1$  or  $i < r' < r$ , whence  $i < i' < r' < r$ , and under this condition  $w$  is reduced fully commutative;
- if  $r' = i' + 1 \leq r$ , we can write

$$w = [i, 1][r, n] a_{n+1} \sigma_{i'} \sigma_{i'+1} [i' - 1, 1][i' + 2, n].$$

We claim that no braid relation involving  $\sigma_{i'}$  or  $\sigma_{i'+1}$  can occur. On the right of the product  $\sigma_{i'} \sigma_{i'+1}$  in the expression just above, this is clear. This same product  $\sigma_{i'} \sigma_{i'+1}$  can be pushed to its left as long as it commutes with its left neighbour. If  $r' = i' + 1 < r - 1$  we arrive at

$$w = [i, 1] \sigma_{i'} \sigma_{i'+1} [r, n] a_{n+1} [i' - 1, 1][i' + 2, n]$$

where we see that our claim holds. If  $r' = i' + 1 = r - 1$  we arrive at

$$w = [i, 1] \sigma_{r-2} \sigma_r \sigma_{r-1} [r + 1, n] a_{n+1} [i' - 1, 1][i' + 2, n]$$

and our claim holds again. Finally if  $r' = i' + 1 = r$  we arrive at

$$\begin{aligned} w &= [i, 1] \sigma_r \sigma_{r-1} \sigma_{r+1} \sigma_r [r + 2, n] a_{n+1} [i' - 1, 1][i' + 2, n] \quad \text{if } r < n, \\ w &= [i, 1] \sigma_n a_{n+1} \sigma_{n-1} \sigma_n [n - 2, 1] \quad \text{if } r = n, \end{aligned}$$

and our claim is proved, hence  $w$  is reduced fully commutative.  $\square$

**Lemma 2.10.** *Let  $w \in W^c(\tilde{A}_n)$  with  $L(w) = m \geq 2$ . Write  $w$  as in (2) and assume that  $h(i_t, r_t) \neq [n, 1]$  for  $2 \leq t \leq m$ . There exist nonnegative integers  $p$  and  $k$  satisfying  $p + k = m$  and an integer  $j \in \{1, \dots, n - 1\}$  such that  $w$  has the following form:*

$$(3) \quad \begin{aligned} \text{if } p = 0 : w &= (h(j, j + 1) a_{n+1})^k w_r, \\ \text{if } p > 0 : w &= h(i_1, r_1) a_{n+1} \dots h(i_p, r_p) a_{n+1} (h(j, j + 1) a_{n+1})^k w_r, \end{aligned}$$

with  $w_r \in W^c(A_n)$  and, if  $p > 0$ :

- $1 \leq i_2 < \dots < i_p < r_p < \dots < r_2 \leq n$  and  $r_p - i_p \geq 2$ ,
- if  $k > 0$ : either  $i_p < j < j + 1 < r_p$  or  $j + 1 = r_p$ .

The element  $w_r$  can be described as follows:

- if  $k > 0$ : for some  $z \in \{0, \dots, n\}$  such that  $j + z - 1 \leq n$ , we have
  - if  $z = 0$  :  $w_r = 1$ ;
  - if  $z \geq 1$  :  $w_r = [j, d_1][j + 1, d_2] \dots [j + z - 1, d_z]$   
with  $1 \leq d_1 < \dots < d_z \leq n$  and  $j + c \geq d_{c+1}$  for  $0 \leq c \leq z - 1$ ;
- if  $k = 0$  (hence  $p > 0$ ): for some  $t \in \{0, \dots, n\}$  we have
  - if  $t = 0$  :  $w_r = 1$ ;



- if  $t \geq 1$ :  $w_r = [l_1, g_1][l_2, g_2] \dots [l_t, g_t]$  with:
  - \*  $1 \leq l_1 < \dots < l_t \leq n$ ,
  - \*  $1 \leq g_1 < \dots < g_t \leq n$ ,
  - \*  $l_i \geq g_i$  for  $1 \leq i \leq t$ ,
  - \*  $i_p < l_1 < r_p$ ,
  - \* for any  $i$ ,  $2 \leq i \leq t$ , such that  $l_i > l_{i-1} + 1$  we have  $l_i < r_p$ .

*Proof.* We apply Lemma 2.9 repeatedly from left to right, i.e. letting  $t$  increase in form (2) (Lemma 2.7). Case (1) forces the inequality  $i < i' < r' < r$ , it cannot happen more than  $\lfloor n/2 \rfloor$  times. Case (2) can only be followed by  $h(i', i' + 1)$  again. We thus get (3).

To determine  $w_r$  we use form (1) (Theorem 2.3) and we repeatedly apply Lemma 2.8, remembering that if  $l_i > l_{i-1} + 1$ , there is a reduced expression for  $w_r$  that begins with  $\sigma_{l_i}$ . If  $k > 0$  any reduced expression for  $w_r$  has to begin with  $\sigma_j$  on the left, we thus obtain a simple condition.  $\square$

We can now start the classification. Let  $w \in W(\tilde{A}_n)$  with  $L(w) = m \geq 1$ , written as in (2) in Lemma 2.7. We discuss according to the first term  $h(i_1, r_1)$  which can have one, and only one, of the following forms:

- (1)  $i_1 = n, r_1 = n + 1$  (extremal, equal to  $\lfloor n, 1 \rfloor$ );
- (2)  $1 \leq i_1 < r_1 \leq n$  (extremal, different from  $\lfloor n, 1 \rfloor$ );
- (3)  $i_1 = 0$  and  $2 \leq r_1 \leq n$  ( $\sigma_n$  appears but not  $\sigma_1$ );
- (4)  $1 \leq i_1 \leq n - 1$  and  $r_1 = n + 1$  ( $\sigma_1$  appears but not  $\sigma_n$ );
- (5)  $i_1 = 0$  and  $r_1 = n + 1$ .

We now examine each case.

- (1) By Lemma 2.8, after  $h(n, n + 1) = \sigma_n \sigma_{n-1} \dots \sigma_1$  there is only one choice for  $h(i_2, r_2)$ , namely  $h(n, n + 1)$  itself, and this repeats until we reach the rightmost term. This term must be 1 or some  $\lfloor n, i \rfloor$ . So  $w$  has the form:

$$(h(n, n + 1) a_{n+1})^k w_r \text{ with } w_r = 1 \text{ or } w_r = \lfloor n, i \rfloor.$$

- (2) By Lemma 2.8 all elements  $h(i_t, r_t)$  must also be extremal and different from  $\lfloor n, 1 \rfloor$ , hence the previous discussion applies and we get the forms in Lemma 2.10, either with  $p = 0$  (if  $r_1 = i_1 + 1$ ), or with  $p > 0$ . We only have to extend the condition of the lemma to  $(i_1, r_1)$ , that is:

$1 \leq i_1 < \dots < i_p < r_p < \dots < r_1 \leq n$ ,  $r_p - i_p \geq 2$  and, if  $k > 0$ , either  $i_p < j < j + 1 < r_p$  or  $j + 1 = r_p$ .

- (3) Assume first  $m \geq 2$ . By Lemma 2.8 again we must have  $i_2 < r_1$  hence, as before, all elements  $h(i_t, r_t)$  in form (2) must be extremal and different from  $\lfloor n, 1 \rfloor$ . Furthermore, if  $r_2 - i_2 > 1$ , then  $\sigma_{r_2}$  can be pushed to the left of

$h(i_2, r_2)$  so Lemma 2.8 implies  $r_2 < r_1$ . We obtain the possible forms from Lemma 2.10 with  $p > 0$ , with the condition:

$$0 = i_1 < i_2 < \cdots < i_p < r_p < \cdots < r_2 < r_1 \leq n, \quad r_p - i_p \geq 2$$

and, if  $k > 0$ , either  $i_p < j < j + 1 < r_p$  or  $j + 1 = r_p$ .

If  $m = 1$ , we have to describe the rightmost term  $v_2$  in expression (2). Writing  $v_2$  in form (1) as in the proof of Lemma 2.10, we find exactly the same expression as in the case  $k = 0$  for  $w_r$  in this lemma, with  $p = 1$ .

- (4) If  $m \geq 2$ , Lemma 2.8 gives  $i_2 > i_1$ . If  $i_2 = n$ , from the same lemma we obtain

$$w = h(i_1, n + 1) a_{n+1} (h(n, n + 1) a_{n+1})^k w_r$$

with  $k > 0$  and  $w_r = 1$  or  $w_r = \lfloor n, i \rfloor$  for some  $i$ ,  $1 \leq i \leq n$ .

If  $i_2 < n$  we are back to the case of extremal elements different from  $\lfloor n, 1 \rfloor$  and we obtain the possible forms from Lemma 2.10 with  $p > 0$ , with the condition:

$$1 \leq i_1 < i_2 < \cdots < i_p < r_p < \cdots < r_2 < r_1 = n + 1, \quad r_p - i_p \geq 2 \text{ and, if } k > 0, \text{ either } i_p < j < j + 1 < r_p \text{ or } j + 1 = r_p.$$

If  $m = 1$ , Lemma 2.10 provides, for the rightmost term, the same expression as in the case  $k = 0$  for  $w_r$  in this lemma, with  $p = 1$ .

- (5) Here if  $m \geq 2$ , in case  $i_2 < n$ , then in the notation of Lemma 2.10 we have  $h(i_1, r_1) = 1$ , and in case  $i_2 = n$ , we have a form similar to case (1), namely

$$a_{n+1} (h(n, n + 1) a_{n+1})^k w_r \text{ with } w_r = 1 \text{ or } w_r = \lfloor n, i \rfloor.$$

If  $m = 1$  the rightmost term can be any fully commutative element  $w_r$  in  $W(A_n)$ . Actually the description of  $w_r$  given in Lemma 2.10, with  $k = 0$ ,  $p = 1$ ,  $i_1 = 0$  and  $r_1 = n + 1$ , applies here as well.

We subsume this discussion in the following theorem:

**Theorem 2.11.** *Let  $w \in W^c(\tilde{A}_n)$  with  $L(w) \geq 1$ . Then  $w$  can be written in a unique way as a reduced word of the following form, for nonnegative integers  $p$  and  $k$  and for  $j \in \{1, \dots, n\}$ :*

$$(4) \quad \begin{aligned} &\text{if } p = 0 : w = (h(j, j + 1) a_{n+1})^k w_r \text{ (with } k > 0), \\ &\text{if } p > 0 : w = h(i_1, r_1) a_{n+1} \cdots h(i_p, r_p) a_{n+1} (h(j, j + 1) a_{n+1})^k w_r, \end{aligned}$$

with  $w_r \in W^c(A_n)$  and, if  $p > 0$ :

- $0 \leq i_1 < \cdots < i_p < r_p < \cdots < r_1 \leq n + 1$  and  $r_p - i_p \geq 2$ ,
- if  $k > 0$ : either  $i_p < j < j + 1 < r_p$  or  $j + 1 = r_p$ .


In particular we have  $p \leq \lfloor \frac{n+1}{2} \rfloor$ .

The element  $w_r$  can be described as follows:

- If  $k > 0$ : for some  $z \in \{0, \dots, n\}$  such that  $j + z - 1 \leq n$ , we have
  - if  $z = 0$  :  $w_r = 1$ ;
  - if  $z \geq 1$  :  $w_r = [j, d_1][j+1, d_2] \dots [j+z-1, d_z]$   
with  $1 \leq d_1 < \dots < d_z \leq n$  and  $j+c \geq d_{c+1}$  for  $0 \leq c \leq z-1$ .
- If  $k = 0$  (hence  $p > 0$ ): for some  $t \in \{0, \dots, n\}$  we have
  - if  $t = 0$  :  $w_r = 1$ ;
  - if  $t \geq 1$  :  $w_r = [l_1, g_1][l_2, g_2] \dots [l_t, g_t]$  with:
    - \*  $1 \leq l_1 < \dots < l_t \leq n$ ,
    - \*  $1 \leq g_1 < \dots < g_t \leq n$ ,
    - \*  $l_i \geq g_i$  for  $1 \leq i \leq t$ ,
    - \*  $i_p < l_1 < r_p$ ;
    - \* for any  $i$ ,  $2 \leq i \leq t$ , such that  $l_i > l_{i-1} + 1$  we have  $l_i < r_p$ .

Conversely, any word written as above is reduced and fully commutative.

**Example 2.12.** Let  $w$  be in  $W^c(\tilde{A}_2)$ . Then there exists  $k \geq 0$  such that  $w$  has one and only one of the following forms:



We finish this section with simple remarks. Firstly, the affine length of a fully commutative element  $w$  written in form (4) is  $p+k$ . Secondly, the family of elements corresponding to the case  $k > 0$  and  $j = n$  is particularly simple: their form is

$$h(i, n+1) a_{n+1} (h(n, n+1) a_{n+1})^t w_r$$

for some  $i$  such that  $0 \leq i \leq n$  and with  $w_r = 1$  or  $[n, l]$  with  $n \geq l \geq 1$ .

### 3. FULLY COMMUTATIVE AFFINE BRAIDS

We denote by  $B(\tilde{A}_n)$  the affine braid group with  $n+1$  generators of type  $\tilde{A}$ , while we denote by  $B(A_n)$  the braid group with  $n$  generators of type  $A$ , where  $n \geq 1$ . By

definition  $B(\tilde{A}_n)$  has  $\{\sigma_1, \sigma_2, \dots, \sigma_n, a_{n+1}\}$  as a set of generators together with the following defining relations:

- (1)  $\sigma_i \sigma_j = \sigma_j \sigma_i$  for  $1 \leq i, j \leq n$  and  $|i - j| \geq 2$ ,
- (2)  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$  for  $1 \leq i \leq n - 1$ ,
- (3)  $\sigma_i a_{n+1} = a_{n+1} \sigma_i$  for  $2 \leq i \leq n - 1$ ,
- (4)  $\sigma_1 a_{n+1} \sigma_1 = a_{n+1} \sigma_1 a_{n+1}$  for  $n \geq 2$ ,
- (5)  $\sigma_n a_{n+1} \sigma_n = a_{n+1} \sigma_n a_{n+1}$  for  $n \geq 2$ ,

while  $B(A_n)$  is generated by  $\{\sigma_1, \sigma_2, \dots, \sigma_n\}$  with relations (1) and (2).

**Lemma 3.1.** *The following map:*

$$\begin{aligned} R_n : B(\tilde{A}_{n-1}) &\longrightarrow B(\tilde{A}_n) \\ \sigma_i &\longmapsto \sigma_i \quad \text{for } 1 \leq i \leq n - 1 \\ a_n &\longmapsto \sigma_n a_{n+1} \sigma_n^{-1} \end{aligned}$$

*is a group monomorphism.*

*Proof.* Graham and Lehrer give in [12, §2] a presentation of the  $B$ -type braid group  $B(B_{n+1})$  with generators  $\{\tau_{n+1}, \sigma_1, \dots, \sigma_n, a_{n+1}\}$  and relations (1) to (5) above, plus

$$\tau_{n+1} \sigma_i \tau_{n+1}^{-1} = \sigma_{i+1} \text{ for } 1 \leq i \leq n - 1, \quad \tau_{n+1} \sigma_n \tau_{n+1}^{-1} = a_{n+1}, \quad \tau_{n+1} a_{n+1} \tau_{n+1}^{-1} = \sigma_1.$$

This presentation is related to the usual presentation of the  $B$ -type braid group, with generators  $\{t, \sigma_1, \sigma_2, \dots, \sigma_n\}$  and braid relations, by

$$\tau_{n+1} = t \sigma_1 \sigma_2 \dots \sigma_n, \quad a_{n+1} = \tau_{n+1} \sigma_n \tau_{n+1}^{-1} = t \sigma_1 \sigma_2 \dots \sigma_n \sigma_{n-1}^{-1} \dots \sigma_1^{-1} t^{-1}.$$

They show that the subgroup of  $B(B_{n+1})$  generated by  $\{\sigma_1, \dots, \sigma_n, a_{n+1}\}$  is isomorphic to  $B(\tilde{A}_n)$  and fits into the exact sequence [12, Corollary 2.7]

$$1 \longrightarrow B(\tilde{A}_n) \longrightarrow B(B_{n+1}) \longrightarrow \mathbb{Z} \longrightarrow 1.$$

We thus view  $B(\tilde{A}_n)$  as a subgroup of the braid group  $B(B_{n+1})$  for  $n \geq 1$ .

There is a natural injection from  $B(B_n)$  generated by  $\{t, \sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$  into  $B(B_{n+1})$  generated by  $\{t, \sigma_1, \sigma_2, \dots, \sigma_n\}$  (see e.g. [9, §2.2]). We claim that this injection restricts to  $R_n$  on  $B(\tilde{A}_{n-1})$ . To prove this claim we only need to check that

the image of  $a_n$  is  $\sigma_n a_{n+1} \sigma_n^{-1}$ . Since  $\sigma_n$  commutes with  $t, \sigma_1, \dots, \sigma_{n-2}$ , we have:

$$\begin{aligned} \sigma_n a_{n+1} \sigma_n^{-1} &= \sigma_n t \sigma_1 \sigma_2 \dots \sigma_n \sigma_{n-1}^{-1} \dots \sigma_1^{-1} t^{-1} \sigma_n^{-1} \\ &= t \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_n \sigma_{n-1} \sigma_n \sigma_{n-1}^{-1} \sigma_n^{-1} \sigma_{n-2}^{-1} \dots \sigma_1^{-1} t^{-1} \\ &= t \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1} \sigma_n \sigma_{n-1} \sigma_{n-1}^{-1} \sigma_n^{-1} \sigma_{n-2}^{-1} \dots \sigma_1^{-1} t^{-1} \\ &= t \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1} \sigma_{n-2}^{-1} \dots \sigma_1^{-1} t^{-1} \\ &= a_n \end{aligned}$$

as announced.  $\square$

Using the injective morphism  $R_n$  we now view  $B(\tilde{A}_{n-1})$  as a subgroup of  $B(\tilde{A}_n)$ . Let  $\psi_n$  be the Dynkin automorphism of  $B(\tilde{A}_{n-1})$  that shifts the generators of the Dynkin diagram one step counterclockwise ( $\sigma_1 \mapsto a_n \mapsto \sigma_{n-1} \mapsto \dots \mapsto \sigma_2 \mapsto \sigma_1$ ). It generates a subgroup of  $\text{Aut}(B(\tilde{A}_{n-1}))$  of order  $n$ . We simply refer to it by  $\psi$  in what follows.

**Lemma 3.2.** *The element  $c_n = \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1}$  of  $B(\tilde{A}_n)$  normalizes  $B(\tilde{A}_{n-1})$ . It acts by conjugacy on  $B(\tilde{A}_{n-1})$  in the same way as  $\psi$ . We will write:*

$$c_n^t h = \psi^t[h] c_n^t \text{ for any } h \in B(\tilde{A}_{n-1}).$$

*Proof.* Applying braid relations in  $B(\tilde{A}_n)$  we see that, for  $2 \leq i \leq n-1$ , we have:

$$\begin{aligned} \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1} \sigma_i &= \sigma_n \dots \sigma_{i+1} \sigma_i \sigma_{i-1} \sigma_i \dots \sigma_1 a_{n+1} \\ &= \sigma_n \dots \sigma_{i+1} \sigma_{i-1} \sigma_i \sigma_{i-1} \dots \sigma_1 a_{n+1} \\ &= \sigma_{i-1} \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1}. \end{aligned}$$

In a similar way, we use the braid relations involving  $a_{n+1}$  and the following relations worth keeping in mind:

$$(5) \quad a_{n+1} a_n = \sigma_n^{-1} a_n \sigma_n a_n = \sigma_n^{-1} \sigma_n a_n \sigma_n = a_n \sigma_n = \sigma_n a_{n+1}$$

to obtain:

$$\begin{aligned} \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1} \sigma_1 &= a_n \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1}, \\ \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1} a_n &= \sigma_{n-1} \sigma_n \sigma_{n-1} \dots \sigma_1 a_{n+1} \end{aligned}$$

hence our result.  $\square$

Let  $n \geq 1$ . Recall from [4, IV.1, Proposition 5] that there exists a map

$$g : W(\tilde{A}_n) \longrightarrow B(\tilde{A}_n)$$

such that, given any reduced expression of  $w \in W(\tilde{A}_n)$ , the image  $g(w)$  of  $w$  is defined by the same expression in the braid group (keeping the same symbols for the generators of the affine braid group and their images via the natural surjection onto the affine Coxeter group).

**Definition 3.3.** We call fully commutative braids the images under  $g$  of the fully commutative elements in  $W(\tilde{A}_n)$ , i.e. the elements of  $\{g(w) \mid w \in W^c(\tilde{A}_n)\}$ .

We now proceed to give a general form for fully commutative braids in which the element  $c_n$  is singled out, as a consequence of the normal form for fully commutative elements in Theorem 2.11. We use the notation in Section 2, equally valid for the braid groups, with an additional index  $n$  in  $B(\tilde{A}_n)$  and  $n-1$  in  $B(\tilde{A}_{n-1})$ .

**Lemma 3.4.** Consider the element  $y = h_n(j, j+1) a_{n+1}$  of  $B(\tilde{A}_n)$  where  $1 \leq j \leq n$ . Let  $k \geq 1$  be an integer and write  $k = m(n-j+1) + r$  with  $0 \leq r < n-j+1$ . Then if  $j = n$  we have  $y^k = c_n^k$  while for  $j < n$  we have (with the convention  $[n, n+1-r] = 1$  if  $r = 0$ ):

$$(1) \text{ if } m = 0: \quad y^k = (h_{n-1}(j, j+1) a_n)^r [n, n+1-r];$$

$$(2) \text{ if } m > 0:$$

$$y^k = \left( \prod_{i=0}^{i=m-1} \psi^i \left[ (h_{n-1}(j, j+1) a_n)^{n-j} [j, n-1] \right] \right) \psi^m \left[ (h_{n-1}(j, j+1) a_n)^r \right] c_n^m [n, n+1-r].$$

*Proof.* We have for  $j < n$ , using relations (5):

$$y = \sigma_j \dots \sigma_1 \sigma_{j+1} \dots \sigma_{n-1} \sigma_n a_{n+1} = \sigma_j \dots \sigma_1 \sigma_{j+1} \dots \sigma_{n-1} a_n \sigma_n = h_{n-1}(j, j+1) a_n \sigma_n.$$

Observe that for  $j+1 < s \leq n$  we have  $\sigma_s y = y \sigma_{s-1}$ . Hence if  $j+1 < n$ :

$$y^2 = h_{n-1}(j, j+1) a_n \sigma_n y = h_{n-1}(j, j+1) a_n y \sigma_{n-1} = (h_{n-1}(j, j+1) a_n)^2 \sigma_n \sigma_{n-1}.$$

Continuing this way, we can see that whenever  $0 \leq t \leq n-j$  we have

$$y^t = (h_{n-1}(j, j+1) a_n)^t [n, n+1-t].$$

This holds in particular for  $t = n-j$  thus:

$$y^{n-j+1} = (h_{n-1}(j, j+1) a_n)^{n-j} \sigma_n \sigma_{n-1} \dots \sigma_{j+1} \sigma_j \dots \sigma_2 \sigma_1 \sigma_{j+1} \dots \sigma_{n-1} \sigma_n a_{n+1}.$$

We now use:  $\sigma_n \sigma_{n-1} \dots \sigma_1 \sigma_u = \sigma_{u-1} \sigma_n \sigma_{n-1} \dots \sigma_1$  for  $2 \leq u \leq n$ , and get:

$$y^{n-j+1} = (h_{n-1}(j, j+1) a_n)^{n-j} [j, n-1] c_n.$$

An easy induction using Lemma 3.2 leads to:

$$y^{m(n-j+1)} = \left( \prod_{i=0}^{i=m-1} \psi^i \left[ (h_{n-1}(j, j+1) a_n)^{n-j} [j, n-1] \right] \right) c_n^m.$$

Finally, let  $k = m(n - j + 1) + r$ , where  $0 \leq r < n - j + 1$ . We have:

$$\begin{aligned} y^k &= \left( \prod_{i=0}^{i=m-1} \psi^i \left[ (h_{n-1}(j, j+1)a_n)^{n-j} [j, n-1] \right] \right) c_n^m (h_{n-1}(j, j+1)a_n)^r [n, n+1-r] \\ &= \left( \prod_{i=0}^{i=m-1} \psi^i \left[ (h_{n-1}(j, j+1)a_n)^{n-j} [j, n-1] \right] \right) \psi^m \left[ (h_{n-1}(j, j+1)a_n)^r \right] c_n^m [n, n+1-r] \end{aligned}$$

as claimed.  $\square$

**Theorem 3.5.** *Let  $n \geq 2$ . Let  $w$  be a fully commutative braid in  $B(\tilde{A}_n)$ . Then  $w$  can be written in one and only one of the following two forms:*

$$\begin{aligned} & u c_n^t v \\ \text{or} \quad & [i, 1] a_{n+1} u c_n^t v, \end{aligned}$$

with  $u \in B(\tilde{A}_{n-1})$ ,  $v \in B(A_n)$ ,  $t \geq 0$  and  $0 \leq i \leq n-1$ .

*Proof.* We have  $w = g(\bar{w})$  with  $\bar{w}$  in  $W^c(\tilde{A}_n)$ , and we use the reduced expression of  $w$  corresponding to the normal form of  $\bar{w}$  given by Theorem 2.11, of which we use the notations. We may and do ignore the rightmost term  $w_r$  that belongs to  $B(A_n)$  and we remark that if  $p = 0$  the element has indeed the first claimed form, by Lemma 3.4. We proceed with  $p > 0$  and set

$$x = h_n(i_1, r_1) a_{n+1} \dots h_n(i_p, r_p) a_{n+1}$$

so that  $w = xy^k$  where  $y = h_n(j, j+1) a_{n+1}$  as in the previous lemma.

We remark that for  $1 \leq i \leq p$  we have  $r_i \leq r_1 - i + 1$ ; since  $r_1 \leq n+1$  this gives:  $r_i \leq n - i + 2$ . We have two main cases to consider:

- $r_1 \leq n$ , that is  $\sigma_n$  belongs to the support of  $\sigma_{r_1} \dots \sigma_{n-1} \sigma_n$  ;
- $r_1 = n+1$ , that is  $\sigma_n$  does not belong to the support of  $\sigma_{r_1} \dots \sigma_{n-1} \sigma_n$ .

We start with the first case  $r_1 \leq n$ , so that  $r_i \leq n - i + 1$ . Using first the relation  $\sigma_n a_{n+1} = a_n \sigma_n$ , then the braid relations  $\sigma_u \sigma_{u-1} \sigma_u = \sigma_{u-1} \sigma_u \sigma_{u-1}$  for  $u = n, n-1, \dots, n-p+2$ , we push to the right the first  $\sigma_n$  from the left and get:

$$x = [i_1, 1] [r_1, n-1] a_n h(i_2, r_2) a_{n+1} \dots h(i_p, r_p) a_{n+1} \sigma_{n-(p-1)}.$$

We repeat this operation with the first  $\sigma_n$  from the left in this new expression, and so on. Proceeding from  $i = 1$  to  $i = p$ , we obtain:

$$x = h_{n-1}(i_1, r_1) a_n \dots h_{n-1}(i_p, r_p) a_n \sigma_n \sigma_{n-1} \dots \sigma_{n-(p-1)}.$$

We set  $\rho = h_{n-1}(i_1, r_1) a_n \dots h_{n-1}(i_p, r_p) a_n$ , an element of  $B(\tilde{A}_{n-1})$ . If  $k = 0$  we have  $w = x = \rho[n, n - (p-1)]$  and our claim holds. Let  $k \geq 1$  and set  $\epsilon = n - (p-1)$ . We have  $\epsilon \geq j+1$  since  $j+1 \leq r_p \leq n - (p-1)$  and we can write:

$$w = \rho \sigma_n \sigma_{n-1} \dots \sigma_\epsilon y^k, \text{ with } \rho \in B(\tilde{A}_{n-1}) \text{ and } \epsilon \geq j+1.$$

We recall that  $y = h_n(j, j+1) a_{n+1}$  acts on  $\sigma_i$  in the following way:  $\sigma_i y = y \sigma_{i-1}$  for  $j+1 < i \leq n$ . We distinguish two cases:

$$(1) \ 1 \leq k \leq \epsilon - (j+1),$$

$$(2) \ \epsilon - (j+1) < k.$$

We start with (1). We have:  $\sigma_n \sigma_{n-1} \dots \sigma_\epsilon y^k = y^k \sigma_{n-k} \sigma_{n-1-k} \dots \sigma_{\epsilon-k}$ , with

$$k \leq \epsilon - (j+1) = n - j - p < n - j.$$

Thus we are in case (1) of Lemma 3.4, that is:

$$w = \rho(h_{n-1}(j, j+1) a_n)^k [n, n+1-k] [n-k, \epsilon-k] \in B(\tilde{A}_{n-1}) B(A_n)$$

as required (this is the first form in the theorem with  $t = 0$ ).

Now we deal with case (2) and set  $h = k - (\epsilon - (j+1)) \geq 1$ . Using the computation from case (1) for  $\epsilon - (j+1)$  we get, using again Lemma 3.4 (1):

$$\begin{aligned} \sigma_n \sigma_{n-1} \dots \sigma_\epsilon y^k &= \sigma_n \sigma_{n-1} \dots \sigma_\epsilon y^{\epsilon-(j+1)} y^h = y^{\epsilon-(j+1)} \sigma_{j+p} \sigma_{j+p-1} \dots \sigma_{j+1} y^h \\ &= (h_{n-1}(j, j+1) a_n)^{\epsilon-(j+1)} [n, j+p+1] [j+p, j+1] y^h \\ &= (h_{n-1}(j, j+1) a_n)^{\epsilon-(j+1)} [n, j+1] y y^{h-1}. \end{aligned}$$

We now compute:

$$\begin{aligned} [n, j+1] y y^{h-1} &= (\sigma_n \dots \sigma_{j+1})(\sigma_j \dots \sigma_1)(\sigma_{j+1} \dots \sigma_n) a_{n+1} y^{h-1} \\ &= (\sigma_n \dots \sigma_1)(\sigma_{j+1} \dots \sigma_n) a_{n+1} y^{h-1} \\ &= (\sigma_j \dots \sigma_{n-1})(\sigma_n \dots \sigma_1) a_{n+1} y^{h-1} \end{aligned}$$

since  $(\sigma_n \dots \sigma_1) \sigma_k = \sigma_{k-1} (\sigma_n \dots \sigma_1)$  for  $2 \leq k \leq n$ . Setting

$$\eta = \rho(h_{n-1}(j, j+1) a_n)^{\epsilon-(j+1)} \sigma_j \dots \sigma_{n-1} \in B(\tilde{A}_{n-1}),$$

we get  $w = \eta c_n y^{h-1}$ .

(a) If  $h-1 \leq n-j$ , we see, using Lemma 3.4, that:

$$\begin{aligned} w &= \eta c_n (h_{n-1}(j, j+1) a_n)^{h-1} [n, n+1-(h-1)] \\ &= \eta \psi \left[ (h_{n-1}(j, j+1) a_n)^{h-1} \right] c_n [n, n+1-(h-1)], \end{aligned}$$



which is of the first claimed form.

- (b) If  $n - j < h - 1$ , we write  $h - 1 = m(n - j + 1) + r$  with  $0 \leq r < n - j + 1$  as in Lemma 3.4 and get:

$$\begin{aligned} w &= \eta c_n \left( \prod_{i=0}^{i=m-1} \psi^i \left[ (h_{n-1}(j, j+1)a_n)^{n-j} [j, n-1] \right] \right) \psi^m \left[ (h_{n-1}(j, j+1)a_n)^r \right] \\ &\quad c_n^m [n, n+1-r] \\ &= \eta \left( \prod_{i=1}^{i=m} \psi^i \left[ (h_{n-1}(j, j+1)a_n)^{n-j} [j, n-1] \right] \right) \psi^{m+1} \left[ (h_{n-1}(j, j+1)a_n)^r \right] \\ &\quad c_n^{m+1} [n, n+1-r] \end{aligned}$$

which has indeed the required form (first form in the theorem).

We are left with the second main case  $r_1 = n + 1$ : the element under study has a normal form  $w = [i_1, 1]a_{n+1}w'$  with  $0 \leq i_1 \leq n$ . In fact  $i_1 = n$  is the case of positive powers of  $c_n = \sigma_n \dots \sigma_1 a_{n+1}$  that have the first form in the theorem. For  $0 \leq i_1 < n$  the element  $w'$  actually belongs to one of the previous cases thus we get the second form in the theorem.  $\square$

Using the natural surjection of  $B(\tilde{A}_n)$  onto  $W(\tilde{A}_n)$ , we deduce from Theorem 3.5 two possible forms for fully commutative elements in  $W(\tilde{A}_n)$ . In the above notation, the image of  $v \in B(A_n)$  belongs to  $W(A_n)$  and, considering the left classes of  $W(A_{n-1})$  in  $W(A_n)$  as in [20, p.1288], this image can be written uniquely as a product  $v'\sigma_n\sigma_{n-1}\dots\sigma_s$  with  $v' \in W(A_{n-1})$  and  $1 \leq s \leq n+1$  (with the convention that for  $s = n+1$ , the product  $\sigma_n\sigma_{n-1}\dots\sigma_s$  is 1). Since  $c_n$  normalizes  $W(\tilde{A}_{n-1})$  the element  $v'$  can be moved to the left of  $c_n^t$  into an element that incorporates with the image of  $u$ . We obtain:

**Corollary 3.6.** *Let  $w$  be a fully commutative element in  $W(\tilde{A}_n)$  for  $n \geq 2$ . Then  $w$  can be written in one and only one of the following two forms:*

$$\begin{aligned} &d c_n^t \sigma_n \sigma_{n-1} \dots \sigma_s \\ \text{or } &[i, 1] a_{n+1} d c_n^t \sigma_n \sigma_{n-1} \dots \sigma_s, \end{aligned}$$

where  $d$  is in  $W(\tilde{A}_{n-1})$  and  $t, s$  and  $i$  are integers with  $t \geq 0$ ,  $1 \leq s \leq n+1$  and  $0 \leq i \leq n-1$ .

## 4. THE TOWER OF FULLY COMMUTATIVE ELEMENTS

The injection of braid groups  $R_n : B(\tilde{A}_{n-1}) \longrightarrow B(\tilde{A}_n)$  of Lemma 3.1 induces the group homomorphism:

$$\begin{aligned} R_n : W(\tilde{A}_{n-1}) &\longrightarrow W(\tilde{A}_n) \\ \sigma_i &\longmapsto \sigma_i \text{ for } 1 \leq i \leq n-1 \\ a_n &\longmapsto \sigma_n a_{n+1} \sigma_n. \end{aligned}$$

**Lemma 4.1.** *The morphism  $R_n : W(\tilde{A}_{n-1}) \longrightarrow W(\tilde{A}_n)$  is an injection.*

*Proof.* This fact is most easily seen using the description of  $W(\tilde{A}_n)$  as the group of  $(n+1)$ -periodic permutations of  $\mathbb{Z}$  with total shift equal to 0, which we briefly recall. A permutation  $u$  of  $\mathbb{Z}$  is  $m$ -periodic if

$$u(i+m) = u(i) + m \text{ for any } i \in \mathbb{Z}.$$

We define the total shift of an  $m$ -periodic permutation  $u$  to be:

$$\frac{1}{m} \sum_{i=1}^{i=m} (u(i) - i).$$

We set  ${}_0^m\mathbb{Z}$  to be the set of  $m$ -periodic permutations with total shift equal to 0. It forms a subgroup of the group of permutations of  $\mathbb{Z}$ . Let  $i$  be an integer such that  $1 \leq i \leq m$  and let  ${}_m s_i$  be the  $m$ -periodic permutation defined by:  ${}_m s_i(k) = k+1$  for  $k \equiv i \pmod{m}$ ,  ${}_m s_i(k) = k-1$  for  $k \equiv i+1 \pmod{m}$ ,  ${}_m s_i(k) = k$  for  $k \not\equiv i, i+1 \pmod{m}$ . Then

$$\begin{aligned} \sigma_i &\longmapsto {}^{n+1}s_i \quad \text{for } 1 \leq i \leq n \\ a_{n+1} &\longmapsto {}^{n+1}s_{n+1} \end{aligned} \tag{6}$$

is an isomorphism from  $W(\tilde{A}_n)$  onto  ${}_0^{n+1}\mathbb{Z}$  (see e.g. [5, Proposition 3.2]).

It is easy to check that  ${}_0^n\mathbb{Z}$  injects into  ${}_0^{n+1}\mathbb{Z}$  as follows. We define an injection  $\phi : \mathbb{Z} \longrightarrow \mathbb{Z}$  by letting  $\phi(i+kn) = i+k(n+1)$  for  $1 \leq i \leq n$  and  $k \in \mathbb{Z}$ . We then map an  $n$ -periodic permutation  $v$  to the  $(n+1)$ -periodic permutation  $v'$  defined by  $v'(\phi(x)) = \phi(v(x))$  ( $x \in \mathbb{Z}$ ) and  $v'(k(n+1)) = k(n+1)$  ( $k \in \mathbb{Z}$ ). For  $v \in {}_0^n\mathbb{Z}$  and  $1 \leq i \leq n$ , write  $v(i) = j_i + k_i n$  with  $1 \leq j_i \leq n$ ; the total shift of  $v$  is  $(\sum_{i=1}^{i=n} k_i) + \frac{1}{n} \sum_{i=1}^{i=n} (j_i - i)$ . Since  $v$  is an  $n$ -periodic permutation it induces a permutation

mod  $n$ , hence  $\sum_{i=1}^{i=n} j_i = \sum_{i=1}^{i=n} i$  and consequently  $\sum_{i=1}^{i=n} k_i = 0$ . Now the total shift of  $v'$  is

$$\begin{aligned} \frac{1}{n+1} \sum_{i=1}^{i=n+1} (v'(i) - i) &= \frac{1}{n+1} \left( \sum_{i=1}^{i=n} (\phi(v(i)) - i) \right) + \frac{1}{n+1} (n+1 - (n+1)) \\ &= \frac{1}{n+1} \sum_{i=1}^{i=n} (j_i + k_i(n+1) - i) = 0 \end{aligned}$$

This injection maps  ${}^n s_i$  to  ${}^{n+1} s_i$  for  $1 \leq i \leq n-1$ , and  ${}^n s_n$  to  ${}^{n+1} s_n {}^{n+1} s_{n+1} {}^{n+1} s_n$ . Going back to  $W(\tilde{A}_{n-1})$  and  $W(\tilde{A}_n)$  through the isomorphisms (6) above, we find the morphism  $R_n$ , hence itself an injection.  $\square$

The Coxeter group  $W(A_{n-1})$  with Coxeter generators  $(\sigma_1, \dots, \sigma_{n-1})$  is a parabolic subgroup of  $W(A_n)$ . This is no longer the case for  $W(\tilde{A}_{n-1})$  and  $W(\tilde{A}_n)$ , indeed proper parabolic subgroups of  $W(\tilde{A}_n)$  are finite. This is an important difficulty when dealing with the affine case.  $W(A_n)$  though, with Coxeter generators  $(\sigma_1, \dots, \sigma_n)$ , is a parabolic subgroup of  $W(\tilde{A}_n)$  and fully commutative elements of  $W(A_n)$  are fully commutative in  $W(\tilde{A}_n)$  as well. As for  $W(\tilde{A}_{n-1})$ , the injection  $R_n : W(\tilde{A}_{n-1}) \rightarrow W(\tilde{A}_n)$  of Lemma 4.1 is a group homomorphism, but it does not preserve full commutativity, as can be seen directly on the image of  $a_n$ , namely  $\sigma_n a_{n+1} \sigma_n$ . When dealing with fully commutative elements, the notion of homomorphism of groups may thus become irrelevant. We introduce the following maps:

**Theorem 4.2.** *For  $w \in W^c(\tilde{A}_{n-1})$ , let  $I(w)$  (resp.  $J(w)$ ) be the element of  $W(\tilde{A}_n)$  obtained by substituting  $\sigma_n a_{n+1}$  (resp.  $a_{n+1} \sigma_n$ ) to  $a_n$  in the normal form for  $w$ . Both expressions are reduced and both  $I(w)$  and  $J(w)$  are fully commutative, with the same affine length as  $w$ . The maps:*

$$I, J : W^c(\tilde{A}_{n-1}) \longrightarrow W^c(\tilde{A}_n)$$

*are injective and satisfy  $l(I(w)) = l(J(w)) = l(w) + L(w)$ . Furthermore the images of  $I$  and  $J$  intersect exactly on  $W^c(A_{n-1})$ .*

*Proof.* We use the notation in Section 2 with an additional index  $n$  in  $W(\tilde{A}_n)$  and  $n-1$  in  $W(\tilde{A}_{n-1})$ . We then see that  $I(h_{n-1}(i, j) a_n) = h_n(i, j) a_{n+1}$ .

Let  $w \in W^c(\tilde{A}_{n-1})$ . By inspection of the normal forms in  $W^c(\tilde{A}_{n-1})$  and  $W^c(\tilde{A}_n)$  given in Theorem 2.11 one sees directly that  $I(w)$  is the normal form of a fully

commutative element in  $W(\tilde{A}_n)$  and that this process is injective. Indeed we have:

$$\begin{aligned} I(h_{n-1}(i_1, r_1) a_n \dots h_{n-1}(i_p, r_p) a_n (h_{n-1}(j, j+1) a_n)^k w_r) \\ = h_n(i_1, r_1) a_{n+1} \dots h_n(i_p, r_p) a_{n+1} (h_n(j, j+1) a_{n+1})^k w_r. \end{aligned}$$

As for  $J(w)$ , the corresponding normal form is obtained as follows. We first observe that

$$J(h_{n-1}(j, j+1) a_n) = [j, 1][j+1, n-1]a_{n+1}\sigma_n = [j, 1]a_{n+1}[j+1, n],$$

which implies, since  $\sigma_j, \dots, \sigma_1$  commute with  $\sigma_{j+2}, \dots, \sigma_n$ :

$$\begin{aligned} J((h_{n-1}(j, j+1) a_n)^2) &= [j, 1]a_{n+1}[j+1, n][j, 1]a_{n+1}[j+1, n] \\ &= [j, 1]a_{n+1}[j+1, 1][j+2, n]a_{n+1}[j+1, n], \end{aligned}$$

and inductively:

$$J((h_{n-1}(j, j+1) a_n)^k) = [j, 1]a_{n+1} (h_n(j+1, j+2)a_{n+1})^{k-1} [j+1, n].$$

For  $k > 0$  we thus get

$$(7) \quad J((h_{n-1}(j, j+1) a_n)^k w_r) = h_n(j, n+1)a_{n+1} (h_n(j+1, j+2)a_{n+1})^{k-1} [j+1, n]w_r$$

and we observe that either  $w_r = 1$ , in which case  $[j+1, n]$  has the shape required by Theorem 2.11, or  $w_r = [j, d_1][j+1, d_2] \dots [j+z-1, d_z]$ , in which case

$$[j+1, n]w_r = [j+1, d_1][j+2, d_2] \dots [j+z, d_z][j+z+1, n]$$

which is again the shape of the rightmost element required in Theorem 2.11. The expression (7) is thus the normal form of  $J(w)$  when  $p = 0$  in Theorem 2.11 (4).

If  $p$  is positive in Theorem 2.11 (4), we have:

$$\begin{aligned} J(h_{n-1}(i_p, r_p) a_n)[j, 1]a_{n+1} &= [i_p, 1][r_p, n-1]a_{n+1}\sigma_n[j, 1]a_{n+1} \\ &= [i_p, 1]a_{n+1}[r_p, n][j, 1]a_{n+1}. \end{aligned}$$

Thus if  $j < r_p - 1$ :

$$\begin{aligned} J(h_{n-1}(i_p, r_p) a_n)[j, 1]a_{n+1} &= [i_p, 1]a_{n+1}[j, 1][r_p, n]a_{n+1} \\ &= [i_p, 1]a_{n+1}h_n(j, r_p)a_{n+1} \end{aligned}$$

while if  $j = r_p - 1$ :

$$\begin{aligned} J(h_{n-1}(i_p, r_p) a_n)[j, 1]a_{n+1} &= [i_p, 1]a_{n+1}[r_p, 1][r_p+1, n]a_{n+1} \\ &= [i_p, 1]a_{n+1}h_n(j+1, j+2)a_{n+1}. \end{aligned}$$

We proceed in the same way from right to left, noticing that when going from  $(i_t, r_t)$  to  $(i_{t-1}, r_{t-1})$  the case  $i_t = r_{t-1} - 1$  cannot occur, so we get:

$$J(h_{n-1}(i_{t-1}, r_{t-1}) a_n) [i_t, 1] a_{n+1} = [i_{t-1}, 1] a_{n+1} h_n(i_t, r_{t-1}) a_{n+1}.$$

We can now write the normal form of  $J(w)$  when  $p > 0$  in Theorem 2.11 (4):

$J(h_{n-1}(i_1, r_1) a_n \dots h_{n-1}(i_p, r_p) a_n (h_{n-1}(j, j+1) a_n)^k w_r)$  is equal to:

$$\begin{aligned} & h_n(i_1, n+1) a_{n+1} \dots h_n(i_p, r_{p-1}) a_{n+1} h_n(j, r_p) a_{n+1} (h_n(j+1, j+2) a_{n+1})^{k-1} [j+1, n] w_r \\ & \quad \text{if } k > 0 \text{ and } j < r_p - 1, \\ & h_n(i_1, n+1) a_{n+1} \dots h_n(i_p, r_{p-1}) a_{n+1} (h_n(j+1, j+2) a_{n+1})^k [j+1, n] w_r \\ & \quad \text{if } k > 0 \text{ and } j = r_p - 1, \\ & h_n(i_1, n+1) a_{n+1} \dots h_n(i_p, r_{p-1}) a_{n+1} [r_p, n] w_r \quad \text{if } k = 0. \end{aligned}$$

We see as before, by a suitable right shift of  $\sigma_{r_p}, \dots, \sigma_n$ , that for  $k = 0$  the rightmost term  $[r_p, n] w_r$  has again the shape required by Theorem 2.11.

The fact that the substitution process adds to the original length the number of occurrences of  $a_n$ , i.e. the affine length, is clear. As for the intersection of the images, one only needs to notice that if a reduced expression of a fully commutative element contains an element  $\sigma_n$  to the left of the first  $a_{n+1}$  from left to right, then all reduced expressions for this element have the same property.  $\square$

We remark that the injection  $I$  is well defined only on the set of fully commutative elements. Indeed, substituting  $\sigma_n a_{n+1}$  to  $a_n$  in the two reduced expressions  $\sigma_{n-1} a_n \sigma_{n-1}$  and  $a_n \sigma_{n-1} a_n$  gives rise to different elements of  $W(\tilde{A}_n)$ . It might be the case that  $I$  is well defined on the set of elements for which the number of occurrences of  $a_n$  in any reduced expression is the same, but as we do not need this, we will not examine it further.

## 5. THE TOWER OF TEMPERLEY-LIEB ALGEBRAS

Let  $K$  be an integral domain of characteristic 0 and  $q$  be an invertible element in  $K$ . We mean by algebra in what follows  $K$ -algebra. For  $x, y$  in a given ring with identity we define

$$V(x, y) = xyx + xy + yx + x + y + 1.$$

For  $n \geq 2$ , we define  $\widehat{TL}_{n+1}(q)$  to be the algebra with unit given by the set of generators  $\{g_{\sigma_1}, \dots, g_{\sigma_n}, g_{a_{n+1}}\}$ , with the following relations (see [11, 0.1, 0.5]):

$$(8) \quad \left\{ \begin{array}{l} g_{\sigma_i} g_{\sigma_j} = g_{\sigma_j} g_{\sigma_i} \text{ for } 1 \leq i, j \leq n \text{ and } |i - j| \geq 2, \\ g_{\sigma_i} g_{a_{n+1}} = g_{a_{n+1}} g_{\sigma_i} \text{ for } 2 \leq i \leq n - 1, \\ g_{\sigma_i} g_{\sigma_{i+1}} g_{\sigma_i} = g_{\sigma_{i+1}} g_{\sigma_i} g_{\sigma_{i+1}} \text{ for } 1 \leq i \leq n - 1, \\ g_{\sigma_i} g_{a_{n+1}} g_{\sigma_i} = g_{a_{n+1}} g_{\sigma_i} g_{a_{n+1}} \text{ for } i = 1, n, \\ g_{\sigma_i}^2 = (q - 1)g_{\sigma_i} + q \text{ for } 1 \leq i \leq n, \\ g_{a_{n+1}}^2 = (q - 1)g_{a_{n+1}} + q, \\ V(g_{\sigma_i}, g_{\sigma_{i+1}}) = V(g_{\sigma_1}, g_{a_{n+1}}) = V(g_{\sigma_n}, g_{a_{n+1}}) = 0 \text{ for } 1 \leq i \leq n - 1. \end{array} \right.$$

We set  $\widehat{TL}_1(q) = K$ . For  $n = 1$ , the algebra  $\widehat{TL}_2(q)$  is generated by two elements:  $g_{\sigma_1}, g_{a_2}$ , with only Hecke quadratic relations. That is:

$$g_{\sigma_1}^2 = (q - 1)g_{\sigma_1} + q \quad \text{and} \quad g_{a_2}^2 = (q - 1)g_{a_2} + q.$$

Let  $w$  be a fully commutative element in  $W(\tilde{A}_n)$ . Pick a reduced expression of  $w$ , say  $w = s_1 \cdots s_k$  with  $s_i \in \{\sigma_1, \dots, \sigma_n, a_{n+1}\}$  for  $1 \leq i \leq k$ . Then  $g_w := g_{s_1} \cdots g_{s_k}$  is a well defined element in  $\widehat{TL}_{n+1}(q)$  that does not depend on the choice of a reduced expression of  $w$ , and the set  $\{g_w \mid w \in W^c(\tilde{A}_n)\}$  is a  $K$ -basis of  $\widehat{TL}_{n+1}(q)$  [7, Proposition 1]. The multiplication associated to this basis satisfies, for  $w, v$  in  $W^c(\tilde{A}_n)$  and  $s$  in  $\{\sigma_1, \dots, \sigma_n, a_{n+1}\}$ :

$$\begin{aligned} g_w g_v &= g_{wv} && \text{whenever } l(wv) = l(w) + l(v) \text{ and } wv \in W^c(\tilde{A}_n), \\ g_s g_w &= (q - 1)g_w + qg_{sw} && \text{whenever } l(sw) = l(w) - 1. \end{aligned}$$

The classical Temperley-Lieb algebra of type  $A$  with  $n$  generators,  $TL_n(q)$ , can be regarded as the subalgebra of  $\widehat{TL}_{n+1}(q)$  generated by  $\{g_{\sigma_1}, \dots, g_{\sigma_n}\}$ . A  $K$ -basis of  $TL_n(q)$  is given by  $\{g_w \mid w \in W^c(A_n)\}$ . We set  $TL_0(q) = K$ .

The affine Temperley-Lieb algebra  $\widehat{TL}_{n+1}(q)$  is the quotient of the affine braid group algebra  $K[B(\tilde{A}_n)]$  by the ideal generated by Relations (8). The injection of braid groups  $R_n : B(\tilde{A}_{n-1}) \longrightarrow B(\tilde{A}_n)$  of Lemma 3.1 induces the injection of group algebras:

$$\begin{aligned} G_n : K[B(\tilde{A}_{n-1})] &\longrightarrow K[B(\tilde{A}_n)] \\ \sigma_i &\longmapsto \sigma_i && \text{for } 1 \leq i \leq n - 1 \\ a_n &\longmapsto \sigma_n a_{n+1} \sigma_n^{-1} \end{aligned}$$

which, as we will see shortly, induces an algebra homomorphism at the Temperley-Lieb level. Yet the possible lack of injectivity of this homomorphism forces us to

use different notations for the generators of  $\widehat{TL}_n(q)$  and  $\widehat{TL}_{n+1}(q)$ . In the following proposition we use  $\{t_{\sigma_1}, \dots, t_{\sigma_{n-1}}, t_{a_n}\}$  as the set of generators for  $\widehat{TL}_n(q)$  satisfying relations (8) (with  $t$  replacing  $g$  and  $n$  replacing  $n+1$ ).

**Proposition 5.1.** *The injection  $G_n$  induces the following morphism of algebras:*

$$\begin{aligned} R_n : \widehat{TL}_n(q) &\longrightarrow \widehat{TL}_{n+1}(q) \\ t_{\sigma_i} &\longmapsto g_{\sigma_i} \text{ for } 1 \leq i \leq n-1 \\ t_{a_n} &\longmapsto g_{\sigma_n} g_{a_{n+1}} g_{\sigma_n}^{-1}. \end{aligned}$$

The restriction of  $R_n$  to  $TL_{n-1}(q)$  is an injective morphism into  $TL_n(q)$  and satisfies  $R_n(t_w) = g_{I(w)} = g_{J(w)}$  for  $w \in W^c(A_{n-1})$ .

*Proof.* We first have to show that the defining relations (8) for  $\widehat{TL}_n(q)$ , with  $t$  replacing  $g$  and  $n$  replacing  $n+1$ , are preserved for the images in  $TL_n(q)$ . This is immediate for those relations that do not involve  $a_n$ , using directly relations (8) for  $\widehat{TL}_{n+1}(q)$ . For the others, this is easily checked. For instance we have:

$$\begin{aligned} R_n(V(t_{\sigma_1}, t_{a_n})) &= g_{\sigma_n} V(g_{\sigma_1}, g_{a_{n+1}}) g_{\sigma_n}^{-1} = 0, \\ R_n(V(t_{\sigma_{n-1}}, t_{a_n})) &= g_{\sigma_n} g_{\sigma_{n-1}} V(g_{\sigma_n}, g_{a_{n+1}}) g_{\sigma_{n-1}}^{-1} g_{\sigma_n}^{-1} = 0. \end{aligned}$$

The last assertion comes from the fact that  $W(A_{n-1})$  is parabolic in  $W(A_n)$ : fully commutative elements of type  $A_{n-1}$  embed in those of type  $A_n$  and this embedding preserves the normal form.  $\square$

**Remark 5.2.** *The injection  $G_n : K[B(\tilde{A}_{n-1})] \longrightarrow K[B(\tilde{A}_n)]$  also induces a homomorphism of the corresponding Hecke algebras. We have shown in [1, Proposition 4.3.3] that this homomorphism is injective for  $K = \mathbb{Z}[q, q^{-1}]$  where  $q$  is an indeterminate. We will not need this fact in what follows.*

**Proposition 5.3.** *For any  $w \in W^c(\tilde{A}_{n-1})$  of positive affine length, the element  $R_n(t_w)$  has the following form:*

$$R_n(t_w) = (-1)^{L(w)} g_{I(w)} + \left(-\frac{1}{q}\right)^{L(w)} g_{J(w)} + \sum_{\substack{l(x) < l(I(w)) \\ L(x) \leq L(w)}} \alpha_x g_x \quad (\alpha_x \in K).$$

*Proof.* We prove the statement by induction on the affine length  $L(w)$  of an element  $w \in W^c(\tilde{A}_{n-1})$ . We first assume that  $L(w) = 1$ . We can write  $w = ua_nv$  where  $u$

and  $v$  belong to  $W^c(A_{n-1})$ . We compute:

$$\begin{aligned}
R_n(t_{a_n}) &= g_{\sigma_n} g_{a_{n+1}} g_{\sigma_n}^{-1} = g_{\sigma_n} g_{a_{n+1}} \left( \frac{1}{q} g_{\sigma_n} + \left( \frac{1}{q} - 1 \right) g_1 \right) \\
&= \frac{1}{q} g_{\sigma_n} g_{a_{n+1}} g_{\sigma_n} + \left( \frac{1}{q} - 1 \right) g_{\sigma_n} g_{a_{n+1}} \\
&= -\frac{1}{q} g_{\sigma_n} g_{a_{n+1}} - \frac{1}{q} g_{a_{n+1}} g_{\sigma_n} - \frac{1}{q} g_{\sigma_n} - \frac{1}{q} g_{a_{n+1}} - \frac{1}{q} g_1 + \left( \frac{1}{q} - 1 \right) g_{\sigma_n} g_{a_{n+1}} \\
&= -g_{\sigma_n} g_{a_{n+1}} - \frac{1}{q} g_{a_{n+1}} g_{\sigma_n} - \frac{1}{q} g_{\sigma_n} - \frac{1}{q} g_{a_{n+1}} - \frac{1}{q} g_1.
\end{aligned}$$

Since  $R_n$  is a homomorphism of algebras we have  $R_n(t_w) = R_n(t_u) R_n(t_{a_n}) R_n(t_v)$  and, by Proposition 5.1, we have  $R_n(t_v) = g_{I(v)} = g_{J(v)}$ . Furthermore, multiplying on the left the element  $v \in W^c(A_{n-1})$  by any element of  $\{\sigma_n a_{n+1}, a_{n+1} \sigma_n, \sigma_n, a_{n+1}\}$  produces a reduced fully commutative word, hence:

$$R_n(t_{a_n}) R_n(t_v) = -g_{\sigma_n a_{n+1} I(v)} - \frac{1}{q} g_{a_{n+1} \sigma_n I(v)} - \frac{1}{q} g_{\sigma_n I(v)} - \frac{1}{q} g_{a_{n+1} I(v)} - \frac{1}{q} g_{I(v)}.$$

Finally, since  $u$  also belongs to  $W^c(A_{n-1})$  we get, as claimed:

$$\begin{aligned}
R_n(t_w) &= g_{I(u)} \left( -g_{\sigma_n a_{n+1} I(v)} - \frac{1}{q} g_{a_{n+1} \sigma_n I(v)} - \frac{1}{q} g_{\sigma_n I(v)} - \frac{1}{q} g_{a_{n+1} I(v)} - \frac{1}{q} g_{I(v)} \right) \\
&= -g_{I(w)} - \frac{1}{q} g_{J(w)} - \frac{1}{q} \sum_{\substack{l(x) < l(I(w)) \\ L(x) \leq 1}} \alpha_x g_x \quad (\alpha_x \in K).
\end{aligned}$$

We now assume that the property holds for any  $u$  of positive affine length at most  $k$ . Any  $w \in W^c(\tilde{A}_{n-1})$  with  $L(w) = k + 1$  can be written as  $w = u a_n v$  where  $u \in W^c(\tilde{A}_{n-1})$ ,  $L(u) = k$ ,  $v \in W^c(A_{n-1})$ , and  $l(w) = l(u) + l(v) + 1$ . We have  $R_n(t_w) = R_n(t_u) R_n(t_{a_n}) R_n(t_v)$ . Using our previous computation of  $R_n(t_{a_n}) R_n(t_v)$  and the induction hypothesis we write:

$$\begin{aligned}
R_n(t_w) &= \left( (-1)^{L(u)} g_{I(u)} + \left( -\frac{1}{q} \right)^{L(u)} g_{J(u)} + \sum_{\substack{l(x) < l(I(u)) \\ L(x) \leq L(u)}} \alpha_x g_x \right) \\
&\quad \left( -g_{\sigma_n a_{n+1} I(v)} - \frac{1}{q} g_{a_{n+1} \sigma_n I(v)} - \frac{1}{q} g_{\sigma_n I(v)} - \frac{1}{q} g_{a_{n+1} I(v)} - \frac{1}{q} g_{I(v)} \right).
\end{aligned}$$

We know from Theorem 4.2 that  $I(w) = I(u) \sigma_n a_{n+1} I(v)$  and  $J(w) = J(u) a_{n+1} \sigma_n I(v)$  and that both are reduced fully commutative words, hence  $g_{I(u)} g_{\sigma_n a_{n+1} I(v)} = g_{I(w)}$  and



$g_{J(u)}g_{a_{n+1}\sigma_n I(v)} = g_{J(w)}$ . We obtain the two leading terms

$$(-1)^{L(w)}g_{I(w)} + \left(-\frac{1}{q}\right)^{L(w)}g_{J(w)}$$

in the formula that we are looking for.

We now observe that the other terms in the development of the product above have affine length at most  $L(w)$  and Coxeter length at most  $l(I(w))$ . The only terms that might have length  $l(I(w))$  come from  $I(u)$  and  $J(u)$  in the first parenthesis, together with  $\sigma_n a_{n+1} I(v)$  and  $a_{n+1} \sigma_n I(v)$  in the second. The cases of  $I(w)$  and  $J(w)$  being settled, it remains to prove that  $g_{I(u)}g_{a_{n+1}\sigma_n I(v)}$  and  $g_{J(u)}g_{\sigma_n a_{n+1} I(v)}$  are linear combinations of basis elements  $g_x$  where the length of  $x \in W^c(\tilde{A}_n)$  is strictly less than  $l(I(w))$ .

Remember from Lemma 2.7 that between two consecutive appearances of  $a_{n+1}$  we must see one and only one occurrence of  $\sigma_n$ . So the word  $I(u)a_{n+1}\sigma_n I(v)$  either is not reduced, hence of length strictly less than  $l(I(w))$ , or is not fully commutative. In the latter case it contains a braid so the corresponding product  $g_{I(u)}g_{a_{n+1}\sigma_n I(v)}$  decomposes, in the Temperley Lie algebra, into a linear combination of elements  $g_z$  with  $l(z) < l(I(w))$ . Similarly  $J(u)\sigma_n a_{n+1} I(v)$  has two occurrences of  $\sigma_n$  between the rightmost occurrence of  $a_{n+1}$  and the previous one on the left: it cannot be fully commutative reduced hence the product  $g_{J(u)}g_{\sigma_n a_{n+1} I(v)}$  decomposes as before into terms of strictly smaller length. The result follows.  $\square$

**Theorem 5.4.** *The tower of affine Temperley-Lieb algebras*

$$\widehat{TL}_1(q) \xrightarrow{R_1} \widehat{TL}_2(q) \xrightarrow{R_2} \widehat{TL}_3(q) \longrightarrow \cdots \longrightarrow \widehat{TL}_n(q) \xrightarrow{R_n} \widehat{TL}_{n+1}(q) \longrightarrow \cdots$$

*is a tower of faithful arrows.*

*Proof.* We need to show that  $R_n$  is an injective homomorphism of algebras. A basis for  $\widehat{TL}_n(q)$  is given by the elements  $t_w$  where  $w$  runs over  $W^c(\tilde{A}_{n-1})$ . Assume that there are non trivial dependence relations between the images of these basis elements. Pick one such relation, say  $\sum_w \lambda_w R_n(t_w) = 0$ , and let  $k = \max\{l(w) + L(w) \mid \lambda_w \neq 0\}$ . Using Proposition 5.3 we can write this relation as follows:

$$\sum_{\substack{l(w)+L(w)=k \\ L(w)>0}} \lambda_w \left( (-1)^{L(w)} g_{I(w)} + \left(-\frac{1}{q}\right)^{L(w)} g_{J(w)} \right) + \sum_{\substack{l(w)+L(w)=k \\ L(w)=0}} \lambda_w g_{I(w)} + \sum_{l(x)<k} \lambda'_x g_x = 0$$

for suitable coefficients  $\lambda'_x$  (where the  $x$ 's are elements of  $W^c(\tilde{A}_n)$  and  $l(x)$  is the length in  $W(\tilde{A}_n)$ ). Since the elements  $g_y$  for  $y \in W^c(\tilde{A}_n)$  form a basis of  $\widehat{TL}_{n+1}(q)$ , and since  $I$  and  $J$  are injective and the intersection of their images is  $W^c(A_{n-1})$ , we see that all the coefficients  $\lambda_w$  for  $l(w) + L(w) = k$  must be 0, a contradiction.  $\square$

## REFERENCES

- [1] S. Al Harbat, On the affine braid group, affine Temperley-Lieb algebra and Markov trace, PH.D Thesis, Université Paris-Diderot-Paris 7, 2013.
- [2] S. Al Harbat, Markov trace on a tower of affine Temperley-Lieb algebras of type  $\tilde{A}$ , J. Knot Theory Ramifications 24 (9) (2015), 1–28.
- [3] S. C. Billey and G. S. Warrington, Kazhdan-Lusztig polynomials for 321-hexagon-avoiding permutations, J. Algebraic Combin. 13 (2) (2001), 111–136.
- [4] N. Bourbaki, Groupes et algèbres de Lie, Chapitres 4, 5 et 6, Masson, Paris, 1981.
- [5] F. Digne, A Garside presentation for Artin-Tits groups of type  $\tilde{C}_n$ , Ann. Inst. Fourier (Grenoble) 62 (2) (2012), 641–666.
- [6] C. K. Fan, A Hecke Algebra Quotient and Properties of Commutative Elements of a Weyl group, Ph.D. Thesis, MIT, 1995.
- [7] C. K. Fan, A Hecke algebra quotient and some combinatorial applications, J. Algebraic Combin. 5 (3) (1996), 175–189.
- [8] C. K. Fan and R. M. Green, On the affine Temperley-Lieb algebras, J. Lond. Math. Soc. 60 (2) (1999), 366–380.
- [9] M. Geck and S. Lambropoulou, Markov traces and knot invariants related to Iwahori-Hecke algebras of type  $B$ , J. Reine Angew. Math. 482 (1997), 191–213.
- [10] J. J. Graham, Modular representations of Hecke algebras and related algebras, Ph.D. Thesis, University of Sydney, 1995.
- [11] J. J. Graham and G. I. Lehrer, The representation theory of affine Temperley-Lieb algebras, Enseign. Math. 44 (1998), 173–218.
- [12] J. J. Graham and G. I. Lehrer, Diagram algebras, Hecke algebras and decomposition numbers at roots of unity, Ann. Sci. Éc. Norm. Supér. 36 (4) (2003), 479–524.
- [13] R. M. Green and J. Losonczy, Fully commutative Kazhdan-Lusztig cells, Ann. Inst. Fourier (Grenoble) 51 (4) (2001), 1025–1045.
- [14] R. M. Green, Generalized Jones traces and Kazhdan-Lusztig bases, J. Pure Appl. Algebra 211 (3) (2007), 744–772.
- [15] M. Hagiwara, M. Ishikawa and H. Tagawa, A Characterization of the Simply-Laced FC-Finite Coxeter Groups, Ann. Comb. 8 (2) (2004), 177–196.
- [16] C. R. H. Hanusa and B. C. Jones, The enumeration of fully commutative affine permutations, European J. Combin. 31 (5) (2010), 1342–1359.
- [17] V. F. R. Jones, A polynomial invariant for knots via Von Neumann algebras, Bull. Amer. Math. Soc. 12 (1) (1985), 103–111.
- [18] J.-Y. Shi, Fully commutative elements and Kazhdan-Lusztig cells in the finite and affine Coxeter groups, Proc. Amer. Math. Soc. 131 (11) (2003), 3371–3378.
- [19] J. R. Stembridge, On the fully commutative elements of Coxeter groups, J. Algebraic Combin. 5 (4) (1996), 353–385.

- [20] J. R. Stembridge, Some combinatorial aspects of reduced words in finite Coxeter groups, Trans. Amer. Math. Soc. 349 (4) (1997), 1285–1332.

*E-mail address:* `sadikharbat@math.univ-paris-diderot.fr`