

Q1. Which is not an example of a quantitative variable?

- ☐ (A) Age
- ☐ (B) Marital status
- ☐ (C) Salary
- ☐ (D) Weight

Q2. The ordinal scale is used when:

- ☐ (A) Differences between values are meaningful
- ☐ (B) There is a true zero
- ☐ (C) Categories can be ranked
- ☐ (D) Data can only be labeled

Q3. Which of the following is a discrete variable?

- ☐ (A) Temperature
- ☐ (B) Height
- ☐ (C) Number of siblings
- ☐ (D) Distance

Q4. The sample is:

- ☐ (A) Always larger than the population
- ☐ (B) The entire group being studied
- ☒ (C) A subset of the population
- ☐ (D) Chosen only randomly

Q5. Which graph is best suited for quantitative, continuous data?

- ☐ (A) Bar graph
- ☒ (B) Histogram
- ☐ (C) Pie chart
- ☐ (D) Pareto chart

Q6. Query times (ms): 5, 6, 7, 7, 8, 9, 50. Which measure changes MOST if 50 becomes 500?

- ☐ (A) Median
- ☐ (B) Mode
- ☐ (C) Mean
- ☐ (D) First Quartile

Q7. If 95th percentile response time is 220 ms, it means:

- ☐ (A) 95% of responses are exactly 220 ms
- ☐ (B) 5% of responses exceed 220 ms
- ☐ (C) 95% of responses exceed 220 ms
- ☐ (D) Median is 220 ms

Q8. If CPU utilization data shows: Mean = 75%, Median = 82% and Mode = 90%. The system usage distribution is:

- ☐ (A) Positively skewed
- ☒ (B) Negatively skewed
- ☐ (C) Symmetric
- ☐ (D) Uniform

Q9. Two sorting algorithms are tested (in ms):

Algorithm A: 19, 20, 21, 22, 23, 30

Algorithm B: 15, 16, 21, 21, 24, 30

Which statement is correct?

- ☐ (A) Both algorithms have equal performance
- ☐ (B) Algorithm B is more stable
- ☐ (C) Mean of B is heavily influenced by an outlier
- ☐ (D) Median of A is larger than B

Q10. The deciles divide the data into:

- ☐ (A) 4 equal parts
- ☐ (B) 5 equal parts
- ☒ (C) 10 equal parts
- ☐ (D) 100 equal parts