

## Vector Calculus

### 1) What is nabla?

**Ans.** Del or **nabla**, is an operator used in mathematics, in particular in vector calculus, as a vector differential operator, usually represented by the **nabla** symbol  $\nabla$ .

$$\left[ \nabla = \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right]$$

2) **Gradient:**  $\text{grad } \phi = \nabla \phi = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right) \phi = \left( \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k} \right)$

3) **Divergence:** Suppose  $V(x, y, z) = V_1(x, y, z)\mathbf{i} + V_2(x, y, z)\mathbf{j} + V_3(x, y, z)\mathbf{k}$

$$\text{div } \mathbf{V} = \nabla \cdot \mathbf{V} = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right) \cdot (V_1 \mathbf{i} + V_2 \mathbf{j} + V_3 \mathbf{k}) = \frac{\partial V_1}{\partial x} + \frac{\partial V_2}{\partial y} + \frac{\partial V_3}{\partial z}$$

4) **Curl:** Suppose  $V(x, y, z) = V_1(x, y, z)\mathbf{i} + V_2(x, y, z)\mathbf{j} + V_3(x, y, z)\mathbf{k}$

$$\begin{aligned} \text{curl } \mathbf{V} &= \nabla \times \mathbf{V} = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right) \times (V_1 \mathbf{i} + V_2 \mathbf{j} + V_3 \mathbf{k}) \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ V_1 & V_2 & V_3 \end{vmatrix} = \mathbf{i} \left( \frac{\partial V_3}{\partial y} - \frac{\partial V_2}{\partial z} \right) - \mathbf{j} \left( \frac{\partial V_3}{\partial x} - \frac{\partial V_1}{\partial z} \right) + \mathbf{k} \left( \frac{\partial V_2}{\partial x} - \frac{\partial V_1}{\partial y} \right) \end{aligned}$$

**Ex:1)** If  $\phi = 3xyz - x^2y^2z^2$ , then find  $\nabla \phi$  at the point  $(1, 1, 1)$ .

**Solution:** Given,  $\phi = 3xyz - x^2y^2z^2$

$$\therefore \nabla \phi = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right) (3xyz - x^2y^2z^2)$$

$$= \mathbf{i} \frac{\partial}{\partial x} (3xyz - x^2y^2z^2) + \mathbf{j} \frac{\partial}{\partial y} (3xyz - x^2y^2z^2) + \mathbf{k} \frac{\partial}{\partial z} (3xyz - x^2y^2z^2)$$

$$= \mathbf{i}(3yz - 2xy^2z^2) + \mathbf{j}(3xz - 2yx^2z^2) + \mathbf{k}(3xy - 2zx^2y^2)$$

At the point,  $(1,1,1)$ ,  $\nabla\phi = \mathbf{i} + \mathbf{j} + \mathbf{k}$

**Ex.2) If  $\mathbf{A} = [2x^2, -3yz, xz^2]$  and  $\phi = 2z - x^3y$ , then prove at  $(1, -1, 1)$**

(i)  $\nabla \cdot \mathbf{A} = 3$                       (ii)  $\nabla \times \mathbf{A} = [-3, -1, 0]$

(iii)  $\mathbf{A} \cdot \nabla\phi = 5$                       (iv)  $\mathbf{A} \times \nabla\phi = [7, -1, -11]$

**Solution: (i)** Given,  $\mathbf{A} = [2x^2, -3yz, xz^2]$

$$\begin{aligned}\text{We know, } \nabla \cdot \mathbf{A} &= \frac{\partial A_1}{\partial x} + \frac{\partial A_2}{\partial y} + \frac{\partial A_3}{\partial z} \\ &= \frac{\partial}{\partial x}(2x^2) + \frac{\partial}{\partial y}(-3yz) + \frac{\partial}{\partial z}(xz^2) = 4x - 3z + 2xz\end{aligned}$$

At the point  $(1,-1,1)$ ,  $\nabla \cdot \mathbf{A} = 4 - 3 + 2 = 6 - 3 = 3$  (proved)

**(ii)** Given,  $\mathbf{A} = [2x^2, -3yz, xz^2]$

We know,

$$\begin{aligned}\nabla \times \mathbf{A} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x^2 & -3yz & xz^2 \end{vmatrix} \\ &= \mathbf{i} \left\{ \frac{\partial}{\partial y}(xz^2) - \frac{\partial}{\partial z}(-3yz) \right\} - \mathbf{j} \left\{ \frac{\partial}{\partial x}(xz^2) - \frac{\partial}{\partial z}(2x^2) \right\} + \mathbf{k} \left\{ \frac{\partial}{\partial x}(-3yz) - \frac{\partial}{\partial y}(2x^2) \right\} \\ &= \mathbf{i}(0 + 3z) - \mathbf{j}(z^2 - 0) + \mathbf{k}(0 - 0) = 3z\mathbf{i} - z^2\mathbf{j}\end{aligned}$$

At  $(1,-1,1)$ ,  $\nabla \times \mathbf{A} = [-3, -1, 0]$  (proved)

**(iii)** Given,  $\mathbf{A} = [2x^2, -3yz, xz^2]$  and  $\phi = 2z - x^3y$

$$\text{We know, } \nabla\phi = \mathbf{i} \frac{\partial}{\partial x}(2z - x^3y) + \mathbf{j} \frac{\partial}{\partial y}(2z - x^3y) + \mathbf{k} \frac{\partial}{\partial z}(2z - x^3y) = -3x^2y\mathbf{i} - x^3\mathbf{j} + 2\mathbf{k}$$

$$\mathbf{A} \cdot \nabla\phi = [2x^2, -3yz, xz^2] \cdot (-3x^2y\mathbf{i} - x^3\mathbf{j} + 2\mathbf{k}) = -6x^4y + 3x^3yz + 2xz^2 = 6 - 3 + 2 = 5$$

$$(iv) \nabla \phi = \mathbf{i} \frac{\partial}{\partial x} (2z - x^3y) + \mathbf{j} \frac{\partial}{\partial y} (2z - x^3y) + \mathbf{k} \frac{\partial}{\partial z} (2z - x^3y) = -3x^2y\mathbf{i} - x^3\mathbf{j} + 2\mathbf{k}$$

$$A \times \nabla \phi = [2x^2, -3yz, xz^2] \times [-3x^2y, -x^3, 2]$$

$$\Rightarrow \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2x^2 & -3yz & xz^2 \\ -3x^2y & -x^3 & 2 \end{vmatrix} = \mathbf{i}(-6yz + x^4z^2) - \mathbf{j}(4x^2 + 3x^3yz^2) + \mathbf{k}(-2x^5 - 9x^2y^2z^2)$$

$$= 7\mathbf{i} - \mathbf{j} - 11\mathbf{k}$$

**Ex.3) If  $V = [x + 3y, y - 2z, x + az]$  is solenoidal, then find the value of  $a$ .**

**Solution:** If  $V$  is solenoidal then,  $\nabla \cdot V = 0$

$$\Rightarrow \frac{\partial}{\partial x}(x + 3y) + \frac{\partial}{\partial y}(y - 2z) + \frac{\partial}{\partial z}(x + az) = 0 \Rightarrow 1 + 1 + a = 0 \Rightarrow a = -2$$

**Ex.4) Find the unit tangent vector to any point on the curve  $x = t^2 + 1$ ,  $y = 4t - 3$ ,  $z = 2t^2 - 6t$  and determine the unit tangent at the point  $t = 2$ .**

**Solution:** Let,  $r = [x, y, z] = [t^2 + 1, 4t - 3, 2t^2 - 6t]$

$$\therefore \frac{dr}{dt} = [2t, 4, 4t - 6]$$

$$\left| \frac{dr}{dt} \right| = \sqrt{4t^2 + 16 + (4t - 6)^2}$$

$$\text{Then, the unit tangent vector} = \frac{\frac{dr}{dt}}{\left| \frac{dr}{dt} \right|} = \frac{[2t, 4, 4t - 6]}{\sqrt{4t^2 + 16 + (4t - 6)^2}}$$

At  $t=2$ ,

$$\text{The unit tangent vector} = \frac{\frac{dr}{dt}}{\left| \frac{dr}{dt} \right|} = \frac{[4,4,2]}{\sqrt{16+16+4}} = \frac{[4,4,2]}{6} = \frac{4}{6}\mathbf{i} + \frac{4}{6}\mathbf{j} + \frac{2}{6}\mathbf{k}$$

**Ex. 5) Find the directional derivative of  $f(x,y)$**

$$= \tan^{-1} \frac{y}{x} \text{ at the point } (-2, 2) \text{ in the direction}$$

**of  $\mathbf{a} = -\mathbf{i} - \mathbf{j}$ .**

**Solution:** Given,

$$f(x,y) = \tan^{-1} \frac{y}{x}$$

$$\nabla f = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} \right) \left( \tan^{-1} \frac{y}{x} \right) = \mathbf{i} \frac{1}{1 + \frac{y^2}{x^2}} \left( -\frac{y}{x^2} \right) + \mathbf{j} \frac{1}{1 + \frac{y^2}{x^2}} \frac{1}{x}$$

$$= -\frac{\mathbf{i} y}{\frac{x^2 + y^2}{x^2}} \left( \frac{1}{x^2} \right) + \frac{\mathbf{j}}{\frac{x^2 + y^2}{x^2}} \frac{1}{x} = -\frac{\mathbf{i} y}{x^2 + y^2} + \frac{\mathbf{j} x}{x^2 + y^2}$$

At the point  $(-2, 2)$ ,

$$\nabla f = -\frac{\mathbf{i} y}{x^2 + y^2} + \frac{\mathbf{j} x}{x^2 + y^2} = -\frac{\mathbf{i} 2}{8} + \frac{\mathbf{j} (-2)}{8} = -\frac{\mathbf{i}}{4} - \frac{\mathbf{j}}{4}$$

$$\text{Hence, the directional derivative of } f = \nabla f \frac{\mathbf{a}}{|\mathbf{a}|} = \left( -\frac{\mathbf{i}}{4} - \frac{\mathbf{j}}{4} \right) \frac{-\mathbf{i} - \mathbf{j}}{\sqrt{1+1}} = \frac{1}{\sqrt{2}} \left( \frac{1}{4} + \frac{1}{4} \right) = \frac{1}{2\sqrt{2}}$$

**Ex. 6) Find the directional derivative of  $f(x,y)$**

$$= x^2 + xy \text{ at the point } (1, 2) \text{ in the direction}$$

$$\text{of the unit vector } \hat{\mathbf{a}} = \frac{\mathbf{i}}{\sqrt{2}} + \frac{\mathbf{j}}{\sqrt{2}}$$

**Solution:** Given,

$$f(x,y) = x^2 + xy$$

$$\nabla f = \left( \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} \right) (x^2 + xy) = \mathbf{i} (2x + y) + \mathbf{j} x$$

At the point (1,2),

$$\nabla f = 4\mathbf{i} + \mathbf{j}$$

$$\text{Hence, the directional derivative of } f = \nabla f \cdot \hat{a} = (4\mathbf{i} + \mathbf{j}) \left( \frac{\mathbf{i}}{\sqrt{2}} + \frac{\mathbf{j}}{\sqrt{2}} \right) = \frac{4}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \frac{5}{\sqrt{2}}$$

**Ex. 7) Find the angle between the surfaces  $xy^2z = 3x + z^2$  and  $3x^2 - y^2 + 2z = 1$  at the point  $(1, -2, 1)$ .**

**Solution:** Let,

$$\phi = xy^2z - 3x - z^2 \text{ and } \psi = 3x^2 - y^2 + 2z - 1$$

$$\text{Angle between the surfaces, } \cos \theta = \frac{\nabla \phi \cdot \nabla \psi}{|\nabla \phi| \cdot |\nabla \psi|}$$

$$\nabla \phi = \mathbf{i}(y^2z - 3) + \mathbf{j}(2xyz) + \mathbf{k}(xy^2 - 2z)$$

$$\nabla \psi = \mathbf{i}(6x) + \mathbf{j}(-2y) + \mathbf{k}(2)$$

$$\text{At } (1, -2, 1), \nabla \phi = \mathbf{i} - 4\mathbf{j} + 2\mathbf{k} \text{ and } \nabla \psi = 6\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$$

$$\nabla \phi \cdot \nabla \psi = 6 - 16 + 4 = -6$$

$$|\nabla \phi| = \sqrt{21} \text{ and } |\nabla \psi| = 2\sqrt{14}$$

$$\text{Hence, the angle between the surfaces, } \cos \theta = \frac{\nabla \phi \cdot \nabla \psi}{|\nabla \phi| \cdot |\nabla \psi|} = \frac{-6}{\sqrt{21} \cdot 2\sqrt{14}} = \frac{-3}{\sqrt{21} \cdot \sqrt{14}}$$

$$\Rightarrow \cos \theta = \frac{-3}{\sqrt{294}} \Rightarrow \theta = \cos^{-1} \left( \frac{-3}{\sqrt{294}} \right)$$

**H.W:**

1) If  $\phi = 3x^2 - 4y^3 + z^2$ , then find  $\nabla\phi$  at the point (1,2,3) and  $|\nabla\phi|$       Ans.  $6i - 48j + 6k$

2) If  $A = 2xz^2i - yzj + 3xz^3k$ , Then find the value of  $\nabla \times (\nabla \times A)$  at (1,1,1)      Ans.  $5i+3k$

3) If  $A = 3xz^2i - yzj + (x + 2z)k$ , then find curl (curl A)      Ans.  $-6xi+(6z-1)k$

4) Find the directional derivative of  $f(x,y) = xe^y - ye^x$  at the point (0,0)      Ans.  $\frac{7}{\sqrt{29}}$

in the direction of  $a = 5i - 2j$ .

5) Find the directional derivative of  $f(x,y)$   
 $= x^3z - yz^2 + z^2$  at the point (2, -1,1)      Ans.  $5\sqrt{14}$

in the direction of  $a = 3i - j + 2k$ .

6) Find the directional derivative of  $f(x,y) = 4xz^3 - 3x^2y^2z$  at the point (2, -1,2)      Ans.  $\frac{376}{7}$

in the direction of  $a = 2i - 3j + 6k$ .

7) Find the angle between the surfaces  $x^2y + z = 3$  and  $x \ln z + y^2 = 4$       Ans.  $\theta$   
 $= \cos^{-1} \left( \frac{1}{\sqrt{34}} \right)$

at the point (-1,2,1)