Chapter 5
Random Vectors

Chien-Kang Huang (黃乾綱)
臺大工科海洋系

Outline

5.1 Probability Models of N Random Variables

5.2 Vector Notation

5.3 Marginal Probability Functions

5.4 Independence of Random Variables and Random Vectors

5.5 Functions of Random Vectors

5.6 Expected Value Vector and Correlation Matrix

5.7 Gaussian Random Vectors

5.1 Probability Models of N Random Variables

- Generalize the concepts presented in Chapter 4 to any number of random variables.
- A random vector treats a collection of n random variable as a single entity. Thus vector notation provides a concise representation of relationships that would otherwise be extremely difficult to represent.

Definition 5.1 ***Multivariate Joint CDF***

The joint CDF *of* $X_1, \ldots, X_n$ *is*

$$F_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = P\left[X_1 \leq x_1, \ldots, X_n \leq x_n\right].$$

Definition 5.2 ***Multivariate Joint PMF***

The joint PMF *of the discrete random variables* $X_1, \ldots, X_n$ *is*

$$P_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = P[X_1 = x_1, \ldots, X_n = x_n].$$

Definition 5.3 Multivariate Joint PDF

- *The* **joint PDF** *of the continuous random variable* $X_1, \ldots, X_n$ *is the function*

$$f_{X_1,\ldots,X_n}(x_1,\ldots,x_n) = \frac{\partial^n F_{X_1,\ldots,X_n}(x_1,\ldots,x_n)}{\partial x_1 \ldots \partial x_n}$$

Theorem 5.1

If $X_1, \ldots, X_n$ are discrete random variables with joint PMF $P_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$,

(a) $P_{X_1,\ldots,X_n}(x_1, \ldots, x_n) \geq 0$,

(b) $\displaystyle\sum_{x_1 \in S_{X_1}} \cdots \sum_{x_n \in S_{X_n}} P_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = 1.$

Theorem 5.2

If $X_1, \ldots, X_n$ are continuous random variables with joint PDF $f_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$,

(a) $f_{X_1,\ldots,X_n}(x_1, \ldots, x_n) \geq 0$,

(b) $F_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = \int_{-\infty}^{x_1} \cdots \int_{-\infty}^{x_n} f_{X_1,\ldots,X_n}(u_1, \ldots, u_n)\, du_1 \cdots du_n$,

(c) $\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f_{X_1,\ldots,X_n}(x_1, \ldots, x_n)\, dx_1 \cdots dx_n = 1$.

Theorem 5.3

The probability of an event A expressed in terms of the random variables $X_1, \ldots, X_n$ is

multiple sum

Discrete: $$P[A] = \sum_{(x_1,\ldots,x_n)\in A} P_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$$

Continuous: $$P[A] = \int \cdots \int_A f_{X_1,\ldots,X_n}(x_1, \ldots, x_n)\, dx_1\, dx_2 \ldots dx_n.$$

- Often we consider an event A described in terms of a property of X_1, …, X_n , such as $|X_1 + X_2 + \ldots + X_n| < 1$, or $\max_i X_i \leq 100$. To find the probability of the event A, we sum the joint PMF or integrate the joint PDF over all x_1, …, x_n that belong to A.
- Although, we have written the discrete version with a single summation, we must remember that in fact it is a multiple sum over the n variables.

Example 5.1 Problem

Consider a set of n independent trials in which there are r possible outcomes $s_1, \ldots, s_r$ for each trial. In each trial, $P[s_i] = p_i$. Let N_i equal the number of times that outcome s_i occurs over n trials. What is the joint PMF of $N_1, \ldots, N_r$?

Example 5.1 Solution

The solution to this problem appears in Theorem 1.19 and is repeated here:

$$P_{N_1,\ldots,N_r}(n_1, \ldots, n_r) = \binom{n}{n_1, \ldots, n_r} p_1^{n_1} p_2^{n_2} \cdots p_r^{n_r}.$$

Example 5.2 Problem

In response to requests for information, a company sends faxes that can be 1, 2, or 3 pages in length, depending on the information requested. The PMF of L, the length of one fax is

$$P_L(l) = \begin{cases} 1/3 & l = 1, \\ 1/2 & l = 2, \\ 1/6 & l = 3, \\ 0 & \text{otherwise.} \end{cases}$$

For a set of four independent information requests:

(a) What is the joint PMF of the random variables, X, Y,and Z, the number of 1-page, 2-page, and 3-page faxes, respectively?

(b) What is $P[A] = P[$total length of four faxes is 8 pages$]$?

(c) What is $P[B] = P[$at least half of the four faxes has more than 1 page$]$?

Example 5.2 Solution

Each fax sent is an independent trial with three possible outcomes: $L = 1$, $L = 2$, and $L = 3$. Hence, the number of faxes of each length out of four faxes is described by the multinomial PMF of Example 5.1:

$$P_{X,Y,Z}(x, y, z) = \binom{4}{x, y, z} \left(\frac{1}{3}\right)^x \left(\frac{1}{2}\right)^y \left(\frac{1}{6}\right)^z.$$

The PMF is displayed numerically in Table 5.1. The final column of the table indicates that there are three outcomes in event A and 12 outcomes in event B. Adding the probabilities in the two events, we have $P[A] = 107/432$ and $P[B] = 8/9$.

Table 5.1

x (1 page)	y (2 pages)	z (3 pages)	$P_{X,Y,Z}(x, y, z)$	total pages	events
0	0	4	$1/1296$	12	B
0	1	3	$1/108$	11	B
0	2	2	$1/24$	10	B
0	3	1	$1/12$	9	B
0	4	0	$1/16$	8	AB
1	0	3	$1/162$	10	B
1	1	2	$1/18$	9	B
1	2	1	$1/6$	8	AB
1	3	0	$1/6$	7	B
2	0	2	$1/54$	8	AB
2	1	1	$1/9$	7	B
2	2	0	$1/6$	6	B
3	0	1	$2/81$	6	
3	1	0	$2/27$	5	
4	0	0	$1/81$	4	

The PMF $P_{X,Y,Z}(x, y, z)$ and the events A and B for Example 5.2.

Example 5.3 Problem

The random variables $X_1, \ldots, X_n$ have the joint PDF

$$f_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = \begin{cases} 1 & 0 \le x_i \le 1, i = 1, \ldots, n, \\ 0 & \text{otherwise.} \end{cases}$$

Let A denote the event that $\max_i X_i \le 1/2$. Find $P[A]$.

Example 5.3 Solution

$$P[A] = P\left[\max_i X_i \le 1/2\right] = P[X_1 \le 1/2, \ldots, X_n \le 1/2]$$
$$= \int_0^{1/2} \cdots \int_0^{1/2} 1\, dx_1 \cdots dx_n = \frac{1}{2^n}.$$

Here we have n independent uniform $(0, 1)$ random variables. As n grows, the probability that the maximum is less than $1/2$ rapidly goes to 0.

Quiz 5.1

The random variables $Y_1, \ldots, Y_4$ have the joint PDF

$$f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4) = \begin{cases} 4 & 0 \le y_1 \le y_2 \le 1, 0 \le y_3 \le y_4 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let C denote the event that $\max_i Y_i \le 1/2$. Find $P[C]$.

Quiz 5.1 Solution

We find $P[C]$ by integrating the joint PDF over the region of interest. Specifically,

$$
\begin{aligned}
P[C] &= \int_0^{1/2} dy_2 \int_0^{y_2} dy_1 \int_0^{1/2} dy_4 \int_0^{y_4} 4 dy_3 \\
&= 4\left(\int_0^{1/2} y_2\, dy_2\right)\left(\int_0^{1/2} y_4\, dy_4\right) = 1/16
\end{aligned}
$$

5.2 Vector Notation

- We use boldface notation $\mathbf{x}$ for a column vector.
- Row vectors are transpose column vectors; $\mathbf{x}'$ is a row vector.

Definition 5.4 Random Vector

A random vector *is a column vector* $\mathbf{X} = \begin{bmatrix} X_1 & \cdots & X_n \end{bmatrix}'$. *Each* X_i *is a random variable.*

- A random variable is a random vector with $n = 1$.

Definition 5.5 Vector Sample Value

A sample value of a random vector *is a column vector* $\mathbf{x} = \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}'$. *The* i*th component,* x_i*, of the vector* $\mathbf{x}$ *is a sample value of a random variable,* X_i*.*

- Following out convention for random variables, the uppercase $\mathbf{X}$ is the random vector and the lowercase $\mathbf{x}$ is a sample value of $\mathbf{X}$.
- However, we also use boldface capitals such as $\mathbf{A}$ and $\mathbf{B}$ to denote matrices with components that are not random variables. It will be clear from the context whether $\mathbf{A}$ is a matrix of numbers, a matrix of random variables, or a random vector.

Definition 5.6 Functions

(a) *The* CDF of a random vector $\mathbf{X}$ *is* $F_{\mathbf{X}}(\mathbf{x}) = F_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$.

(b) *The* PMF of a discrete random vector $\mathbf{X}$ *is* $P_{\mathbf{X}}(\mathbf{x}) = P_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$.

(c) *The* PDF of a continuous random vector $\mathbf{X}$ *is* $f_{\mathbf{X}}(\mathbf{x}) = f_{X_1,\ldots,X_n}(x_1, \ldots, x_n)$.

- We use similar notation for a function $g(\mathbf{X}) = g(X_1, \ldots, X_n)$ of n random variables and a function $g(\mathbf{x}) = g(x_1, \ldots, x_n)$ of n numbers.

Definition 5.7 Random Vectors

For random vectors $\mathbf{X}$ *with* n *components and* $\mathbf{Y}$ *with* m *components:*

(a) *The* joint CDF *of* $\mathbf{X}$ *and* $\mathbf{Y}$ *is*

$$F_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) = F_{X_1,\ldots,X_n,Y_1,\ldots,Y_m}(x_1,\ldots,x_n,y_1,\ldots,y_m);$$

(b) *The* joint PMF *of discrete random vectors* $\mathbf{X}$ *and* $\mathbf{Y}$ *is*

$$P_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) = P_{X_1,\ldots,X_n,Y_1,\ldots,Y_m}(x_1,\ldots,x_n,y_1,\ldots,y_m);$$

(c) *The* joint PDF *of continuous random vectors* $\mathbf{X}$ *and* $\mathbf{Y}$ *is*

$$f_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) = f_{X_1,\ldots,X_n,Y_1,\ldots,Y_m}(x_1,\ldots,x_n,y_1,\ldots,y_m).$$

- The logic of Definition 5.7 is that the pair of random vectors $\mathbf{X}$ and $\mathbf{Y}$ is the same as $\mathbf{W} = [\mathbf{X}' \ \mathbf{Y}']' = [X_1 \ \ldots \ X_n \ \ Y_1 \ \ldots \ Y_m]'$, a concatenation of $\mathbf{X}$ and $\mathbf{Y}$. Thus a probability function of the pair $\mathbf{X}$ and $\mathbf{Y}$ corresponds to the same probability function of $\mathbf{W}$; for example, $F_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y})$ is the same CDF as $F_{\mathbf{W}}(\mathbf{w})$.

Example 5.4 Problem

Random vector $\mathbf{X}$ has PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} 6e^{-\mathbf{a}'\mathbf{x}} & \mathbf{x} \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

where $\mathbf{a} = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}'$. What is the CDF of $\mathbf{X}$?

Example 5.4 Solution

Because $\mathbf{a}$ has three components, we infer that $\mathbf{X}$ is a 3-dimensional random vector. Expanding $\mathbf{a}'\mathbf{x}$, we write the PDF as a function of the vector components,

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} 6e^{-x_1-2x_2-3x_3} & x_i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Applying Definition 5.7, we integrate the PDF with respect to the three variables to obtain

$$F_{\mathbf{X}}(\mathbf{x}) = \begin{cases} (1-e^{-x_1})(1-e^{-2x_2})(1-e^{-3x_3}) & x_i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Quiz 5.2

Discrete random vectors $\mathbf{X} = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix}'$ and $\mathbf{Y} = \begin{bmatrix} y_1 & y_2 & y_3 \end{bmatrix}'$ are related by $\mathbf{Y} = \mathbf{A}\mathbf{X}$. Find the joint PMF $P_{\mathbf{Y}}(\mathbf{y})$ if $\mathbf{X}$ has joint PMF

$$P_{\mathbf{X}}(\mathbf{x}) = \begin{cases} (1-p)\,p^{x_3} & x_1 < x_2 < x_3; \\ & x_1, x_2, x_3 \in \{1, 2, \ldots\}, \\ 0 & \text{otherwise,} \end{cases} \quad \mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}.$$

Quiz 5.2 Solution

By definition of $\mathbf{A}$, $Y_1 = X_1$, $Y_2 = X_2 - X_1$ and $Y_3 = X_3 - X_2$. Since $0 < X_1 < X_2 < X_3$, each Y_i must be a strictly positive integer. Thus, for $y_1, y_2, y_3 \in \{1, 2, \ldots\}$,

$$\begin{aligned} P_{\mathbf{Y}}(\mathbf{y}) &= P\left[Y_1 = y_1, Y_2 = y_2, Y_3 = y_3\right] \\ &= P\left[X_1 = y_1, X_2 - X_1 = y_2, X_3 - X_2 = y_3\right] \\ &= P\left[X_1 = y_1, X_2 = y_2 + y_1, X_3 = y_3 + y_2 + y_1\right] \\ &= (1-p)^3 p^{y_1+y_2+y_3} \end{aligned}$$

By defining the vector $\mathbf{a} = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}'$, the complete expression for the joint PMF of $\mathbf{Y}$ is

$$P_{\mathbf{Y}}(\mathbf{y}) = \begin{cases} (1-p)\, p^{\mathbf{a}'\mathbf{y}} & y_1, y_2, y_3 \in \{1, 2, \ldots\} \\ 0 & \text{otherwise} \end{cases}$$

5.3 Marginal Probability Functions

Theorem 5.4

For a joint PMF $P_{W,X,Y,Z}(w, x, y, z)$ of discrete random variables W, X, Y, Z, some marginal PMFs are

$$P_{X,Y,Z}(x, y, z) = \sum_{w \in S_W} P_{W,X,Y,Z}(w, x, y, z),$$

$$P_{W,Z}(w, z) = \sum_{x \in S_X} \sum_{y \in S_Y} P_{W,X,Y,Z}(w, x, y, z),$$

$$P_X(x) = \sum_{w \in S_W} \sum_{y \in S_Y} \sum_{z \in S_Z} P_{W,X,Y,Z}(w, x, y, z).$$

- Consider an experiment with 4 random variables W, X, Y, Z.

Theorem 5.5

For a joint PDF $f_{W,X,Y,Z}(w, x, y, z)$ of continuous random variables W, X, Y, Z, some marginal PDFs are

$$f_{X,Y,Z}(x, y, z) = \int_{-\infty}^{\infty} f_{W,X,Y,Z}(w, x, y, z)\, dw,$$

$$f_{W,Z}(w, z) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{W,X,Y,Z}(w, x, y, z)\, dx\, dy,$$

$$f_X(x) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{W,X,Y,Z}(w, x, y, z)\, dw\, dy\, dz.$$

- Theorems 5.4 and 5.5 can be generalized in a straightforward way to any marginal PMF or marginal PDF or an arbitrary number of random variables.
- For a probability model described by the set of random variables $\{X_1, \ldots, X_n\}$, each nonempty strict subset of those random variables has a marginal probability model.
- There are 2^n subsets of $\{X_1, \ldots, X_n\}$. After excluding the entire set and the null set ϕ, we can find that there are $2^n - 2$ marginal probability models.

Example 5.5 Problem

As in Quiz 5.1, the random variables $Y_1, \ldots, Y_4$ have the joint PDF

$$f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4) = \begin{cases} 4 & 0 \le y_1 \le y_2 \le 1, 0 \le y_3 \le y_4 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find the marginal PDFs $f_{Y_1,Y_4}(y_1, y_4)$, $f_{Y_2,Y_3}(y_2, y_3)$, and $f_{Y_3}(y_3)$.

Example 5.5 Solution

$$f_{Y_1,Y_4}(y_1, y_4) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4)\, dy_2\, dy_3.$$

In the foregoing integral, the hard part is identifying the correct limits. These limits will depend on y_1 and y_4. For $0 \le y_1 \le 1$ and $0 \le y_4 \le 1$,

$$f_{Y_1,Y_4}(y_1, y_4) = \int_{y_1}^{1}\int_{0}^{y_4} 4\, dy_3\, dy_2 = 4(1 - y_1)y_4.$$

The complete expression for $f_{Y_1,Y_4}(y_1, y_4)$ is

$$f_{Y_1,Y_4}(y_1, y_4) = \begin{cases} 4(1 - y_1)y_4 & 0 \le y_1 \le 1, 0 \le y_4 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Similarly, for $0 \le y_2 \le 1$ and $0 \le y_3 \le 1$,

$$f_{Y_2,Y_3}(y_2, y_3) = \int_{0}^{y_2}\int_{y_3}^{1} 4\, dy_4\, dy_1 = 4y_2(1 - y_3).$$

[Continued]

Example 5.5 Solution (continued)

The complete expression for $f_{Y_2,Y_3}(y_2, y_3)$ is

$$f_{Y_2,Y_3}(y_2, y_3) = \begin{cases} 4y_2(1-y_3) & 0 \le y_2 \le 1, 0 \le y_3 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Lastly, for $0 \le y_3 \le 1$,

$$f_{Y_3}(y_3) = \int_{-\infty}^{\infty} f_{Y_2,Y_3}(y_2, y_3)\, dy_2 = \int_0^1 4y_2(1-y_3)\, dy_2 = 2(1-y_3).$$

The complete expression is

$$f_{Y_3}(y_3) = \begin{cases} 2(1-y_3) & 0 \le y_3 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Quiz 5.3

The random vector $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$ has PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} 6 & 0 \leq x_1 \leq x_2 \leq x_3 \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find the marginal PDFs $f_{X_1,X_2}(x_1, x_2)$, $f_{X_1,X_3}(x_1, x_3)$, $f_{X_2,X_3}(x_2, x_3)$, and $f_{X_1}(x_1)$, $f_{X_2}(x_2)$, $f_{X_3}(x_3)$.

Quiz 5.3 Solution

First we note that each marginal PDF is nonzero only if any subset of the x_i obeys the ordering contraints $0 \le x_1 \le x_2 \le x_3 \le 1$. Within these constraints, we have

$$f_{X_1,X_2}(x_1,x_2) = \int_{-\infty}^{\infty} f_{\mathbf{X}}(\mathbf{x})\, dx_3 = \int_{x_2}^{1} 6\, dx_3 = 6(1-x_2),$$
$$f_{X_2,X_3}(x_2,x_3) = \int_{-\infty}^{\infty} f_{\mathbf{X}}(\mathbf{x})\, dx_1 = \int_{0}^{x_2} 6\, dx_1 = 6x_2,$$
$$f_{X_1,X_3}(x_1,x_3) = \int_{-\infty}^{\infty} f_{\mathbf{X}}(\mathbf{x})\, dx_2 = \int_{x_1}^{x_3} 6\, dx_2 = 6(x_3-x_1).$$

In particular, we must keep in mind that $f_{X_1,X_2}(x_1,x_2) = 0$ unless $0 \le x_1 \le x_2 \le 1$, $f_{X_2,X_3}(x_2,x_3) = 0$ unless $0 \le x_2 \le x_3 \le 1$, and that $f_{X_1,X_3}(x_1,x_3) = 0$ unless $0 \le x_1 \le x_3 \le 1$. The complete expressions are

$$f_{X_1,X_2}(x_1,x_2) = \begin{cases} 6(1-x_2) & 0 \le x_1 \le x_2 \le 1 \\ 0 & \text{otherwise} \end{cases}$$
$$f_{X_2,X_3}(x_2,x_3) = \begin{cases} 6x_2 & 0 \le x_2 \le x_3 \le 1 \\ 0 & \text{otherwise} \end{cases}$$
$$f_{X_1,X_3}(x_1,x_3) = \begin{cases} 6(x_3-x_1) & 0 \le x_1 \le x_3 \le 1 \\ 0 & \text{otherwise} \end{cases}$$

[Continued]

Quiz 5.3 Solution (continued)

Now we can find the marginal PDFs. When $0 \leq x_i \leq 1$ for each x_i,

$$f_{X_1}(x_1) = \int_{-\infty}^{\infty} f_{X_1,X_2}(x_1, x_2)\, dx_2 = \int_{x_1}^{1} 6(1 - x_2)\, dx_2 = 3(1 - x_1)^2$$

$$f_{X_2}(x_2) = \int_{-\infty}^{\infty} f_{X_2,X_3}(x_2, x_3)\, dx_3 = \int_{x_2}^{1} 6x_2\, dx_3 = 6x_2(1 - x_2)$$

$$f_{X_3}(x_3) = \int_{-\infty}^{\infty} f_{X_2,X_3}(x_2, x_3)\, dx_2 = \int_{0}^{x_3} 6x_2\, dx_2 = 3x_3^2$$

The complete expressions are

$$f_{X_1}(x_1) = \begin{cases} 3(1 - x_1)^2 & 0 \leq x_1 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$f_{X_2}(x_2) = \begin{cases} 6x_2(1 - x_2) & 0 \leq x_2 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$f_{X_3}(x_3) = \begin{cases} 3x_3^2 & 0 \leq x_3 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

5.4 Independence of Random Variables and Random Vectors

Definition 5.8 ***N Independent Random Variables***

Random variables $X_1, \ldots, X_n$ *are* independent *if for all* $x_1, \ldots, x_n$,

Discrete: $$P_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = P_{X_1}(x_1) P_{X_2}(x_2) \cdots P_{X_N}(x_n)$$

Continuous: $$f_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = f_{X_1}(x_1) f_{X_2}(x_2) \cdots f_{X_n}(x_n).$$

Example 5.6 Problem

As in Example 5.5, random variables $Y_1, \ldots, Y_4$ have the joint PDF

$$f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4) = \begin{cases} 4 & 0 \leq y_1 \leq y_2 \leq 1, 0 \leq y_3 \leq y_4 \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Are $Y_1, \ldots, Y_4$ independent random variables?

Example 5.6 Solution

In Equation (5.15) of Example 5.5, we found the marginal PDF $f_{Y_1,Y_4}(y_1, y_4)$. We can use this result to show that

$$f_{Y_1}(y_1) = \int_0^1 f_{Y_1,Y_4}(y_1, y_4)\, dy_4 = 2(1 - y_1), \qquad 0 \le y_1 \le 1,$$

$$f_{Y_4}(y_4) = \int_0^1 f_{Y_1,Y_4}(y_1, y_4)\, dy_1 = 2y_4, \qquad 0 \le y_4 \le 1.$$

The full expressions for the marginal PDFs are

$$f_{Y_1}(y_1) = \begin{cases} 2(1 - y_1) & 0 \le y_1 \le 1, \\ 0 & \text{otherwise}, \end{cases}$$

$$f_{Y_4}(y_4) = \begin{cases} 2y_4 & 0 \le y_4 \le 1, \\ 0 & \text{otherwise}. \end{cases}$$

Similarly, the marginal PDF $f_{Y_2,Y_3}(y_2, y_3)$ found in Equation (5.17) of Example 5.5 implies that for $0 \le y_2 \le 1$,

$$f_{Y_2}(y_2) = \int_{-\infty}^{\infty} f_{Y_2,Y_3}(y_2, y_3)\, dy_3 = \int_0^1 4y_2(1 - y_3)\, dy_3 = 2y_2$$

[Continued]

Example 5.6 Solution (continued)

It follows that the marginal PDF of Y_2 is

$$f_{Y_2}(y_2) = \begin{cases} 2y_2 & 0 \le y_2 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

From Equation (5.19) for the PDF $f_{Y_3}(y_3)$ derived in Example 5.5, we have

$$\begin{aligned} f_{Y_1}(y_1) f_{Y_2}(y_2) f_{Y_3}(y_3) f_{Y_4}(y_4) \\ &= \begin{cases} 16(1-y_1)y_2(1-y_3)y_4 & 0 \le y_1, y_2, y_3, y_4 \le 1, \\ 0 & \text{otherwise,} \end{cases} \\ &\neq f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4). \end{aligned} \tag{1}$$

Therefore $Y_1, \ldots, Y_4$ are not independent random variables.

- Independence of n random variables is typically a property of an experiment consisting of n independent subexperiments. In this case, subexperiment i produces the random variable X_i. If all subexperiments follow the same procedure, all of the have the same PMF or PDF. In this case, we say the random variables are **identically distributed**.

Independent and Identically Distributed (iid)

Definition 5.9

Random variables $X_1, \ldots, X_n$ *are* independent and identically distributed (iid) *if*

Discrete: $$P_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = P_X(x_1)P_X(x_2)\cdots P_X(x_n)$$

Continuous: $$f_{X_1,\ldots,X_n}(x_1, \ldots, x_n) = f_X(x_1)f_X(x_2)\cdots f_X(x_n).$$

Definition 5.10 *Independent Random Vectors*

Random vectors $\mathbf{X}$ *and* $\mathbf{Y}$ *are independent if*

Discrete: $$P_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) = P_{\mathbf{X}}(\mathbf{x})\,P_{\mathbf{Y}}(\mathbf{y})$$

Continuous: $$f_{\mathbf{X},\mathbf{Y}}(\mathbf{x},\mathbf{y}) = f_{\mathbf{X}}(\mathbf{x})\,f_{\mathbf{Y}}(\mathbf{y}).$$

Example 5.7 Problem

As in Example 5.5, random variables $Y_1, \ldots, Y_4$ have the joint PDF

$$f_{Y_1,\ldots,Y_4}(y_1, \ldots, y_4) = \begin{cases} 4 & 0 \leq y_1 \leq y_2 \leq 1, 0 \leq y_3 \leq y_4 \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\mathbf{V} = \begin{bmatrix} Y_1 & Y_4 \end{bmatrix}'$ and $\mathbf{W} = \begin{bmatrix} Y_2 & Y_3 \end{bmatrix}'$. Are $\mathbf{V}$ and $\mathbf{W}$ independent random vectors?

Example 5.7 Solution

We first note that the components of $\mathbf{V}$ are $V_1 = Y_1$, and $V_2 = Y_4$. Also, $W_1 = Y_2$, and $W_2 = Y_3$. Therefore,

$$f_{\mathbf{V},\mathbf{W}}(\mathbf{v}, \mathbf{w}) = f_{Y_1,\ldots,Y_4}(v_1, w_1, w_2, v_2) = \begin{cases} 4 & 0 \le v_1 \le w_1 \le 1; \\ & 0 \le w_2 \le v_2 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Since $\mathbf{V} = \begin{bmatrix} Y_1 & Y_4 \end{bmatrix}'$ and $\mathbf{W} = \begin{bmatrix} Y_2 & Y_3 \end{bmatrix}'$,

$$f_{\mathbf{V}}(\mathbf{v}) = f_{Y_1,Y_4}(v_1, v_2) \qquad f_{\mathbf{W}}(\mathbf{w}) = f_{Y_2,Y_3}(w_1, w_2)$$

In Example 5.5. we found $f_{Y_1,Y_4}(y_1, y_4)$ and $f_{Y_2,Y_3}(y_2, y_3)$ in Equations (5.15) and (5.17). From these marginal PDFs, we have

$$f_{\mathbf{V}}(\mathbf{v}) = \begin{cases} 4(1 - v_1)v_2 & 0 \le v_1, v_2 \le 1, \\ 0 & \text{otherwise,} \end{cases}$$

$$f_{\mathbf{W}}(\mathbf{w}) = \begin{cases} 4w_1(1 - w_2) & 0 \le w_1, w_2 \le 1, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore,

$$f_{\mathbf{V}}(\mathbf{v})\, f_{\mathbf{W}}(\mathbf{w}) = \begin{cases} 16(1 - v_1)v_2w_1(1 - w_2) & 0 \le v_1, v_2, w_1, w_2 \le 1, \\ 0 & \text{otherwise,} \end{cases}$$

which is not equal to $f_{\mathbf{V},\mathbf{W}}(\mathbf{v}, \mathbf{w})$. Therefore $\mathbf{V}$ and $\mathbf{W}$ are not independent.

Quiz 5.4

Use the components of $\mathbf{Y} = \left[Y_1, \ldots, Y_4\right]'$ in Example 5.7 to construct two independent random vectors $\mathbf{V}$ and $\mathbf{W}$. Prove that $\mathbf{V}$ and $\mathbf{W}$ are independent.

Quiz 5.4 Solution

In the PDF $f_{\mathbf{Y}}(\mathbf{y})$, the components have dependencies as a result of the ordering constraints $Y_1 \leq Y_2$ and $Y_3 \leq Y_4$. We can separate these constraints by creating the vectors

$$\mathbf{V} = \begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix}, \qquad \mathbf{W} = \begin{bmatrix} Y_3 \\ Y_4 \end{bmatrix}.$$

The joint PDF of $\mathbf{V}$ and $\mathbf{W}$ is

$$f_{\mathbf{V},\mathbf{W}}(\mathbf{v}, \mathbf{w}) = \begin{cases} 4 & 0 \leq v_1 \leq v_2 \leq 1, 0 \leq w_1 \leq w_2 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

5.5 Functions of Random Vectors

- Just as we did for one random variable and two random variables, we can derive a random variable $W = g(\mathbf{X})$ that is a function of an arbitrary number of random variables.
- If W is discrete, the probability model can be calculate as $P_W(w)$, the probability of the event $A = \{W = w\}$ in Theorem 5.3.
- If W is continuous, the probability model can be expressed as $F_W(w) = P[W \leq w]$.

Theorem 5.6

For random variable $W = g(\mathbf{X})$,

Discrete: $$P_W(w) = P[W = w] = \sum_{\mathbf{x}: g(\mathbf{x}) = w} P_{\mathbf{X}}(\mathbf{x})$$

Continuous: $$F_W(w) = P[W \leq w] = \int \cdots \int_{g(\mathbf{x}) \leq w} f_{\mathbf{X}}(\mathbf{x})\, dx_1 \cdots dx_n.$$

Example 5.8 Problem

Consider an experiment that consists of spinning the pointer on the wheel of circumference 1 meter in Example 3.1 n times and observing Y_n meters, the maximum position of the pointer in the n spins. Find the CDF and PDF of Y_n.

Example 5.8 Solution

If X_i is the position of the pointer on the ith spin, then $Y_n = \max\{X_1, X_2, \ldots, X_n\}$. As a result, $Y_n \leq y$ if and only if each $X_i \leq y$. This implies

$$P[Y_n \leq y] = P[X_1 \leq y, X_2 \leq y, \ldots X_n \leq y].$$

If we assume the spins to be independent, the events $\{X_1 \leq y\}$, $\{X_2 \leq y\}$, $\ldots$, $\{X_n \leq y\}$ are independent events. Thus

$$P[Y_n \leq y] = P[X_1 \leq y] \cdots P[X_n \leq y] = (P[X \leq y])^n = (F_X(y))^n.$$

Example 3.2 derives that $F_X(x) = x$ for $0 \leq x < 1$. Furthermore, $F_X(x) = 0$ for $x < 0$ and $F_X(x) = 1$ for $x \geq 1$ since $0 \leq X \leq 1$. Therefore, since the CDF of Y_n is $F_{Y_n}(y) = (F_X(y))^n$, we can write the CDF and corresponding PDF as

$$F_{Y_n}(y) = \begin{cases} 0 & y < 0, \\ y^n & 0 \leq y \leq 1, \\ 1 & y > 1, \end{cases} \qquad f_{Y_n}(y) = \begin{cases} ny^{n-1} & 0 \leq y \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 5.7

Let $\mathbf{X}$ be a vector of n iid random variables each with CDF $F_X(x)$ and PDF $f_X(x)$.

(a) The CDF and the PDF of $Y = \max\{X_1, \ldots, X_n\}$ are

$$F_Y(y) = (F_X(y))^n, \qquad f_Y(y) = n(F_X(y))^{n-1} f_X(y).$$

(b) The CDF and the PDF of $W = \min\{X_1, \ldots, X_n\}$ are

$$F_W(w) = 1 - (1 - F_X(w))^n, \qquad f_W(w) = n(1 - F_X(w))^{n-1} f_X(w).$$

- This theorem is a generalization of Example 5.8. It expresses the PDF of the maximum and minimum values of a sequence of iid random variables in terms of the CDF and PDF of the individual random variables.

Proof: Theorem 5.7

By definition, $f_Y(y) = P[Y \leq y]$. Because Y is the maximum value of $\{X_1, \ldots, X_n\}$, the event $\{Y \leq y\} = \{X_1 \leq y, X_2 \leq y, \ldots, X_n \leq y\}$. Because all the random variables X_i are iid, $\{Y \leq y\}$ is the intersection of n independent events. Each of the events $\{X_i \leq y\}$ has probability $F_X(y)$. The probability of the intersection is the product of the individual probabilities, which implies the first part of the theorem: $F_Y(y) = (F_X(y))^n$. The second part is the result of differentiating $F_Y(y)$ with respect to y. The derivations of $F_W(w)$ and $f_W(w)$ are similar. They begin with the observations that $F_W(w) = 1 - P[W > w]$ and that the event $\{W > w\} = \{X_1 > w, X_2 > w, \ldots X_n > w\}$, which is the intersection of n independent events, each with probability $1 - F_X(w)$.

Theorem 5.8

For a random vector $\mathbf{X}$, the random variable $g(\mathbf{X})$ has expected value

Discrete: $$E[g(\mathbf{X})] = \sum_{x_1 \in S_{X_1}} \cdots \sum_{x_n \in S_{X_n}} g(\mathbf{x}) P_{\mathbf{X}}(\mathbf{x})$$

Continuous: $$E[g(\mathbf{X})] = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) \, dx_1 \cdots dx_n.$$

- In some applications of probability theory, we are interested only in the expected value of a function, not the complete probability model. Although we can always find $E[W]$ by first deriving $P_W(w)$ or $f_W(w)$, it is easier to find $E[W]$ by applying this theorem.
- If $W = g(\mathbf{X})$ is the product of n univariate functions and the components of $\mathbf{X}$ are mutually independent, $E[W]$ is a product of n expected values. (Theorem 5.9)

Theorem 5.9

When the components of $\mathbf{X}$ are independent random variables,

$$E\left[g_1(X_1)g_2(X_2)\cdots g_n(X_n)\right] = E\left[g_1(X_1)\right]E\left[g_2(X_2)\right]\cdots E\left[g_n(X_n)\right].$$

Proof: Theorem 5.9

When $\mathbf{X}$ is discrete, independence implies $P_{\mathbf{X}}(\mathbf{x}) = P_{X_1}(x_1) \cdots P_{X_n}(x_n)$. This implies

$$\begin{aligned} E\left[g_1(X_1) \cdots g_n(X_n)\right] &= \sum_{x_1 \in S_{X_1}} \cdots \sum_{x_n \in S_{X_n}} g_1(x_1) \cdots g_n(x_n) P_{\mathbf{X}}(\mathbf{x}) \\ &= \left(\sum_{x_1 \in S_{X_1}} g_1(x_1) P_{X_1}(x_1) \right) \cdots \left(\sum_{x_n \in S_{X_n}} g_n(x_n) P_{X_n}(x_n) \right) \\ &= E\left[g_1(X_1)\right] E\left[g_2(X_2)\right] \cdots E\left[g_n(X_n)\right]. \end{aligned}$$

The derivation is similar for independent continuous random variables.

- We have considered the case of a single random variable $W = g(\mathbf{X})$ derived from a random vector $\mathbf{X}$. More complicated experiments may yield a new random vector $\mathbf{Y}$ with components $Y_1, \ldots, Y_n$ that are functions of the components of $\mathbf{X}$: $Y_k = g_k(\mathbf{X})$. We can derive the PDF of $\mathbf{Y}$ by first finding the CDF $F_{\mathbf{Y}}(\mathbf{y})$ and then applying Theorem 5.2(b). Theorem 5.10 demonstrates this technique.

Theorem 5.10

Given the continuous random vector $\mathbf{X}$, define the derived random vector $\mathbf{Y}$ such that $Y_k = aX_k + b$ for constants $a > 0$ and b. The CDF and PDF of $\mathbf{Y}$ are

$$F_{\mathbf{Y}}(\mathbf{y}) = F_{\mathbf{X}}\left(\frac{y_1 - b}{a}, \ldots, \frac{y_n - b}{a}\right), \quad f_{\mathbf{Y}}(\mathbf{y}) = \frac{1}{a^n} f_{\mathbf{X}}\left(\frac{y_1 - b}{a}, \ldots, \frac{y_n - b}{a}\right).$$

Proof: Theorem 5.10

We observe $\mathbf{Y}$ has CDF $F_{\mathbf{Y}}(\mathbf{y}) = P[aX_1 + b \leq y_1, \ldots, aX_n + b \leq y_n]$. Since $a > 0$,

$$F_{\mathbf{Y}}(\mathbf{y}) = P\left[X_1 \leq \frac{y_1 - b}{a}, \ldots, X_n \leq \frac{y_n - b}{a}\right] = F_{\mathbf{X}}\left(\frac{y_1 - b}{a}, \ldots, \frac{y_n - b}{a}\right).$$

From Theorem 5.2(b), the joint PDF of $\mathbf{Y}$ is

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{\partial^n F_{Y_1,\ldots,Y_n}(y_1, \ldots, y_n)}{\partial y_1 \cdots \partial y_n} = \frac{1}{a^n} f_{\mathbf{X}}\left(\frac{y_1 - b}{a}, \ldots, \frac{y_n - b}{a}\right).$$

Theorem 5.11

If $\mathbf{X}$ is a continuous random vector and $\mathbf{A}$ is an invertible matrix, then $\mathbf{Y} = \mathbf{AX} + \mathbf{b}$ has PDF

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{1}{|\det(\mathbf{A})|} f_{\mathbf{X}}\left(\mathbf{A}^{-1}(\mathbf{y} - \mathbf{b})\right)$$

- Theorem 5.10 is a special case of a transformation of the form $\mathbf{Y} = \mathbf{AX} + \mathbf{b}$.
- This theorem is a consequence of the change-of-variable theorem (Appendix B, Math Fact B.13) in multivariable calculus.

Proof: Theorem 5.11

Let $B = \{\mathbf{y}|\mathbf{y} \leq \tilde{\mathbf{y}}\}$ so that $F_{\mathbf{Y}}(\tilde{\mathbf{y}}) = \int_B f_{\mathbf{Y}}(\mathbf{y})\, d\mathbf{y}$. Define the vector transformation $\mathbf{x} = T(\mathbf{y}) = \mathbf{A}^{-1}(\mathbf{y} - \mathbf{b})$. It follows that $\mathbf{Y} \in B$ if and only if $\mathbf{X} \in T(B)$, where $T(B) = \{\mathbf{x}|\mathbf{A}\mathbf{x} + \mathbf{b} \leq \tilde{\mathbf{y}}\}$ is the image of B under transformation T. This implies

$$F_{\mathbf{Y}}\left(\tilde{\mathbf{y}}\right) = P\left[\mathbf{X} \in T(B)\right] = \int_{T(B)} f_{\mathbf{X}}(\mathbf{x})\; d\mathbf{x}$$

By the change-of-variable theorem (Math Fact B.13),

$$F_{\mathbf{Y}}\left(\tilde{\mathbf{y}}\right) = \int_B f_{\mathbf{X}}\left(\mathbf{A}^{-1}(\mathbf{y} - \mathbf{b})\right) \left|\det\left(\mathbf{A}^{-1}\right)\right|\, d\mathbf{y}$$

where $|\det(\mathbf{A}^{-1})|$ is the absolute value of the determinant of $\mathbf{A}^{-1}$. Definition 5.6 for the CDF and PDF of a random vector combined with Theorem 5.2(b) imply that $f_{\mathbf{Y}}(\mathbf{y}) = f_{\mathbf{X}}(\mathbf{A}^{-1}(\mathbf{y} - \mathbf{b}))|\det(\mathbf{A}^{-1})|$. The theorem follows since $|\det(\mathbf{A}^{-1})| = 1/|\det(\mathbf{A})|$.

Quiz 5.5(A)

A test of light bulbs produced by a machine has three possible outcomes: L, long life; A, average life; and R, reject. The results of different tests are independent. All tests have the following probability model: $P[L] = 0.3$, $P[A] = 0.6$, and $P[R] = 0.1$. Let X_1, X_2, and X_3 be the number of light bulbs that are L, A, and R respectively in five tests. Find the PMF $P_{\mathbf{X}}(\mathbf{x})$; the marginal PMFs $P_{X_1}(x_1)$, $P_{X_2}(x_2)$, and $P_{X_3}(x_3)$; and the PMF of $W = \max(X_1, X_2, X_3)$.

Quiz 5.5(A) Solution

Referring to Theorem 1.19, each test is a subexperiment with three possible outcomes: L, A and R. In five trials, the vector $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$ indicating the number of outcomes of each subexperiment has the multinomial PMF

$$P_{\mathbf{X}}(\mathbf{x}) = \begin{cases} \binom{5}{x_1, x_2, x_3}(0.3)^{x_1}(0.6)^{x_2}(0.1)^{x_3} & x_1 + x_2 + x_3 = 5; \\ & x_1, x_2, x_3 \in \{0, 1, \ldots, 5\} \\ 0 & \text{otherwise} \end{cases}$$

We can find the marginal PMF for each X_i from the joint PMF $P_{\mathbf{X}}(\mathbf{x})$; however it is simpler to just start from first principles and observe that X_1 is the number of occurrences of L in five independent tests. If we view each test as a trial with success probability $P[L] = 0.3$, we see that X_1 is a binomial $(n, p) = (5, 0.3)$ random variable. Similarly, X_2 is a binomial $(5, 0.6)$ random variable and X_3 is a binomial $(5, 0.1)$ random variable. That is, for $p_1 = 0.3$, $p_2 = 0.6$ and $p_3 = 0.1$,

$$P_{X_i}(x) = \begin{cases} \binom{5}{x} p_i^x (1 - p_i)^{5-x} & x = 0, 1, \ldots, 5 \\ 0 & \text{otherwise} \end{cases}$$

[Continued]

Quiz 5.5(A) Solution (continued)

From the marginal PMFs, we see that X_1, X_2 and X_3 are not independent. Hence, we must use Theorem 5.6 to find the PMF of W. In particular, since $X_1 + X_2 + X_3 = 5$ and since each X_i is non-negative, $P_W(0) = P_W(1) = 0$. Furthermore,

$$\begin{aligned} P_W(2) &= P_{\mathbf{X}}(1,2,2) + P_{\mathbf{X}}(2,1,2) + P_{\mathbf{X}}(2,2,1) \\ &= \frac{5![0.3(0.6)^2(0.1)^2 + 0.3^2(0.6)(0.1)^2 + 0.3^2(0.6)^2(0.1)]}{2!2!1!} \\ &= 0.1458 \end{aligned}$$

In addition, for $w = 3$, $w = 4$, and $w = 5$, the event $W = w$ occurs if and only if one of the mutually exclusive events $X_1 = w$, $X_2 = w$, or $X_3 = w$ occurs. Thus,

$$\begin{aligned} P_W(3) &= P_{X_1}(3) + P_{X_2}(3) + P_{X_3}(3) = 0.486 \\ P_W(4) &= P_{X_1}(4) + P_{X_2}(4) + P_{X_3}(4) = 0.288 \\ P_W(5) &= P_{X_1}(5) + P_{X_2}(5) + P_{X_3}(5) = 0.0802 \end{aligned}$$

Quiz 5.5(B)

The random vector $\mathbf{X}$ has PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} e^{-x_3} & 0 \le x_1 \le x_2 \le x_3, \\ 0 & \text{otherwise.} \end{cases}$$

Find the PDF of $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$. where $\mathbf{A} = \text{diag}[2, 2, 2]$ and $\mathbf{b} = \begin{bmatrix} 4 & 4 & 4 \end{bmatrix}'$.

Quiz 5.5(B) Solution

Since each $Y_i = 2X_i + 4$, we can apply Theorem 5.10 to write

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{1}{2^3} f_{\mathbf{X}}\left(\frac{y_1 - 4}{2}, \frac{y_2 - 4}{2}, \frac{y_3 - 4}{2}\right)$$
$$= \begin{cases} (1/8)e^{-(y_3-4)/2} & 4 \leq y_1 \leq y_2 \leq y_3 \\ 0 & \text{otherwise} \end{cases}$$

Note that for other matrices $\mathbf{A}$, the constraints on $\mathbf{y}$ resulting from the constraints $0 \leq X_1 \leq X_2 \leq X_3$ can be much more complicated.

5.6 Expected Value Vector and Correlation Matrix

- Corresponding the expected value of a single random variable, the expected value of a random vector is a column vector in which the components are the expected values of the components of the random vector. There is a corresponding definition of the variance and standard deviation of a random vector.

Definition 5.11 *Expected Value Vector*

The expected value of a random vector $\mathbf{X}$ *is a column vector*

$$E[\mathbf{X}] = \boldsymbol{\mu}_{\mathbf{X}} = \begin{bmatrix} E[X_1] & E[X_2] & \cdots & E[X_n] \end{bmatrix}'.$$

- The correlation and covariance (Definition 4.5 and Definition 4.4) are numbers that contain important information about a pair of random variables. Corresponding information about random vectors is reflected in the set of correlations and the set of covariances of all pairs components. These sets are referenced to as **second order statistics**.

- They have a concise matrix notation. To establish the notation, we first observe that for random vectors $\mathbf{X}$ with n components and $\mathbf{Y}$ with m components, the set of all products, $X_i Y_j$, is contained in the $n \times m$ random matrix $\mathbf{XY}'$. If $\mathbf{Y} = \mathbf{X}$, the random matrix $\mathbf{XX}'$ contains all products, $X_i X_j$, of components $\mathbf{X}$.

Example 5.9 Problem

If $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$, what are the components of $\mathbf{X}\mathbf{X}'$?

Example 5.9 Solution

$$\mathbf{X}\mathbf{X}' = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix} = \begin{bmatrix} X_1^2 & X_1X_2 & X_1X_3 \\ X_2X_1 & X_2^2 & X_2X_3 \\ X_3X_1 & X_3X_2 & X_3^2 \end{bmatrix}.$$

Definition 5.12 Matrix

For a random matrix $\mathbf{A}$ *with the random variable* A_{ij} *as its* i, j *th element,* $E[\mathbf{A}]$ *is a matrix with* i, j *th element* $E[A_{ij}]$.

- In definition 5.11, we defined the expected value of a random vector as the vector of expected values, Definition 5.11 can be extended to random matrices (Definition 5.12).
- Applying definition 5.12 to random matrix $\mathbf{X}\mathbf{X}'$, we have a concise way to define the correlation matrix of random vector $\mathbf{X}$.

Definition 5.13 *Vector Correlation*

The correlation of a random vector $\mathbf{X}$ *is an* $n \times n$ *matrix* $\mathbf{R_X}$ *with* i, j*th element* $R_X(i, j) = E[X_i X_j]$. *In vector noation,*

$$\mathbf{R_X} = E\left[\mathbf{X}\mathbf{X}'\right].$$

Example 5.10

If $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$, the correlation matrix of $\mathbf{X}$ is

$$\mathbf{R_X} = \begin{bmatrix} E\left[X_1^2\right] & E\left[X_1 X_2\right] & E\left[X_1 X_3\right] \\ E\left[X_2 X_1\right] & E\left[X_2^2\right] & E\left[X_2 X_3\right] \\ E\left[X_3 X_1\right] & E\left[X_3 X_2\right] & E\left[X_3^2\right] \end{bmatrix} = \begin{bmatrix} E\left[X_1^2\right] & r_{X_1,X_2} & r_{X_1,X_3} \\ r_{X_2,X_1} & E\left[X_2^2\right] & r_{X_2,X_3} \\ r_{X_3,X_1} & r_{X_3,X_2} & E\left[X_3^2\right] \end{bmatrix}.$$

Definition 5.14 *Vector Covariance*

The covariance of a random vector $\mathbf{X}$ *is an* $n \times n$ *matrix* $\mathbf{C_X}$ *with components* $C_X(i, j) = \mathrm{Cov}[X_i, X_j]$. *In vector notation,*

$$\mathbf{C_X} = E\left[(\mathbf{X} - \boldsymbol{\mu}_\mathbf{X})(\mathbf{X} - \boldsymbol{\mu}_\mathbf{X})'\right]$$

- The i, jth element of the correlation matrix is the expected value of the random variable $X_i X_j$. The covariance matrix of $\mathbf{X}$ is a similar generalization of the covariance of two random variables.

Example 5.11

If $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$, the covariance matrix of $\mathbf{X}$ is

$$\mathbf{C_X} = \begin{bmatrix} \text{Var}[X_1] & \text{Cov}\,[X_1, X_2] & \text{Cov}\left[X_1, X_3\right] \\ \text{Cov}\,[X_2, X_1] & \text{Var}[X_2] & \text{Cov}\left[X_2, X_3\right] \\ \text{Cov}\left[X_3, X_1\right] & \text{Cov}\left[X_3, X_2\right] & \text{Var}[X_3] \end{bmatrix}$$

Theorem 5.12

For a random vector $\mathbf{X}$ with correlation matrix $\mathbf{R_X}$, covariance matrix $\mathbf{C_X}$, and vector expected value $\boldsymbol{\mu}_\mathbf{X}$,

$$\mathbf{C_X} = \mathbf{R_X} - \boldsymbol{\mu}_\mathbf{X}\boldsymbol{\mu}'_\mathbf{X}.$$

- Theorem 4.16(a), which connects the correlation and covariance of a pair of random variables, can be extended to random vectors.

$$\mathrm{Cov}[X,Y] = r_{X,Y} - \mu_X \mu_Y$$

Proof: Theorem 5.12

The proof is essentially the same as the proof of Theorem 4.16(a) with vectors replacing scalars. Cross multiplying inside the expectation of Definition 5.14 yields

$$\begin{aligned}\mathbf{C}_{\mathbf{X}} &= E\left[\mathbf{X}\mathbf{X}' - \mathbf{X}\boldsymbol{\mu}_{\mathbf{X}}' - \boldsymbol{\mu}_{\mathbf{X}}\mathbf{X}' + \boldsymbol{\mu}_{\mathbf{X}}\boldsymbol{\mu}_{\mathbf{X}}'\right] \\ &= E\left[\mathbf{X}\mathbf{X}'\right] - E\left[\mathbf{X}\boldsymbol{\mu}_{\mathbf{X}}'\right] - E\left[\boldsymbol{\mu}_{\mathbf{X}}\mathbf{X}'\right] + E\left[\boldsymbol{\mu}_{\mathbf{X}}\boldsymbol{\mu}_{\mathbf{X}}'\right].\end{aligned}$$

Since $E[\mathbf{X}] = \boldsymbol{\mu}_{\mathbf{X}}$ is a constant vector,

$$\mathbf{C}_{\mathbf{X}} = \mathbf{R}_{\mathbf{X}} - E\left[\mathbf{X}\right]\boldsymbol{\mu}_{\mathbf{X}}' - \boldsymbol{\mu}_{\mathbf{X}}E\left[\mathbf{X}'\right] + \boldsymbol{\mu}_{\mathbf{X}}\boldsymbol{\mu}_{\mathbf{X}}' = \mathbf{R}_{\mathbf{X}} - \boldsymbol{\mu}_{\mathbf{X}}\boldsymbol{\mu}_{\mathbf{X}}'.$$

Example 5.12 Problem

Find the expected value $E[\mathbf{X}]$, the correlation matrix $\mathbf{R_X}$, and the covariance matrix $\mathbf{C_X}$ of the 2-dimensional random vector $\mathbf{X}$ with PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} 2 & 0 \leq x_1 \leq x_2 \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Example 5.12 Solution

The elements of the expected value vector are

$$E\left[X_i\right] = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x_i f_{\mathbf{X}}\left(\mathbf{x}\right)\, dx_1 dx_2 = \int_0^1\int_0^{x_2} 2x_i\, dx_1 dx_2, \quad i = 1, 2.$$

The integrals are $E[X_1] = 1/3$ and $E[X_2] = 2/3$, so that $\boldsymbol{\mu}_{\mathbf{X}} = E[\mathbf{X}] = \begin{bmatrix}1/3 & 2/3\end{bmatrix}'$. The elements of the correlation matrix are

$$E\left[X_1^2\right] = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x_1^2 f_{\mathbf{X}}\left(\mathbf{x}\right)\, dx_1 dx_2 = \int_0^1\int_0^{x_2} 2x_1^2\, dx_1 dx_2,$$

$$E\left[X_2^2\right] = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x_2^2 f_{\mathbf{X}}\left(\mathbf{x}\right)\, dx_1 dx_2 = \int_0^1\int_0^{x_2} 2x_2^2\, dx_1 dx_2,$$

$$E\left[X_1 X_2\right] = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x_1 x_2 f_{\mathbf{X}}\left(\mathbf{x}\right)\, dx_1 dx_2 = \int_0^1\int_0^{x_2} 2x_1 x_2\, dx_1 dx_2.$$

[Continued]

Example 5.12 Solution (continued)

These integrals are $E[X_1{}^2] = 1/6$, $E[X_2{}^2] = 1/2$, and $E[X_1X_2] = 1/4$.

Therefore,

$$\mathbf{R_X} = \begin{bmatrix} 1/6 & 1/4 \\ 1/4 & 1/2 \end{bmatrix}.$$

We use Theorem 5.12 to find the elements of the covariance matrix.

$$\mathbf{C_X} = \mathbf{R_X} - \boldsymbol{\mu}_\mathbf{X}\boldsymbol{\mu}'_\mathbf{X} = \begin{bmatrix} 1/6 & 1/4 \\ 1/4 & 1/2 \end{bmatrix} - \begin{bmatrix} 1/9 & 2/9 \\ 2/9 & 4/9 \end{bmatrix} = \begin{bmatrix} 1/18 & 1/36 \\ 1/36 & 1/18 \end{bmatrix}.$$

- In addition to the correlations and covariances of the elements of one random vector, it is useful to refer to the correlations and covariances of elements of two random vectors.

Definition 5.15 *Vector Cross-Correlation*

The cross-correlation of random vectors, $\mathbf{X}$ *with* n *components and* $\mathbf{Y}$ *with* m *components, is an* $n \times m$ *matrix* $\mathbf{R_{XY}}$ *with* i, j*th element* $R_{XY}(i, j) = E[X_i Y_j]$, *or, in vector notation,*

$$\mathbf{R_{XY}} = E\left[\mathbf{X}\mathbf{Y}'\right].$$

Definition 5.16 *Vector Cross-Covariance*

The cross-covariance of a pair of random vectors $\mathbf{X}$ *with* n *components and* $\mathbf{Y}$ *with* m *components is an* $n \times m$ *matrix* $\mathbf{C}_{\mathbf{XY}}$ *with* i, j *th element* $C_{\mathbf{XY}}(i, j) = \mathrm{Cov}[X_i, Y_j]$*, or, in vector notation,*

$$\mathbf{C}_{\mathbf{XY}} = E\left[(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})(\mathbf{Y} - \boldsymbol{\mu}_{\mathbf{Y}})'\right].$$

- To distinguish the correlation or covariance of a random vector from the correlation or covariance of a pair of random vectors, we sometimes use the terminology **autocorrelation** and **autocovariance** then there is one random vector and **cross-correlation** and **cross-covariance** when there is a pair of random vectors.
- Note that when $\mathbf{X} = \mathbf{Y}$ the autocorrelation and cross-correlation are identical (as are the covariances). Recognizing this identity, some texts use the notation $\mathbf{R}_{\mathbf{XX}}$ and $\mathbf{C}_{\mathbf{XX}}$ for the correlation and covariance of a random vector.
- When $\mathbf{Y}$ is a linear transformation of $\mathbf{X}$, the following theorem states the relationship of the second-order statistics of $\mathbf{Y}$ to the corresponding statistics of $\mathbf{X}$.

Theorem 5.13

$\mathbf{X}$ is an n-dimensional random vector with expected value $\boldsymbol{\mu}_{\mathbf{X}}$, correlation $\mathbf{R}_{\mathbf{X}}$, and covariance $\mathbf{C}_{\mathbf{X}}$. The m-dimensional random vector $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$, where $\mathbf{A}$ is an $m \times n$ matrix and $\mathbf{b}$ is an m-dimensional vector, has expected value $\boldsymbol{\mu}_{\mathbf{Y}}$, correlation matrix $\mathbf{R}_{\mathbf{Y}}$, and covariance matrix $\mathbf{C}_{\mathbf{Y}}$ given by

$$\begin{aligned}
\boldsymbol{\mu}_{\mathbf{Y}} &= \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}} + \mathbf{b}, \\
\mathbf{R}_{\mathbf{Y}} &= \mathbf{A}\mathbf{R}_{\mathbf{X}}\mathbf{A}' + (\mathbf{A}\boldsymbol{\mu}_{\mathbf{X}})\mathbf{b}' + \mathbf{b}(\mathbf{A}\boldsymbol{\mu}_{\mathbf{X}})' + \mathbf{b}\mathbf{b}', \\
\mathbf{C}_{\mathbf{Y}} &= \mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'.
\end{aligned}$$

Proof: Theorem 5.13

We derive the formulas for the expected value and covariance of $\mathbf{Y}$. The derivation for the correlation is similar. First, the expected value of $\mathbf{Y}$ is

$$\boldsymbol{\mu}_{\mathbf{Y}} = E\left[\mathbf{AX} + \mathbf{b}\right] = \mathbf{A}E\left[\mathbf{X}\right] + E\left[\mathbf{b}\right] = \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}} + \mathbf{b}.$$

It follows that $\mathbf{Y} - \boldsymbol{\mu}_{\mathbf{Y}} = \mathbf{A}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})$. This implies

$$\begin{aligned}\mathbf{C}_{\mathbf{Y}} &= E\left[(\mathbf{A}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}}))(\mathbf{A}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}}))'\right] \\ &= E\left[\mathbf{A}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}}))(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})'\mathbf{A}'\right] = \mathbf{A}E\left[(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})'\right]\mathbf{A}' = \mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'.\end{aligned}$$

$$\begin{aligned}\mathbf{R}_{\mathbf{Y}} &= E\left[\left(\mathbf{AX}+\mathbf{b}\right)\left(\mathbf{AX}+\mathbf{b}\right)'\right] = E\left[\left(\mathbf{AX}+\mathbf{b}\right)\left(\mathbf{X}'\mathbf{A}'+\mathbf{b}'\right)\right] \\ &= E\left[\mathbf{AXX}'\mathbf{A}' + \mathbf{AXb}' + \mathbf{bX}'\mathbf{A}' + \mathbf{bb}'\right] \\ &= \mathbf{A}E\left[\mathbf{XX}'\right]\mathbf{A}' + \mathbf{A}E\left[\mathbf{X}\right]\mathbf{b}' + \mathbf{b}E\left[\mathbf{X}'\right]\mathbf{A}' + \mathbf{bb}' \\ &= \mathbf{AR}_{\mathbf{X}}\mathbf{A}' + \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}}\mathbf{b}' + \mathbf{b}\boldsymbol{\mu}_{\mathbf{X}}'\mathbf{A}' + \mathbf{bb}'\end{aligned}$$

Example 5.13 Problem

Given random vector $\mathbf{X}$ defined in Example 5.12, let $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 6 & 3 \\ 3 & 6 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 0 \\ -2 \\ -2 \end{bmatrix}.$$

Find the expected value $\boldsymbol{\mu}_{\mathbf{Y}}$, the correlation $\mathbf{R}_{\mathbf{Y}}$, and the covariance $\mathbf{C}_{\mathbf{Y}}$.

Example 5.13 Solution

From the matrix operations of Theorem 5.13, we obtain $\boldsymbol{\mu}_{\mathbf{Y}} = \begin{bmatrix} 1/3 & 2 & 3 \end{bmatrix}'$ and

$$\mathbf{R}_{\mathbf{Y}} = \begin{bmatrix} 1/6 & 13/12 & 4/3 \\ 13/12 & 7.5 & 9.25 \\ 4/3 & 9.25 & 12.5 \end{bmatrix}; \qquad \mathbf{C}_{\mathbf{Y}} = \begin{bmatrix} 1/18 & 5/12 & 1/3 \\ 5/12 & 3.5 & 3.25 \\ 1/3 & 3.25 & 3.5 \end{bmatrix}.$$

Theorem 5.14

The vectors $\mathbf{X}$ and $\mathbf{Y} = \mathbf{AX} + \mathbf{b}$ have cross-correlation $\mathbf{R_{XY}}$ and cross-covariance $\mathbf{C_{XY}}$ given by

$$\mathbf{R_{XY}} = \mathbf{R_X A}' + \boldsymbol{\mu}_\mathbf{X}\mathbf{b}', \qquad \mathbf{C_{XY}} = \mathbf{C_X A}'.$$

- The cross-correlation and cross-covariance of two random vectors can be derived using algebra similar to the proof of Theorem 5.13.

Example 5.14 Problem

Continuing Example 5.13 for random vectors $\mathbf{X}$ and $\mathbf{Y} = \mathbf{AX} + \mathbf{b}$, calculate

(a) The cross-correlation matrix $\mathbf{R_{XY}}$ and the cross-covariance matrix $\mathbf{C_{XY}}$.

(b) The correlation coefficients ρ_{Y_1,Y_3} and ρ_{X_2,Y_1}.

Example 5.14 Solution

(a) Direct matrix calculation using Theorem 5.14 yields

$$\mathbf{R_{XY}} = \begin{bmatrix} 1/6 & 13/12 & 4/3 \\ 1/4 & 5/3 & 29/12 \end{bmatrix}; \qquad \mathbf{C_{XY}} = \begin{bmatrix} 1/18 & 5/12 & 1/3 \\ 1/36 & 1/3 & 5/12 \end{bmatrix}.$$

(b) Referring to Definition 4.8 and recognizing that $\mathrm{Var}[Y_i] = C_{\mathbf{Y}}(i, i)$, we have

$$\rho_{Y_1,Y_3} = \frac{\mathrm{Cov}\left[Y_1, Y_3\right]}{\sqrt{\mathrm{Var}[Y_1]\,\mathrm{Var}[Y_3]}} = \frac{C_Y(1,3)}{\sqrt{C_Y(1,1)C_Y(3,3)}} = 0.756$$

Similarly,

$$\rho_{X_2,Y_1} = \frac{\mathrm{Cov}\left[X_2, Y_1\right]}{\sqrt{\mathrm{Var}[X_2]\,\mathrm{Var}[Y_1]}} = \frac{C_{\mathbf{XY}}(2,1)}{\sqrt{C_{\mathbf{X}}(2,2)C_{\mathbf{Y}}(1,1)}} = 1/2.$$

Quiz 5.6

In Quiz 5.3, the 3-dimensional random vector $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$ has PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} 6 & 0 \leq x_1 \leq x_2 \leq x_3 \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find the expected value $E[\mathbf{X}]$, and the correlation and covariance matrices $\mathbf{R_X}$ and $\mathbf{C_X}$.

Quiz 5.6 Solution

We start by finding the components $E[X_i] = \int_{-\infty}^{\infty} x f_{X_i}(x)\, dx$ of $\boldsymbol{\mu}_X$. To do so, we use the marginal PDFs $f_{X_i}(x)$ found in Quiz 5.3:

$$E[X_1] = \int_0^1 3x(1-x)^2\, dx = 1/4,$$

$$E[X_2] = \int_0^1 6x^2(1-x)\, dx = 1/2,$$

$$E[X_3] = \int_0^1 3x^3\, dx = 3/4.$$

To find the correlation matrix $\mathbf{R}_X$, we need to find $E[X_i X_j]$ for all i and j. We start with the second moments:

$$E\left[X_1^2\right] = \int_0^1 3x^2(1-x)^2\, dx = 1/10.$$

$$E\left[X_2^2\right] = \int_0^1 6x^3(1-x)\, dx = 3/10.$$

$$E\left[X_3^2\right] = \int_0^1 3x^4\, dx = 3/5.$$

[Continued]

Quiz 5.6 Solution (continued)

Using marginal PDFs from Quiz 5.3, the cross terms are

$$E[X_1X_2] = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} x_1x_2 f_{X_1,X_2}(x_1, x_2)\,,\ dx_1\,dx_2$$

$$= \int_0^1 \left(\int_{x_1}^1 6x_1x_2(1-x_2)\,dx_2\right) dx_1 = \int_0^1 [x_1 - 3x_1^3 + 2x_1^4]\,dx_1 = 3/20.$$

$$E[X_2X_3] = \int_0^1\int_{x_2}^1 6x_2^2x_3\,dx_3\,dx_2 = \int_0^1 [3x_2^2 - 3x_2^4]\,dx_2 = 2/5$$

$$E[X_1X_3] = \int_0^1\int_{x_1}^1 6x_1x_3(x_3 - x_1)\,dx_3\,dx_1.$$

$$= \int_0^1 \left((2x_1x_3^3 - 3x_1^2x_3^2)\Big|_{x_3=x_1}^{x_3=1}\right) dx_1 = \int_0^1 [2x_1 - 3x_1^2 + x_1^4]\,dx_1 = 1/5.$$

Summarizing the results, $\mathbf{X}$ has correlation matrix

$$\mathbf{R}_X = \begin{bmatrix} 1/10 & 3/20 & 1/5 \\ 3/20 & 3/10 & 2/5 \\ 1/5 & 2/5 & 3/5 \end{bmatrix}.$$

[Continued]

Quiz 5.6 Solution (continued)

Vector $\mathbf{X}$ has covariance matrix

$$\begin{aligned}
\mathbf{C}_X &= \mathbf{R}_X - E[\mathbf{X}]\,E[\mathbf{X}]' \\
&= \begin{bmatrix} 1/10 & 3/20 & 1/5 \\ 3/20 & 3/10 & 2/5 \\ 1/5 & 2/5 & 3/5 \end{bmatrix} - \begin{bmatrix} 1/4 \\ 1/2 \\ 3/4 \end{bmatrix} \begin{bmatrix} 1/4 & 1/2 & 3/4 \end{bmatrix} \\
&= \begin{bmatrix} 1/10 & 3/20 & 1/5 \\ 3/20 & 3/10 & 2/5 \\ 1/5 & 2/5 & 3/5 \end{bmatrix} - \begin{bmatrix} 1/16 & 1/8 & 3/16 \\ 1/8 & 1/4 & 3/8 \\ 3/16 & 3/8 & 9/16 \end{bmatrix} = \frac{1}{80} \begin{bmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 3 \end{bmatrix}.
\end{aligned}$$

This problem shows that even for fairly simple joint PDFs, computing the covariance matrix by calculus can be a time consuming task.

5.7 Gaussian Random Vectors

- Multiple Gaussian random variables appear in many practical applications of probability theory. The **multivariate Gaussian distribution** is a probability model to n random variables with the property that the marginal PDFs are all Gaussian. A set of random variables described by the multivariate Gaussian PDF is said to be **jointly Gaussian**. A vector whose components are jointly Gaussian random variables is said to be a **Gaussian random vector**. The PDF of a Gaussian random vector has a particularly concise notation. The following definition is a generalization of Definition 3.9 and Definition 4.17.

Definition 5.17 Gaussian Random Vector

$\mathbf{X}$ *is the Gaussian* $(\boldsymbol{\mu}_{\mathbf{X}}, \mathbf{C}_{\mathbf{X}})$ *random vector with expected value* $\boldsymbol{\mu}_{\mathbf{X}}$ *and covariance* $\mathbf{C}_{\mathbf{X}}$ *if and only if*

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{n/2}[det(\mathbf{C}_{\mathbf{X}})]^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})'\mathbf{C}_{\mathbf{X}}^{-1}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})\right)$$

where det$(\mathbf{C}_{\mathbf{X}})$*, the determinant of* $\mathbf{C}_{\mathbf{X}}$*, satisfies det*$(\mathbf{C}_{\mathbf{X}}) > 0$*.*

- When $n = 1$, $\mathbf{C_X} = \sigma_X^2$ and $\mathbf{x} - \boldsymbol{\mu}_\mathbf{X} = x - \mu_X$, and the PDF in Definition 5.17 reduces to the ordinary Gaussian PDF of Definition 3.8. That is, a 1-dimensional Gaussian (μ, σ) random vector is a Gaussian (μ, σ) rand variable, notwithstanding that we write their parameters differently.
- In Problem 5.7.4, we ask you to show that for $n = 2$, Definition 5.17 reduces to the bivariate Gaussian PDF in Definition 4.17. The condition that $\det(\mathbf{C_X}) > 0$ is a generalization of the requirement for the bivariate Gaussian PDF that $|\rho| < 1$. Basically, $\det(\mathbf{C_X}) > 0$ reflects the requirement that no random variable X_i is a linear combination of the other random variables X_j.

- For a Gaussian random vector $\mathbf{X}$, an important special case arises when $\mathrm{Cov}[X_i, X_j] = 0$ for all $i \neq j$. In this case, the off-diagonal elements of the covariance matrix $\mathbf{C_X}$ are all zero and the ith diagomal element is simply $\mathrm{Var}[X_i] = \sigma_i^2$. In this case, we write $\mathbf{C_X} = \mathrm{diag}[\sigma_1^2, \sigma_2^2, \ldots, \sigma_n^2]$. When the covariance matrix is diagonal, X_i and X_j are uncorrelated for $i \neq j$. In Theorem 4.32, we showed that uncorrelated bivariate Gaussian random variables are independent. The following theorem generalized this result.

Theorem 5.15

A Gaussian random vector $\mathbf{X}$ has independent components if and only if $\mathbf{C}_{\mathbf{X}}$ is a diagonal matrix.

Proof: Theorem 5.15

First, if the components of $\mathbf{X}$ are independent, then for $i \neq j$, X_i and X_j are independent. By Theorem 4.27(c), $\mathrm{Cov}[X_i, X_j] = 0$. Hence the off-diagonal terms of $\mathbf{C_X}$ are all zero. If $\mathbf{C_X}$ is diagonal, then

$$\mathbf{C_X} = \begin{bmatrix} \sigma_1^2 & & \\ & \ddots & \\ & & \sigma_n^2 \end{bmatrix} \quad \text{and} \quad \mathbf{C_X}^{-1} = \begin{bmatrix} 1/\sigma_1^2 & & \\ & \ddots & \\ & & 1/\sigma_n^2 \end{bmatrix}.$$

It follows that $\mathbf{C_X}$ has determinant $\det(\mathbf{C_X}) = \prod_{i=1}^n \sigma_i^2$ and that

$$(\mathbf{x} - \boldsymbol{\mu}_\mathbf{X})'\mathbf{C_X}^{-1}(\mathbf{x} - \boldsymbol{\mu}_\mathbf{X}) = \sum_{i=1}^n \frac{(X_i - \mu_i)^2}{\sigma_i^2}.$$

From Definition 5.17, we see that

$$\begin{aligned} f_\mathbf{X}(\mathbf{x}) &= \frac{1}{(2\pi)^{n/2}\prod_{i=1}^n \sigma_i^2} \exp\left(-\sum_{i=1}^n (x_i - \mu_i)/2\sigma_i^2\right) \\ &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(-(x_i - \mu_i)^2/2\sigma_i^2\right). \end{aligned}$$

Thus $f_\mathbf{X}(\mathbf{x}) = \prod_{i=1}^n f_{X_i}(x_i)$, implying $X_1, \ldots, X_n$ are independent.

Example 5.15 Problem

Consider the outdoor temperature at a certain weather station. On May 5, the temperature measurements in units of degrees Fahrenheit taken at 6 AM, 12 noon, and 6 PM are all Gaussian random variables, X_1, X_2, X_3 with variance 16 degrees^2. The expected values are 50 degrees, 62 degrees, and 58 degrees respectively. The covariance matrix of the three measurements is

$$\mathbf{C_X} = \begin{bmatrix} 16.0 & 12.8 & 11.2 \\ 12.8 & 16.0 & 12.8 \\ 11.2 & 12.8 & 16.0 \end{bmatrix}.$$

(a) Write the joint PDF of X_1, X_2 using the algebraic notation of Definition 4.17.

(b) Write the joint PDF of X_1, X_2 using vector notation.

(c) Write the joint PDF of $\mathbf{X} = \begin{bmatrix} X_1 & X_2 & X_3 \end{bmatrix}'$ using vector notation.

Example 5.15 Solution

(a) First we note that X_1 and X_2 have expected values $\mu_1 = 50$ and $\mu_2 = 62$, variances $\sigma_1^2 = \sigma_2^2 = 16$, and covariance $\mathrm{Cov}[X_1, X_2] = 12.8$. It follows from Definition 4.8 that the correlation coefficient is

$$\rho_{X_1,X_2} = \frac{\mathrm{Cov}\left[X_1, X_2\right]}{\sigma_1\sigma_2} = \frac{12.8}{16} = 0.8.$$

From Definition 4.17, the joint PDF is

$$f_{X_1,X_2}(x_1, x_2) = \frac{\exp\left(-\frac{(x_1-50)^2-1.6(x_1-50)(x_2-62)+(x_2-62)^2}{19.2}\right)}{60.3}.$$

(b) Let $\mathbf{W} = \begin{bmatrix} X_1 & X_2 \end{bmatrix}'$ denote a vector representation for random variables X_1 and X_2. From the covariance matrix $\mathbf{C_X}$, we observe that the 2×2 submatrix in the upper left corner is the covariance matrix of the random vector $\mathbf{W}$. Thus

[Continued]

Example 5.15 Solution (continued)

$$\boldsymbol{\mu}_{\mathbf{W}} = \begin{bmatrix} 50 \\ 62 \end{bmatrix}, \qquad \mathbf{C}_{\mathbf{W}} = \begin{bmatrix} 16.0 & 12.8 \\ 12.8 & 16.0 \end{bmatrix}.$$

We observe that $\det(\mathbf{C}_{\mathbf{W}}) = 92.16$ and $\det(\mathbf{C}_{\mathbf{W}})^{1/2} = 9.6$. From Definition 5.17, the joint PDF of $\mathbf{W}$ is

$$f_{\mathbf{W}}(\mathbf{w}) = \frac{1}{60.3} \exp\left(-\frac{1}{2}(\mathbf{w} - \boldsymbol{\mu}_{\mathbf{W}})^T \mathbf{C}_{\mathbf{W}}^{-1}(\mathbf{w} - \boldsymbol{\mu}_{\mathbf{W}})\right).$$

(c) For the joint PDF of $\mathbf{X}$, we note that $\mathbf{X}$ has expected value $\boldsymbol{\mu}_{\mathbf{X}} = \begin{bmatrix} 50 & 62 & 58 \end{bmatrix}'$ and that $\det(\mathbf{C}_{\mathbf{X}})^{1/2} = 22.717$. Thus

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{357.8} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})^T \mathbf{C}_{\mathbf{X}}^{-1}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})\right).$$

Theorem 5.16

Given an n-dimensional Gaussian random vector $\mathbf{X}$ with expected value $\boldsymbol{\mu}_{\mathbf{X}}$ and covariance $\mathbf{C}_{\mathbf{X}}$, and an $m \times n$ matrix $\mathbf{A}$ with rank$(\mathbf{A}) = m$,

$$\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$$

is an m-dimensional Gaussian random vector with expected value $\boldsymbol{\mu}_{\mathbf{Y}} = \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}} + \mathbf{b}$ and covariance $\mathbf{C}_{\mathbf{Y}} = \mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'$.

- This theorem is a generalization of Theorem 3.13. It states that a linear transformation of a Gaussian random vector results in another Gaussian random vector.

Proof: Theorem 5.16

The proof of Theorem 5.13 contains the derivations of $\boldsymbol{\mu}_{\mathbf{Y}}$ and $\mathbf{C}_{\mathbf{Y}}$. Our proof that $\mathbf{Y}$ has a Gaussian PDF is confined to the special case when $m = n$ and $\mathbf{A}$ is an invertible matrix. The case of $m < n$ is addressed in Problem 5.7.9. When $m = n$, we use Theorem 5.11 to write

$$\begin{aligned} f_{\mathbf{Y}}(\mathbf{y}) &= \frac{1}{|\det(\mathbf{A})|} f_{\mathbf{X}}\left(\mathbf{A}^{-1}(\mathbf{y}-\mathbf{b})\right) \\ &= \frac{\exp\left(-\frac{1}{2}[\mathbf{A}^{-1}(\mathbf{y}-\mathbf{b})-\boldsymbol{\mu}_{\mathbf{X}}]'\mathbf{C}_{\mathbf{X}}^{-1}[\mathbf{A}^{-1}(\mathbf{y}-\mathbf{b})-\boldsymbol{\mu}_{\mathbf{X}}]\right)}{(2\pi)^{n/2}\,|\det(\mathbf{A})|\,|\det(\mathbf{C}_{\mathbf{X}})|^{1/2}}. \end{aligned}$$

In the exponent of $f_{\mathbf{Y}}(\mathbf{y})$, we observe that

$$\mathbf{A}^{-1}(\mathbf{y}-\mathbf{b})-\boldsymbol{\mu}_{\mathbf{X}} = \mathbf{A}^{-1}[\mathbf{y}-(\mathbf{A}\boldsymbol{\mu}_{\mathbf{X}}+\mathbf{b})] = \mathbf{A}^{-1}(\mathbf{y}-\boldsymbol{\mu}_{\mathbf{Y}}),$$

since $\boldsymbol{\mu}_{\mathbf{Y}} = \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}} + \mathbf{b}$.

[Continued]

Proof: Theorem 5.16 (continued)

Applying (5.79) to (5.78) yields

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{\exp\left(-\frac{1}{2}[\mathbf{A}^{-1}(\mathbf{y}-\boldsymbol{\mu}_{\mathbf{Y}})]'\mathbf{C}_{\mathbf{X}}^{-1}[\mathbf{A}^{-1}(\mathbf{y}-\boldsymbol{\mu}_{\mathbf{Y}})]\right)}{(2\pi)^{n/2}\,|\det(\mathbf{A})|\,|\det(\mathbf{C}_{\mathbf{X}})|^{1/2}}.$$

Using the identities $|\det(\mathbf{A})||\det(\mathbf{C}_{\mathbf{X}})|^{1/2} = |\det(\mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}')|^{1/2}$ and $(\mathbf{A}^{-1})' = (\mathbf{A}')^{-1}$, we can write

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{\exp\left(-\frac{1}{2}(\mathbf{y}-\boldsymbol{\mu}_{\mathbf{Y}})'(\mathbf{A}')^{-1}\mathbf{C}_{\mathbf{X}}^{-1}\mathbf{A}^{-1}(\mathbf{y}-\boldsymbol{\mu}_{\mathbf{Y}})\right)}{(2\pi)^{n/2}\left|\det\left(\mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'\right)\right|^{1/2}}.$$

Since $(\mathbf{A}')^{-1}\mathbf{C}_{\mathbf{X}}^{-1}\mathbf{A}^{-1} = (\mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}')^{-1}$, we see from Equation (5.81) that $\mathbf{Y}$ is a Gaussian vector with expected value $\boldsymbol{\mu}_{\mathbf{Y}}$ and covariance matrix $\mathbf{C}_{\mathbf{Y}} = \mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'$.

Example 5.16 Problem

Continuing Example 5.15, use the formula $Y_i = (5/9)(X_i - 32)$ to convert the three temperature measurements to degrees Celsius.

(a) What is $\boldsymbol{\mu}_{\mathbf{Y}}$, the expected value of random vector $\mathbf{Y}$?

(b) What is $\mathbf{C}_{\mathbf{Y}}$, the covariance of random vector $\mathbf{Y}$?

(c) Write the joint PDF of $\mathbf{Y} = \begin{bmatrix} Y_1 & Y_2 & Y_3 \end{bmatrix}'$ using vector notation.

Example 5.16 Solution

(a) In terms of matrices, we observe that $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$ where

$$\mathbf{A} = \begin{bmatrix} 5/9 & 0 & 0 \\ 0 & 5/9 & 0 \\ 0 & 0 & 5/9 \end{bmatrix}, \qquad \mathbf{b} = -\frac{160}{9}\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

(b) Since $\boldsymbol{\mu}_{\mathbf{X}} = \begin{bmatrix} 50 & 62 & 58 \end{bmatrix}'$, from Theorem 5.16,

$$\boldsymbol{\mu}_{\mathbf{Y}} = \mathbf{A}\boldsymbol{\mu}_{\mathbf{X}} + \mathbf{b} = \begin{bmatrix} 10 \\ 50/3 \\ 130/9 \end{bmatrix}.$$

(c) The covariance of $\mathbf{Y}$ is $\mathbf{C}_{\mathbf{Y}} = \mathbf{A}\mathbf{C}_{\mathbf{X}}\mathbf{A}'$. We note that $\mathbf{A} = \mathbf{A}' = (5/9)\mathbf{I}$ where $\mathbf{I}$ is the 3×3 identity matrix. Thus $\mathbf{C}_{\mathbf{Y}} = (5/9)^2\mathbf{C}_{\mathbf{X}}$ and $\mathbf{C}_{\mathbf{Y}}^{-1} = (9/5)^2\mathbf{C}_{\mathbf{X}}^{-1}$. The PDF of $\mathbf{Y}$ is

$$f_{\mathbf{Y}}(\mathbf{y}) = \frac{1}{24.47}\exp\left(-\frac{81}{50}(\mathbf{y} - \boldsymbol{\mu}_{\mathbf{Y}})^T\mathbf{C}_{\mathbf{X}}^{-1}(\mathbf{y} - \boldsymbol{\mu}_{\mathbf{Y}})\right).$$

Standard Normal Random

Definition 5.18 Vector

The n-dimensional standard normal random vector $\mathbf{Z}$ *is the n-dimensional Gaussian random vector with $E[\mathbf{Z}] = \mathbf{0}$ and $\mathbf{C_Z} = \mathbf{I}$.*

- A standard normal random vector is a generalization of the standard norm random variable in Definition 3.9.
- Form Definition 5.18, each component Z_i of $\mathbf{Z}$ has expected value $E[Z_i] = 0$ and variance $\mathrm{Var}[Z_i] = 1$. Thus Z_i is a Gausian (0, 1) random variable. In addition, $E[Z_i\, Z_j] = 0$ for all $i \neq j$. Since $\mathbf{C_Z}$ is a diagonal matrix, $Z_1, \ldots, Z_n$ are independent.

- In many situations, it is useful to transform the Gaussian (μ, σ) random variable X to the standard normal random variable $Z = (X - \mu_X) / \sigma_X$. For Gaussian vectors, we have a vector transformation to transform $\mathbf{X}$ into a standard random vector.

Theorem 5.17

For a Gaussian $(\boldsymbol{\mu}_{\mathbf{X}}, \mathbf{C}_{\mathbf{X}})$ random vector, let $\mathbf{A}$ be an $n \times n$ matrix with the property $\mathbf{A}\mathbf{A}' = \mathbf{C}_{\mathbf{X}}$. The random vector

$$\mathbf{Z} = \mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})$$

is a standard normal random vector.

Proof: Theorem 5.17

Applying Theorem 5.16 with $\mathbf{A}$ replaced by $\mathbf{A}^{-1}$, and $\mathbf{b} = \mathbf{A}^{-1}\boldsymbol{\mu}_{\mathbf{X}}$, we have that $\mathbf{Z}$ is a Gaussian random vector with expected value

$$E[\mathbf{Z}] = E\left[\mathbf{A}^{-1}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})\right] = \mathbf{A}^{-1}E\left[\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}}\right] = \mathbf{0},$$

and covariance

$$\mathbf{C}_{\mathbf{Z}} = \mathbf{A}^{-1}\mathbf{C}_{\mathbf{X}}(\mathbf{A}^{-1})' = \mathbf{A}^{-1}\mathbf{A}\mathbf{A}'(\mathbf{A}')^{-1} = \mathbf{I}.$$

- The transformation in this theorem is considerably less straightforward than the scalar transformation $Z = (X - \mu_X) / \sigma_X$, because it is necessary to find for a given $\mathbf{C_X}$ a matrix $\mathbf{A}$ with the property $\mathbf{AA}' = \mathbf{C_X}$. The calculation of $\mathbf{A}$ from $\mathbf{C_X}$ can be achieved by applying the linear algebra procedure **singular value decomposition**.
- Section 5.8 describes this procedure in more detail and applies it go generating sample values of Gaussian random vectors.
- The inverse transform of Theorem 5.17 is particularly useful in computer simulations.

Theorem 5.18

Given the n-dimensional standard normal random vector $\mathbf{Z}$, an invertible $n \times n$ matrix $\mathbf{A}$, and an n-dimensional vector $\mathbf{b}$,

$$\mathbf{X} = \mathbf{A}\mathbf{Z} + \mathbf{b}$$

is an n-dimensional Gaussian random vector with expected value $\boldsymbol{\mu}_{\mathbf{X}} = \mathbf{b}$ and covariance matrix $\mathbf{C}_{\mathbf{X}} = \mathbf{A}\mathbf{A}'$.

Proof: Theorem 5.18

By Theorem 5.16, $\mathbf{X}$ is a Gaussian random vector with expected value

$$\boldsymbol{\mu}_{\mathbf{X}} = E[\mathbf{X}] = E\left[\mathbf{A}\mathbf{Z} + \boldsymbol{\mu}_{\mathbf{X}}\right] = \mathbf{A}E[\mathbf{Z}] + \mathbf{b} = \mathbf{b}.$$

The covariance of $\mathbf{X}$ is

$$\mathbf{C}_{\mathbf{X}} = \mathbf{A}\mathbf{C}_{\mathbf{Z}}\mathbf{A}' = \mathbf{A}\mathbf{I}\mathbf{A}' = \mathbf{A}\mathbf{A}'.$$

- Theorem 5.18 says that we can transform the standard norm vector $\mathbf{Z}$ into a Gaussian random vector $\mathbf{X}$ whose covariance matrix is of the form $\mathbf{C_X} = \mathbf{AA}'$.
- The usefulness of Theorem 5.17 and 5.18 depends on whether we can always find a matrix A such tat $\mathbf{C_X} = \mathbf{AA}'$.
- In fact, as we verify below, Theorem 5.19 is possible for every Gaussian vector $\mathbf{X}$.

Theorem 5.19

For a Gaussian vector $\mathbf{X}$ with covariance $\mathbf{C_X}$, there always exists a matrix $\mathbf{A}$ such that $\mathbf{C_X} = \mathbf{A}\mathbf{A}'$.

Proof: Theorem 5.19

To verify this fact, we connect some simple facts:

- In Problem 5.6.9, we ask the reader to show that every random vector $\mathbf{X}$ has a positive semidefinite covariance matrix $\mathbf{C_X}$. By Math Fact B.17, every eigenvalue of $\mathbf{C_X}$ is nonnegative.
- The definition of the Gaussian vector PDF requires the existence of $\mathbf{C_X}^{-1}$. Hence, for a Gaussian vector $\mathbf{X}$, all eigenvalues of $\mathbf{C_X}$ are nonzero. From the previous step, we observe that all eigenvalues of $\mathbf{C_X}$ must be positive.
- Since $\mathbf{C_X}$ is a real symmetric matrix, Math Fact B.15 says it has a singular value decomposition (SVD) $\mathbf{C_X} = \mathbf{UDU}'$ where $\mathbf{D} = \text{diag}[d_1, \ldots, d_n]$ is the diagonal matrix of eigenvalues of $\mathbf{C_X}$. Since each d_i is positive, we can define $\mathbf{D}^{1/2} = \text{diag}[\sqrt{d_1}, \ldots, \sqrt{d_n}]$, and we can write

$$\mathbf{C_X} = \mathbf{UD}^{1/2}\mathbf{D}^{1/2}\mathbf{U}' = \left(\mathbf{UD}^{1/2}\right)\left(\mathbf{UD}^{1/2}\right)'.$$

We see that $\mathbf{A} = \mathbf{UD}^{1/2}$.

Quiz 5.7

$\mathbf{Z}$ is the two-dimensional standard normal random vector. The Gaussian random vector $\mathbf{X}$ has components

$$X_1 = 2Z_1 + Z_2 + 2 \quad \text{and} \quad X_2 = Z_1 - Z_2.$$

Calculate the expected value $\boldsymbol{\mu}_{\mathbf{X}}$ and the covariance matrix $\mathbf{C}_{\mathbf{X}}$.

Quiz 5.7 Solution

We observe that $\mathbf{X} = \mathbf{A}\mathbf{Z} + \mathbf{b}$ where

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}, \qquad \mathbf{b} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}.$$

It follows from Theorem 5.18 that $\boldsymbol{\mu}_X = \mathbf{b}$ and that

$$\mathbf{C}_X = \mathbf{A}\mathbf{A}' = \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 5 & 1 \\ 1 & 2 \end{bmatrix}.$$