

CAUSE-OF-DEATH

CAUSES -OF-DEATH

“Around 56 million peoples have die each year”

- What caused their death?
- How did the causes of death change over time and differ between different countries and world regions?
- What are the risk factors that lead to early death?

These are the big questions we are answering here.

Definitions: Cause of death vs risk factors

In the epidemiological framework of the Global Burden of Disease study each death has one specific cause. In their own words: ‘each death is attributed to a single underlying cause the cause that initiated the series of events leading to death.

This is different from the deaths that happened due to risk factors. These deaths are an estimation of the reduction of the number of deaths that would be achieved if the risk factors to which a population is exposed would be eliminated(in case of tobacco smoking, for example) or reduced to an optimal. Healthy level(in the case of body-mass index).

What do people die from?

The Global Burden of Disease is a major global study on the causes of death and disease published in the medical journal The Lancet. These estimates of the annual number of deaths by cause are shown here.

This is shown for deaths worldwide. But you can explore data on the annual number of deaths by cause for any country or region using the “change country” toggle.

Non-communicable diseases(NCDs) not only dominate mortality figures at a global level, but also account for the majority of deaths in high-income countries.

Deaths from causes such a infectious disease, malnutrition, nutritional deficiencies, neonatal and maternal deaths are common-and in some cases

CAUSE-OF-DEATH

dominant-across low-and middle-income nations. In Kenya, for example, the leading cause of death remains diarrheal diseases. In South Africa and Botswana, the leading cause of death is HIV/AIDS. In high-income countries however the share of deaths caused by these is very low.

Using the timeline on the chart you can also explore how deaths by cause have changed over time.

Death rates related to disease, illness and other health factors tend to change relatively slowly over time. /whilst death rates may fall or decline from year-to-year as part of a general trend, dramatic change in such deaths are typically rare. Natural disaster and terrorism-related deaths are an important exception to this rule, as they can vary significantly between countries. This can make the annual comparison of deaths and death rates between health-related factors and volatile events more challenging. Understanding the relative risk of these events can require a longer-term overview of high and low-mortality years.

Causes of death by category:

At a global level we see that the majority of deaths are caused by non communicable diseases(NCDs). Collectively NCDs account for more than 73% of global deaths. As the world is making progress in the fight against many infectious diseases, and as population age, we expect that NCDs will become increasingly dominant as the cause of death.

Breakdown of deaths by age

In 2017, there were 56.5 million deaths globally just over half of these were people who were 70 years or older ,26% were between 50 and 69 years old,13% were between 15 and 49, only 1% were older than 5 and younger than 14, and almost 9% were children under the age of 5.

The age at which people die has changed significantly since 1990. Fewer people die at a young age. In 1990 nearly one-quarter of all deaths were in children younger than 5. In 2019, this had declined to just under 9%. In contrast, the share

CAUSE-OF-DEATH

of deaths in the over-70s age bracket has increased from a third to half of all deaths over this period.

It is possible to change this chart to any other country or region in the world. In countries with good health the share dying at a young age is very low. In Japan more than 85% are 70 years or older.

Causes of deaths of children younger than 5

Through the combination of neonatal (newborn infants less than 28 days old) disorders, infections and congenital (from birth) defects, we see that the largest share of deaths in under-5s arises from complications at birth or in the first few weeks of life. Under-5s are also highly susceptible to lower respiratory infections, infectious diseases, diarrheal infections, malnutrition and nutritional deficiencies.

This is shown for deaths worldwide. But you can explore data on the annual number of deaths by cause for any country or region using the “change country” toggle.

Death rates in under-5s are typically much lower in high-income countries, and the nature of these deaths is different from lower incomes. In the United Kingdom, for example, child deaths tend to be highly dominated by neonatal complications. Deaths from infectious and diarrheal diseases and malnutrition is very low. In contrast, infectious diseases and nutritional deficiencies are large causes of death in lower-income countries.

Causes of deaths for children between 5 and 14

Globally, deaths in the 5-14 year old age bracket account for a small percentage of the total (1-2%)

There are six dominant causes of deaths in this age category. The leading causes globally in 5-14 year olds are road accidents, cancers and malaria. Lower respiratory infections, HIV/AIDS, diarrheal diseases, and drowning are all dominant causes typically in the range of 40000-50000 deaths in 2017.

CAUSE-OF-DEATH

Again, this distribution varies by country. In the United States, for example, cancers are the leading cause of death. In India, it's diarrheal diseases, in Bangladesh and China it's drowning, and in South Africa HIV/AIDS.

Causes of deaths for 15 to 49 year olds

In the 15 to 49 years old category, we see that non-communicable diseases(NCDs) begin to become dominant. Globally the leading cause of death in this age group is cardiovascular diseases, followed cancers which both account for more than one million deaths. Road accidents, HIV/AIDS and suicide are all significant within this group.

For some countries, such as South Africa, by far the dominant cause of death is HIV/AIDS in 15 to 49 year olds. In a number of countries(in particular across Latin America, including Brazil and Mexico), homicide is the dominant cause for 15-49 years old.

Cause of deaths for 50 to 69 years olds

In 50 to 69 year olds, non-communicable diseases(NCDs) are strongly dominant- here cardiovascular disease, cancers, respiratory disease and diabetes are the top causes. With the exception of HIV/AIDS and tuberculosis which or some countries climb into the top causes, the global variability in death causes for 50-69 year olds is much lower than that of younger age categories.

Causes of deaths for people who were older than 69 years

For the oldest age category(70 years and older), non-communicable diseases(NCDs) still dominate, however other death causes including Alzheimer's/dementias, and diarrheal diseases also become dominant. Diarrheal diseases remain within the few leading causes of deaths in 70+ year olds for many low-income countries, despite being relatively low at higher incomes.

Risk factors for death

It is important to understand what is meant by the cause of death and the risk factor associated with a premature death:

CAUSE-OF-DEATH

In the epidemiological framework of the Global Burden of Disease study each death has one specific cause. In their own words: 'each death is attributed to a single underlying cause-the cause that initiated the series of events leading to death'.

This is different from the deaths that happened due to risk factors. These deaths are an estimation of the reduction of the number of deaths that would be achieved if the risk factors to which a population is exposed would be eliminated(in the case of tobacco smoking, for example) or reduced to an optimal, healthy level(in the case of body-mass index). Risk factors can be grouped into four broad categories: behavioural risks, environmental risks, occupational risks, and metabolic risks.

All of these estimates are developed independently. This means that we cannot sum all 'attributed number of deaths by risk factor in many cases exceeds that of those by cause of death.

Below, in our section on Measurement, we describe in more detail how the epidemiologists of the Global Burden of Disease study attribute risk factors to mortality.

The number of deaths by risk factor

Here we see that there are several dominant risk factors for death: notably, those related to dietary and activity lifestyle factors (including blood pressure, physical activity, body-mass index, blood sugar, and dietary intake), smoking, air pollution(both outdoor and indoor), environmental factors including clean water and sanitation, and safe sex (for the prevention of HIV/AIDS).

This is shown for death worldwide. But you can explore data on the annual number of deaths by cause for any country or region using the "change country" toggle. The contribution of specific risk factors varies significantly by country.

For most high-income countries, the dominant risk factors are those related to healthy diets, smoking and alcohol intake. Other risk factors such as clean water, sanitation, and child wasting or stunting are very low. In low-income countries the

CAUSE-OF-DEATH

inverse is true: in Sierra Leone for example, the top risk factors include child wasting, household air pollution, unsafe water source, poor sanitation, and the lack of access to handwashing facilities. For countries where HIV/AIDS is a major health burden, such as South Africa and Kenya, unsafe sex is the top risk factor.

Causes:-

1.Cardiovascular diseases

Cardiovascular disease(CVD) is a term used to refer to the range of diseases which affect the heart and blood vessels. These include hypertension(high blood pressure), coronary heart disease(heart attack), cerebrovascular disease(stroke), heart failure, and other heart diseases.

Cardiovascular disease is the top cause of death globally.

2.Cancers

Cancers are defined by the National Cancer Institute as a collection of diseases in which abnormal cells can divide and spread to nearby tissue. Cancers can arise in many parts of the body-leading to a range of cancer types, and in some cases spread to other parts of the body through the blood and lymph systems.

3.Dementia

Dementia comprises several forms- the most common being Alzheimer's disease- is an illness which results in a deterioration of cognitive capacity and function beyond what is expected from the normal aging process. It can occur either in a chronic or progressive form. It affects several cognitive functions including memory, comprehension, judgement, language and learning capacity.

4.Diarrheal diseases

Diarrheal diseases are caused primarily by viral and bacterial pathogens. They are particularly dominant at lower incomes where there is poor access to safe sanitation, drinking water and hygiene facilities. Diarrheal diseases are a leading cause of death in children.

CAUSE-OF-DEATH

5.Tuberculosis

Tuberculosis(TB) is an illness caused by the ingestion of bacteria (*Mycobacterium tuberculosis*) which affects the lungs. The World Health Organization(WHO) estimate that up to one quarter of the global population has latent TB, meaning they have been infected with the disease but are not ill with the disease(although this does not inhibit it from becoming active in the future).

People with compromised immune system, such as those suffering from malnutrition, diabetes, or are smokers are more likely to become ill with TB. There is a strong link between HIV/AIDS and TB: those infected with HIV are 20-30 times more likely to develop active tuberculosis.

6.Malnutrition

Malnutrition arises in various forms, with the broad definition capturing undernourishment, micronutrient deficiencies and obesity . In this case, we refer to 'protein-energy malnutrition(PEM) which refers to energy or protein deficiency caused by insufficient food intake. Protein-energy deficiency can also be exacerbated by infection or disease, which can have the effect of increasing nutritional needs, and/or reducing the body's ability to retain energy or nutrients.

7.HIV/AIDS

An infection with HIV(human immunodeficiency virus) can lead to AIDS(acquired immunodeficiency syndrome).AIDS results in a gradual and persistent decline and failure of the immune system, resulting in heightened risk of life-threatening infection and cancers.

In the majority of cases, HIV is a sexually-transmitted infection. However, HIV can also be transmitted from a mother to her child, during pregnancy or childbirth, or through breastfeeding. Non-sexual transmission can also occur through the sharing of injection equipment such as needles.

CAUSE-OF-DEATH

8.Malaria

Malaria is a disease that is transmitted from person to person by infected mosquitoes. The bite of an infected Anopheles mosquito transmits a parasite that enters the victim's blood system and travels into the person's liver where the parasite reproduces. There the parasite causes a high fever that involves shaking chills and pain. In the worst cases malaria leads to coma and death.

9.Smoking

Tobacco smoking is not a direct cause of death, but it nonetheless one of the world's largest health problems.

Smoking is one of the world's leading risk factors for premature death. Tobacco a risk factor for several of the world's leading causes of death, including lung and other forms of cancer, heart disease, and respiratory diseases.

10.Suicide

Every suicide is a tragedy. With timely, evidence-based interventions, suicides can be prevented.

11.Homicides

Intentional homicides are defined as "an unlawful death deliberately inflicted on one person by another person. Civilian and military deaths during interstate wars, civil wars and genocides are not counted as homicides

12.Natural disasters

Natural disasters can occur in many forms-ranging from earthquakes and tsunamis, to extreme weather events, and heat waves.

The largest disaster events are often infrequent, but high-impact meaning there is significant variability in deaths from year-to-year.

13.Road incidents

Road incident deaths include those of drivers-motor vehicles and motorcyclists-in addition to cyclists and pedestrian deaths.

CAUSE-OF-DEATH

14.Drowning

The World Health Organization(WHO) emphasis that drowning is one of the most overlooked, preventable causes of death across the world. For every country in the world, drowning is among the top 10 killers for children. In some countries, such as Bangladesh, it is the top mortality cause for children under 15 years old.

15.Fire

Most countries across the Americas, Western Europe, East Asia and Oceania average death rates below 2 per 100,000. Rates across other regions are typically higher at 2-6 per 100,000. When viewed through time we see a notable decline in fire death rates, particularly across Sub-Saharan Africa and Eastern Europe.

16.Terrorism

Terrorism is defined in the Oxford Dictionary as “the unlawful use of violence and intimidation, especially against civilians, in the pursuit of political aims.” We quickly see that this definition is unspecific and subjective.

17.Deaths by animal

Mosquitoes are by far the world’s deadliest animal. Around 1.5 million people are killed by animals every year.

More than half a million are killed by other humans-in war, homicides, and terrorism. And close to a million people are killed by other animals in any given year.

Mosquitoes are, by far, the world’s deadliest animal for humans: at estimated 780,000 died from the transmission of disease from mosquitoes in 2016.

Mosquito deaths are the sum of deaths(in order, highest to lowest) from: Malaria, Dengue fever, Japanese encephalitis, Yellow fever, Zika virus, Chikungunya, West Nile virus, and Lymphatic filariasis, for which it is the vector.

What we die from, what we Google, what we read in the news?

CAUSE-OF-DEATH

The coverage in both newspapers here is strikingly similar. And the discrepancy between what we actually die from and what we get informed of in the media is what stands out:

Around one-third of the considered causes of deaths resulted from heart disease, yet this cause of death receives only 2-3 percent of Google searches and media coverage.

Just under one-third of the deaths came from cancer, we actually Google cancer a lot (37 percent of searches) and it is a popular entry. Here on our site, but it receives only 13-14 percent of media coverage.

We searched for road incidents more frequently than their share of deaths, however, they receive much less attention in the news.

When it comes to deaths from strokes, Google searches and media coverage are surprisingly balanced.

The largest discrepancies concern violent forms of death: suicide, homicide and terrorism. All three receive much more relative attention in Google searches and media coverage than their relative share of deaths. When it comes to the media coverage on causes of death, violent deaths account for more than two-thirds of coverage in the New York Times and The Guardian but account for less than 3 percent of the total deaths in the US.

What's interesting is that what Americans search on Google is a much closer reflection of what kills us than what is presented in the media. One way to think about it is that media outlets may produce content that they think readers are most interested in, but this is not necessarily reflected in our preferences when we look for information ourselves.

How over-or underrepresented are deaths in the media?

As we can see clearly from the chart above, there is a disconnect between what we die from, and how much coverage these causes get in the media. Another way to summarize this discrepancy is to calculate how over-or underrepresented each

CAUSE-OF-DEATH

cause is in the media. To do this, we simply calculate the ratio between the share of deaths and share of media coverage for each cause.

Should media exposure reflect what we die from?

There are several reasons we would, or should, expect that what we read online, and what is covered in the media wouldn't correspond with what we actually die from.

The thing is that we would expect there to be some preventative aspect to information we access. There's a strong argument that things we search for and gain information on encourages us to take action which prevents a further death. There are several examples where I can imagine this to be true. People who are concerned about cancer may search online for guidance on symptoms and be convinced to see their doctor. Some people with suicidal thoughts may seek help and support online which later results in an averted death from suicide. We'd therefore expect that both intended or unintended exposure to information on particular topics could prevent deaths from a given cause. Some imbalance in the relative proportions therefore makes sense. But clearly there is some bias in our concerns: most people die from heart disease (hence it should be something that concerns us) yet only a small minority seek [possibly preventative] information online.

How to combat our bias for single events

Media and its consumers are stuck in a reinforcing cycle. The new reports on breaking events, which are often based around a compelling story. Consumers want to know what's going on in the world-we are quickly immersed by the latest headline. We come to expect news updates with increasing frequency, and media channels have clear incentives to deliver. This locks us into a cycle of expectation and coverage with a strong bias for outlier events. Most of us are left with a skewed perception of the world, we think the world is much worse than it is

The responsibility in breaking this cycle lies with both media producers and consumers. Will we ever stop reporting and reading the latest news? Unlikely. But

CAUSE-OF-DEATH

we can all be more conscious of how we let this news shape our understanding of the world.

And journalists can do much better in providing context of the broader trends: if reporting on a homicide, for example, include context of how homicide rates are changing over time.

As media consumers we can be much more aware of the fact that relying on the 24/7 news coverage one is wholly insufficient for understanding the state of the world. This requires us to check our (often unconscious) bias for single narratives and seek out sources that provide a fact-based perspective on the world.

This antidote to the news is what we try to provide at Our World in Data. It should be accessible for everyone, which is why our work is completely open access. Whether you are a media producer or consumer, feel free to take and use anything you find here.

Estimating the causes of death

The IHME's Global Burden of Disease (GBD) has developed a standardized approach to the attribution of deaths to specific causes.

Their methodology states that "each death is attributed to a single underlying cause-the cause that initiated the series of events leading to death-in accordance with ICD principles. Data to estimate the causes of deaths is far from complete, particularly in poorer countries, and for this estimation the researchers therefore need to rely on various sources. These sources include vital registration (VR), verbal autopsy (VA), surveillance, census and survey data, cancer registries, and police records. GBD then develops a data standardization and processing methodology within which they define data quality and completeness scores, and where necessary adjust completeness to 100% using cause fractions for a given location-age-sex-year and estimated all-cause mortality for that location-age-sex-year.

CAUSE-OF-DEATH

The GBD assessment is strongly tied to the mortality cause categories as defined within the International Classification of Diseases(ICD). Codes, as used by the World Health Organization(WHO).

CAUSE-OF-DEATH