

STATISTICS WORKSHEET-5

Q1 to Q10 are MCQs with only one correct answer. Choose the correct option.

- 1. Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.
 - a) Mean
 - b) Actual
 - c) Predicted
 - d) Expected
- 2. Chisquare is used to analyse
 - a) Score
 - b) Rank
 - c) Frequencies
 - d) All of these
- 3. What is the mean of a Chi Square distribution with 6 degrees of freedom?
 - a) 4
 - b) 12
 - c) 6
 - d) 8
- 4. Which of these distributions is used for a goodness of fit testing?
 - a) Normal distribution
 - b) Chisqured distribution
 - c) Gamma distribution
 - d) Poission distribution
- 5. Which of the following distributions is Continuous
 - a) Binomial Distribution
 - b) Hypergeometric Distribution
 - c) F Distribution
 - d) Poisson Distribution
- 6. A statement made about a population for testing purpose is called?
 - a) Statistic
 - b) Hypothesis
 - c) Level of Significance
 - d) TestStatistic
- 7. If the assumed hypothesis is tested for rejection considering it to be true is called?
 - a) Null Hypothesis
 - b) Statistical Hypothesis
 - c) Simple Hypothesis
 - d) Composite Hypothesis
- 8. If the Critical region is evenly distributed then the test is referred as?
 - a) Two tailed
 - b) One tailed
 - c) Three tailed
 - d) Zero tailed
- 9. Alternative Hypothesis is also called as?
 - a) Composite hypothesis
 - b) Research Hypothesis
 - c) Simple Hypothesis
 - d) Null Hypothesis



10. In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by ______a) np

b) n

