

CH5440 HW5

Prem Sagar S - AE14B021

April 25, 2017

Problem 1

H is the pure component spectra obtained from NMF. P is the true pure component spectra.

	H(1,:)	H(2,:)	H(3,:)
P(1,:)	0.5545729916	-0.5707058235	0.1294911335
P(2,:)	0.2004820905	0.7188547893	-0.2450129603
P(3,:)	-0.4946498902	0.821385503	0.4928965533

Table 1: Correlation matrix obtained for the case where only the first of the 5 trials were taken. Here the 2nd and 3rd component of H are both related to 3rd component of P. This is inaccurate.

	H(1,:)	H(2,:)	H(3,:)
P(1,:)	0.7586127475	-0.0460677071	-0.6464171415
P(2,:)	-0.0912255659	0.056043019	0.8390951193
P(3,:)	-0.6833443025	0.7950965931	0.7335536387

Table 2: Correlation matrix obtained for the case where the mean of the 5 trials were taken. The mapping is uniquely obtained and the correlation is also better than the previous case.

The spectra are compared below for the mean data.




Figure 1: Pure spectra of Co




Figure 2: Pure spectra of Ni




Figure 3: Pure spectra of Cr

Problem 2

Part a

$E = \text{measabs}'$ (N variables x M samples) = (7x176)

Connectivity matrix A (NxL=7x3)

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Criterion 1: A must be full column rank

From matlab the rank was found to be 3. Hence this condition is satisfied.

Criterion 2: Each column of A must have atleast L-1=2 zeros which is satisfied.

Criterion 3: If $E=AP$, P must be a full row matrix. A necessary condition for this is $L < M$, or $3 < 176$. This condition is satisfied.

Hence the given connectivity is NCA compliant

Part b

The rotation matrix M transforms the PCA estimated connectivity matrix to the real struture.

$$struc(M\hat{A}^T) = struc(A^T)$$

The second column of the connectivity matrix has 3 zeros. This makes it an overdetermined system of equations. Either only two equations can be used or the a least squares estimate can be obtained.

The least squares solution for the rotation matrix is as below:

$$M = \begin{bmatrix} 1 & 0.850658019 & 1.2257287588 \\ -0.0177462979 & 1 & 0.0029501312 \\ 0.3712629247 & -0.9856172775 & 1 \end{bmatrix}$$

A better rotation matrix that provided a near structure of A was obtained by using only the 2nd and 3rd zeros of the 2nd column.

$$M = \begin{bmatrix} 1 & 0.850658019 & 1.2257287588 \\ 0.4343763995 & 1 & -0.83980366252 \\ 0.3712629247 & -0.9856172775 & 1 \end{bmatrix}$$

The pure component spectra extracted and the true pure component spectra had a correlation of **0.7747**.

Part c

The NCA toolbox was used to obtain the pure component spectra. The problem solved is essentially, finding the pure component spectra matrix $P_{3 \times 176}$ having known the connectivity matrix $A_{7 \times 3}$ and the absorbance matrix, $E_{7 \times 176}$, measured for 176 wavelengths.

$$E = AP$$

The correlation coefficient between the true pure component spectra and the measured pure component spectra is **0.9480**.

The spectra is shown below. The spectra match to a scale but has too much noise.




Figure 4: Pure spectra of Co




Figure 5: Pure spectra of Cr




Figure 6: Pure spectra of Ni