

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

---

### 1. WHY AM I USING PHARMACOR DABIGATRAN?

PHARMACOR DABIGATRAN contains the active substance dabigatran etexilate (as dabigatran etexilate mesilate). It belongs to a group of medicines called anticoagulants. Dabigatran works by inhibiting a specific protein in the blood, called thrombin. Thrombin contributes to the formation of blood clots. Dabigatran prevents the formation of blood clots. For more information, see Section [1. Why am I using PHARMACOR Dabigatran?](#) in the full CMI.

### 2. WHAT SHOULD I KNOW BEFORE I USE PHARMACOR DABIGATRAN?

Do not use if you have ever had an allergic reaction to PHARMACOR Dabigatran or any of the ingredients listed at the end of the CMI or if you become pregnant while you are taking PHARMACOR Dabigatran.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.** For more information, see Section [2. What should I know before I use PHARMACOR DABIGATRAN?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with PHARMACOR DABIGATRAN and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I take PHARMACOR DABIGATRAN?

#### AFTER KNEE AND HIP REPLACEMENT SURGERY

The recommended dose of PHARMACOR DABIGATRAN is 220 mg (2 capsules of 110 mg) taken as a single dose once daily.

- Patients with moderately reduced kidney function (e.g. over 75 years) or patients taking certain medicines may have an increased risk of bleeding. The doctor may prescribe the lower dose of 150 mg once daily, taken as 2 capsules of PHARMACOR DABIGATRAN 75 mg.
- Treatment with PHARMACOR Dabigatran should be started within 1 - 4 hours of completed surgery, using a single capsule of 110 mg and continuing with 2 capsules of 110 mg once daily for a total of 10 days.

#### AFTER HIP REPLACEMENT SURGERY

- Treatment with PHARMACOR DABIGATRAN should be started within 1 - 4 hours of completed surgery, using a single capsule of 110 mg and continuing with 2 capsules of 110 mg once daily for a total of 28 - 35 days.
- If, within 4 hours after surgery, post-operative bleedings can still be observed, initiation of treatment should be delayed. If treatment is not started on the day of surgery then treatment should be initiated with 2 capsules of 110 mg once daily.

## FOR STROKE PREVENTION IN PATIENTS WITH ATRIAL FIBRILLATION

- The recommended dose of PHARMACOR DABIGATRAN is 300 mg taken as 1 capsule of 150 mg in the morning and 1 capsule of 150 mg in the evening.
- Patients over 75 years should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening.
- Patients with an increased risk of major bleeding (as determined by your doctor) should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening.

## FOR THE TREATMENT OF BLOOD CLOTS AND PREVENTION OF BLOOD CLOTS RE-OCCURRING IN THE VEINS OF YOUR LEGS AND/OR LUNGS

- The recommended dose of PHARMACOR DABIGATRAN is 300 mg taken as 1 capsule of 150 mg twice a day following treatment with an injectable blood thinner for at least 5 days. To prevent blood clots re-occurring, continue on 1 capsule of 150 mg twice a day.
- More instructions can be found in Section 4. How do I use PHARMACOR Dabigatran? in the full CMI.

### 5. What should I know while using PHARMACOR DABIGATRAN?

|                                    |   |
|------------------------------------|---|
| <b>Things you should do</b>        | <ul style="list-style-type: none"><li>• Remind any doctor, dentist or pharmacist you visit that you are taking PHARMACOR DABIGATRAN.</li><li>• Have your blood cholesterol and triglycerides checked when your doctor says so. Stop taking PHARMACOR Dabigatran and seek medical attention immediately if you experience any of the serious side effects listed in Section 6 in the full CMI.</li></ul> |
| <b>Things you should not do</b>    | <ul style="list-style-type: none"><li>• Do not stop taking PHARMACOR DABIGATRAN unless you have discussed it with your doctor.</li></ul>  |
| <b>Driving or using machines</b>   | <ul style="list-style-type: none"><li>• Be careful driving a car or operating machinery as PHARMACOR DABIGATRAN may cause dizziness in some people.</li></ul>   |
| <b>Drinking alcohol</b>            | <ul style="list-style-type: none"><li>• Excessive alcohol consumption may not be safe in patients taking PHARMACOR DABIGATRAN.</li></ul>  |
| <b>Looking after your medicine</b> | <ul style="list-style-type: none"><li>• Keep your tablets in the blister pack until it is time to take them. Store your medicine in a cool dry place below 25°C. Keep your medicine where children cannot reach it.</li></ul>   |

For more information, see Section 5. What should I know while using PHARMACOR DABIGATRAN? in the full CMI.

### 6. Are there any side effects?

The most common side effects of PHARMACOR DABIGATRAN are bruising, nose bleeds, stomach ache, itchy skin, rash, diarrhoea, indigestion, feeling sick, cough, painful, swollen joints, sore nasal passages and throat, discomfort when swallowing, hair loss, frequent infections such as fever, severe chills, sore throat or mouth ulcers (signs of lack of white blood cells). These side effects are usually mild. **Tell your doctor immediately or go to Emergency at your nearest hospital if you notice any of the following:** long or excessive bleeding, exceptional weakness, tiredness, headaches, dizziness and looking pale (signs of anaemia), chest pain or being short of breath, swelling of hands, ankles and feet, red or dark brown urine, red or black bowel motions, symptoms related to a condition called anticoagulant-related nephropathy: blood in urine, reduced urine output, swelling of the legs, ankles and feet, increased time for blood to clot (high INR test values), heavy bleeding. These are serious side effects. You may need urgent medical attention. For more information, including what to do if you have any side effects, see Section 6. Are there any side effects? in the full CMI.

## **Consumer Medicine Information (CMI)**

This leaflet provides important information about using [PHARMACOR DABIGATRAN]. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using [PHARMACOR DABIGATRAN].**

### **Where to find information in this leaflet:**

1. Why am I taking PHARMACOR DABIGATRAN?

2. What should I know before I use PHARMACOR DABIGATRAN?

3. What if I am taking other medicines?

4. How do I take PHARMACOR DABIGATRAN?

5. What should I know while using PHARMACOR DABIGATRAN?

6. Are there any side effects?

7. Product details

## **1. Why am I using PHARMACOR DABIGATRAN?**

PHARMACOR DABIGATRAN contains the active substance dabigatran etexilate (as dabigatran etexilate mesilate). It belongs to a group of medicines called anticoagulants. Some people refer to anticoagulant medicines as "blood thinners".

Dabigatran works by inhibiting a specific protein in the blood, called thrombin. Thrombin contributes to the formation of blood clots. Dabigatran prevents the formation of blood clots.

PHARMACOR DABIGATRAN has been prescribed to you for one of the following uses:

- to prevent the formation of blood clots in the veins after knee or hip replacement surgery in adults
- to reduce the risk of brain (stroke) and/or other body vessel obstruction by blood clot formation in adults with an abnormal heart beat rhythm called non-valvular atrial fibrillation .
- to treat blood clots in the veins of your legs and lungs and to prevent blood clots from re-occurring in the veins of your legs and/or lungs.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

## **2. What should I know before I take PHARMACOR DABIGATRAN?**

When you must not take it

**Do not take PHARMACOR DABIGATRAN if you are allergic to:**

- dabigatran etexilate or any of the other ingredients in PHARMACOR DABIGATRAN listed at the end of this leaflet. Some of the symptoms of an allergic reaction may include:
- rash, itching or hives on the skin
- swelling of the face, lips, tongue or other parts of the body
- shortness of breath, wheezing or troubled breathing.

**Do not take PHARMACOR DABIGATRAN:**

- if you are currently bleeding
- if you have severely reduced kidney function (your doctor will know how to determine your kidney function)
- if you have an increased tendency of bleeding complications (this may be inherited, of unknown cause or due to other medicines)
- if you have a medical condition which increases your risk of serious bleeding, such as recent brain or spinal injury, and cancer
- if you have active stomach ulcers or have experienced stomach bleeding in the past year, unless the cause has been permanently eliminated, e.g. by surgery
- if you have a history of bleeding in the head, eyes, spine, abdomen and joints
- if you have an indwelling spinal or epidural catheter, and during the first two hours after their removal (your doctor will know about the kind of catheters and precautionary measures)
- if you have liver problems or liver disease
- if you are currently taking oral ketoconazole or itraconazole, medicines used to treat fungal infections

- if you are taking dronedarone, a medicine used to prevent repetition of your problem of irregular heart beat
- if you are taking cyclosporin or tacrolimus, medicines used to prevent organ rejection after transplantation
- if you are taking medicines to prevent blood clotting (e.g. warfarin, rivaroxaban, apixaban or heparin), except when changing anticoagulant treatment or while having a venous or arterial line and you get heparin through this line to keep it open .
- if you have a prosthetic heart valve.

**Do not start PHARMACOR DABIGATRAN and verapamil treatment at the same time.**

**Do not start verapamil if you are currently taking PHARMACOR DABIGATRAN and have just undergone major orthopaedic surgery.**

**Do not take this medicine if you are pregnant.**  
It may affect your developing baby if you take it during pregnancy.

**Do not breastfeed if you are taking this medicine.**

The active ingredient in PHARMACOR DABIGATRAN passes into breast milk.

**Do not give this medicine to a child or adolescent under 18 years old .**

There is no experience with its use in children or adolescents under 18 years old.

**Do not take this medicine after the expiry date printed on the box or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal. If you use this medicine after the expiry date has passed, it may not work as well.

**If you are not sure whether you should start taking this medicine, talk to your doctor.**

*Before you use it*

**Tell your doctor if you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes.**

**Tell your doctor if you are pregnant or intend to become pregnant.**

PHARMACOR Dabigatran should not be used during pregnancy.

**Tell your doctor if you are breastfeeding or intend to breastfeed.**

PHARMACOR DABIGATRAN is not recommended in women who are breastfeeding.

**Tell your doctor if you have had a heart attack or if you have been diagnosed with conditions that increase the risk to develop a heart attack.**

**Tell your doctor if you have reduced liver function, life-threatening liver disease or increased liver enzymes.**

**Tell your doctor if you have an increased bleeding risk, as could be the case in the following situations:**

- if you are older than 75 years, your doctor may prescribe a lower dose of PHARMACOR DABIGATRAN
- if you know you have reduced kidney function, or you are suffering from dehydration (symptoms include feeling thirsty and passing reduced amounts of dark-coloured urine)
- if you have been recently bleeding
- if you have any problems with your blood
- if you have had a recent tissue sampling (biopsy)
- if you have cancer
- if you have had a serious injury (e.g. a bone fracture, head injury or any injury requiring treatment)
- if you are suffering from an inflammation of the food pipe (oesophagus) or stomach
- if you have problems with reflux of gastric juice into the food pipe (oesophagus)
- if you are receiving medicines which could increase the risk of bleeding, such as clopidogrel and warfarin
- if you are taking anti-inflammatory medicines such as diclofenac
- if you are suffering from an infection of the heart (bacterial endocarditis).

**• Tell your doctor if you know that you have a disease called antiphospholipid syndrome (a disorder of the immune system that causes an increased risk of blood clots).**

**If you have not told your doctor about any of the above, tell them before you use PHARMACOR DABIGATRAN.** See additional information under Section 6. Are there any side effects?

### **3. What if I am taking other medicines?**

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and PHARMACOR DABIGATRAN may interfere with each other. These include:

- aspirin, salicylates or other NSAID (anti-inflammatory) medicines
- medicines used to thin your blood (such as warfarin, unfractionated heparins, heparin derivatives (fondaparinux and desirudin), low molecular weight heparins (enoxaparin), clopidogrel, tirofiban, bivalirudin, prasugrel, ticagrelor, eptifibatide, ticlopidine, dextran, sulfapyrazone, rivaroxaban and apixaban)
- amiodarone, dronedarone, medicines used to treat irregular heartbeats
- verapamil, a calcium channel blocker used to treat high blood pressure and angina
- quinidine, a medicine used to treat malaria and irregular heartbeats
- clarithromycin or rifampicin, medicines used to treat infections
- lopinavir, nelfinavir, ritonavir, tipranavir or saquinavir, medicines used to treat HIV infections
- cyclosporin or tacrolimus, medicines used to help the body's immune system
- selective serotonin re-uptake inhibitors (SSRI) (e.g. citalopram, escitalopram, fluoxetine), selective serotonin norepinephrine re-uptake inhibitors (SNRI) (e.g. duloxetine, venlafaxine, desvenlafaxine), medicines used to treat mood disorders
- herbal medicines derived from St John's wort (*Hypericum perforatum*)
- carbamazepine, a medicine used to treat fits or convulsions
- medicines used to treat reflux and stomach ulcers (such as pantoprazole and ranitidine).

These medicines may be affected by PHARMACOR DABIGATRAN or may affect how well it works. You may need different amounts of

your medicines, change the timing of your medicine-taking routine or take different medicines. Your doctor or pharmacist will advise you.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking PHARMACOR DABIGATRAN.

### **4. How do I take PHARMACOR DABIGATRAN?**

**Follow the instructions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

#### *How much to take*

#### **AFTER KNEE AND HIP REPLACEMENT SURGERY**

The recommended dose of PHARMACOR DABIGATRAN is 220 mg (2 capsules of 110 mg) taken as a single dose once daily.

Patients with moderately reduced kidney function (e.g. over 75 years) or patients taking certain medicines may have an increased risk of bleeding. The doctor may prescribe the lower dose of 150 mg once daily, taken as 2 capsules of PHARMACOR DABIGATRAN 75 mg.

#### **AFTER KNEE REPLACEMENT SURGERY:**

Treatment with PHARMACOR DABIGATRAN should be started within 1 - 4 hours of completed surgery, using a single capsule of 110 mg and continuing with 2 capsules of 110 mg once daily for a total of 10 days.

#### **AFTER HIP REPLACEMENT SURGERY:**

Treatment with PHARMACOR DABIGATRAN should be started within 1 - 4 hours of completed surgery, using a single capsule of 110 mg and continuing with 2 capsules of 110 mg once daily for a total of 28 - 35 days.

If, within 4 hours after surgery, post-operative bleedings can still be observed, initiation of treatment should be delayed. If treatment is not

started on the day of surgery then treatment should be initiated with 2 capsules of 110 mg once daily.

**Follow the initiation instructions given to you by your doctor carefully.**

### **FOR STROKE PREVENTION IN PATIENTS WITH ATRIAL FIBRILLATION**

The recommended dose of PHARMACOR DABIGATRAN is 300 mg taken as 1 capsule of 150 mg in the morning and 1 capsule of 150 mg in the evening.

Patients over 75 years should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening. Patients with an increased risk of major bleeding (as determined by your doctor) should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening.

### **FOR THE TREATMENT OF BLOOD CLOTS AND PREVENTION OF BLOOD CLOTS RE-OCCURRING IN THE VEINS OF YOUR LEGS AND/OR LUNGS**

The recommended dose of PHARMACOR DABIGATRAN is 300 mg taken as 1 capsule of 150 mg twice a day following treatment with an injectable blood thinner for at least 5 days. To prevent blood clots re-occurring, continue on 1 capsule of 150 mg twice a day.

Patients over 75 years should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening. To prevent blood clots re-occurring, continue on 1 capsule of 110 mg twice a day.

Patients with an increased risk of major bleeding (as determined by your doctor) should take a lower dose of 220 mg, taken as 1 capsule of 110 mg in the morning and 1 capsule of 110 mg in the evening. To prevent blood clots re-occurring, continue on 1 capsule of 110 mg twice a day.


Your doctor will decide how long you need to be on this treatment for.

#### ***How to take it***


PHARMACOR DABIGATRAN is available in blister packs.

### **REMOVING PHARMACOR DABIGATRAN CAPSULES FROM THE BLISTER PACK**


Prior to removing a capsule from the blister card, separate one blister segment by tearing along the perforations (Figure A).



Once you have separated an individual blister segment, locate the tab marked with the arrow (Figure B).



Immediately before you are ready to take your dose of PHARMACOR DABIGATRAN, peel back the foil using the tab marked with the arrow until the capsule is fully visible (Figure C).



Turn the blister segment upside down and tip the capsule out, tapping the back of the blister segment, if necessary.

**Do not try to push the capsule through an unopened blister segment.**

**Do not cut the foil or use sharp instruments to remove the capsule from the blister.**

Capsules should always be stored in the sealed blister segments and only removed immediately before use. The capsule should be taken immediately after the foil over an individual blister segment is opened, or its effectiveness may be reduced.

**If additional capsules are inadvertently exposed to air, they should not be used and should be discarded.**

**Capsules should not be removed from the blister pack and repackaged in dose administration aids such as dosette boxes, tablet organisers or weekly medication packs.**

**Swallow the capsules whole with a full glass of water.**

**DO NOT CHEW OR OPEN THE CAPSULE. DO NOT SPRINKLE THE PELLETS ON FOOD OR MIX WITH LIQUIDS.**

This may cause an overdose of PHARMACOR DABIGATRAN and increase the risk of bleeding.

#### ***When to take it***

**Take PHARMACOR DABIGATRAN at about the same time each day.**

Taking your capsules at the same time each day will have the best effect. It will also help you remember when to take it. It does not matter if you take this medicine with or without food.

#### ***How long to take it***

**Continue taking your medicine for as long as your doctor tells you.**

PHARMACOR DABIGATRAN will continue to be prescribed while there is a risk of excessive clotting.

#### **AFTER KNEE REPLACEMENT SURGERY:**

This will usually be for a period of 10 days.

#### **AFTER HIP REPLACEMENT SURGERY:**

This will usually be for a period of 28 - 35 days.

**It is important to keep taking your medicine even if you feel well.**

If you stop using PHARMACOR DABIGATRAN before your doctor tells you to stop, you are at risk of developing a blood clot in a vein of your leg which can move to the lungs and be life-threatening.

**Tell your doctor immediately or go to Emergency at your nearest hospital if you notice swelling of the leg or cough and shortness of breath.**

These could be signs of a blood clot.

**Tell your doctor if you intend stopping treatment earlier.**

#### **FOR STROKE PREVENTION IN PATIENTS WITH ATRIAL FIBRILLATION:**

**It is important to keep taking your medicine even if you feel well.**

If you stop using PHARMACOR DABIGATRAN before your doctor tells you to stop, you are at risk of developing a blood clot. This can lead to serious health problems such as strokes.

#### **FOR TREATMENT AND PREVENTION OF BLOOD CLOTS RE-OCCURRING IN THE VEINS OF YOUR LEGS AND LUNGS:**

**It is important to keep taking your medicine even if you feel well.**

If you stop using PHARMACOR DABIGATRAN before your doctor tells you to stop, you are at risk of developing a blood clot. This can lead to serious health problems if those clots stop blood flowing normally.

#### ***If you forget to take it***

**After knee and hip replacement surgery continue with your remaining daily doses of PHARMACOR DABIGATRAN at the same time of the next day.**

**Do not take a double dose to make up for missed individual doses.**

**For stroke prevention in patients with atrial fibrillation a forgotten dose of PHARMACOR DABIGATRAN can still be taken up to 6 hours prior to the next dose.**

**A missed dose should be omitted if the remaining time is less than 6 hours prior to the next dose.**

**Do not take a double dose to make up for missed individual doses.**

**For the treatment and prevention of blood clots re-occurring in the veins of your legs and lungs a forgotten dose of PHARMACOR DABIGATRAN can still be taken up to 6 hours prior to the next dose.**

**A missed dose should be omitted if the remaining time is less than 6 hours prior to the next dose. Do not take a double dose to make up for missed individual doses.  
If you are not sure what to do, ask your doctor or pharmacist.**

**If you have trouble remembering when to take your medicine, ask your pharmacist for hints.**

*If you take too much (Overdose)*

**Immediately telephone your doctor or Poisons Information Centre (Australia 13 11 26) for advice, or go to Emergency at your nearest hospital if you think that you or anyone else may have taken too much PHARMACOR DABIGATRAN.**

**Do this even if there are no signs of discomfort or poisoning.**

If you take too much PHARMACOR DABIGATRAN you may have bleeding. Blood may be seen in stools or urine. Abnormal bruising may also be experienced.

## **5. What should I know while taking PHARMACOR DABIGATRAN?**

*Things you must do*

**Tell all doctors and pharmacists who are treating you that you are taking PHARMACOR DABIGATRAN.**

**Tell your doctor if, for any reason, you have not used PHARMACOR DABIGATRAN exactly as prescribed.**

Otherwise, your doctor may think that it was not effective and change your treatment unnecessarily.

**If you become pregnant while using PHARMACOR DABIGATRAN, tell your doctor immediately.**

**If you are going to have surgery, including dental surgery, tell your doctor or dentist that you are taking PHARMACOR DABIGATRAN.**

Your doctor may decide to temporarily stop your treatment with PHARMACOR DABIGATRAN

**Tell your doctor if you fall or injure yourself during treatment, especially if you hit your head, please seek urgent medical attention.**

You may need to be checked by a doctor, as you may be at increased risk of bleeding.

*Things you must not do*

**Do not give PHARMACOR DABIGATRAN to anyone else, even if they have the same condition as you.**

**Do not stop taking your medicine or lower the dosage without checking with your doctor.**

*Things to be careful of*

No studies on the effects of PHARMACOR DABIGATRAN on the ability to drive and operate machinery have been performed.

Driving or operating machinery should be avoided for a period of time after orthopaedic surgery.

## **6. Are there any side effects?**

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking PHARMACOR DABIGATRAN.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not.

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

**Ask your doctor or pharmacist to answer any questions you may have.**



### Less serious side effects

| Less serious side effects  | What to do   |
|--|--|
| <ul style="list-style-type: none"><li>• bruising</li><li>• nose bleeds</li><li>• stomach ache</li><li>• itchy skin, rash</li><li>• diarrhoea</li><li>• indigestion</li><li>• feeling sick</li><li>• cough</li><li>• painful, swollen joints</li><li>• sore nasal passages and throat</li><li>• discomfort when swallowing. .</li><li>• hair loss</li><li>• frequent infections such as fever, severe chills, sore throat or mouth ulcers (signs of lack of white blood cells).</li></ul> | <b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b> |

### Serious side effects

| Serious side effects   | What to do  |
|--|---|
| <ul style="list-style-type: none"><li>• long or excessive bleeding exceptional weakness</li><li>• tiredness, headaches, dizziness and looking pale (signs of anaemia)</li><li>• chest pain or being short of breath</li><li>• swelling of hands, ankles and feet</li><li>• red or dark brown urine</li><li>• red or black bowel motions. symptoms related to a condition called anticoagulant-related nephropathy: blood in urine, reduced urine output, swelling of the legs, ankles and feet, increased time for blood to clot (high INR test values), heavy bleeding.</li></ul> | <b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b> |

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

### Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

### After using AKM Dabigatran

#### Storage

**Keep your capsules in the blister pack until it is time to take them.**

If you take them out of the blister pack they may not keep well.

**Keep PHARMACOR DABIGATRAN in a cool dry place where the temperature stays below 25°C.**

**Do not store PHARMACOR DABIGATRAN or any other medicine in the bathroom or near a sink.**

**Do not leave it in the car or on window sills.**

Heat and dampness can destroy some medicines.

**Keep PHARMACOR DABIGATRAN where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

#### Disposal

**If your doctor tells you to stop taking PHARMACOR DABIGATRAN or it has passed its expiry date, ask your pharmacist what to do with any that is left over.**

## 7. Product details

This medicine is only available with a doctor's prescription.

### *What it looks like*

PHARMACOR DABIGATRAN is the brand name of your medicine.

PHARMACOR DABIGATRAN is available in three strengths of capsules:

- PHARMACOR DABIGATRAN 75 mg - White to pale yellow pellets filled in HPMC capsule size "1" light cream opaque cap and light cream opaque body imprinted with "75" on body and "DAB" on cap in black ink.
- PHARMACOR DABIGATRAN 110 mg – White to pale yellow pellets filled in HPMC capsule size "0" light blue opaque cap and light blue opaque body imprinted with "110" on body and "Dab" on cap in black ink.
- PHARMACOR DABIGATRAN 150 mg - White to pale yellow pellets filled in HPMC capsule size "0EL" light blue opaque cap and light cream opaque body imprinted with "150" on body and "DAB" on cap in black ink.

PHARMACOR DABIGATRAN 75 mg, 110 mg and 150 mg are available in Alu-Alu dessicant blister packs of 10, 30 and 60 capsules and HDPE bottle pack with dessicant of 60 capsules.

### *Ingredients*

Active ingredient:

- PHARMACOR DABIGATRAN 75 mg - 75 mg dabigatran etexilate given as 86.48 mg dabigatran etexilate mesilate per capsule.
- PHARMACOR DABIGATRAN 110 mg - 110 mg dabigatran etexilate given as 126.83 mg dabigatran etexilate mesilate per capsule.
- PHARMACOR DABIGATRAN 150 mg - 150 mg dabigatran etexilate given as 172.95 mg dabigatran etexilate mesilate per capsule.

Inactive ingredients:

### **Capsule fill**

- Butylated hydroxytoluene
- Microcrystalline cellulose
- Citric acid monohydrate
- Hyprolose (Hydroxy Propyl Cellulose)
- Hypromellose phthalate
- Povidone
- Purified talc
- Tartaric Acid
- Macrogol 600 (Polyethylene Glycol 600)
- Opadry White YS-1-7040
- Opadry 200 White (200F280000)

### **HPMC capsule shell**

- Hypromellose
- Titanium dioxide
- Sunset yellow FCF (for 75 mg, 110 mg and 150 mg strengths)
- Brilliant blue FCF (110 mg and 150 mg strength)
- Allura red AC (110 mg and 150 mg strength)

### **Printing ink**

- TekPrint™ SW-9008 Black Ink
- TekPrint™ SW-9009 Black Ink

PHARMACOR DABIGATRAN does not contain gluten, sucrose or tartrazine.

### **Who distributes PHARMACOR DABIGATRAN**

PHARMACOR PTY LTD  
Chatswood NSW 2067  
Australia

#### **Australian Registration Numbers:**

PHARMACOR DABIGATRAN

PHARMACOR DABIGATRAN 75 mg:

301981 - PHARMACOR DABIGATRAN dabigatran etexilate (as mesilate) 75 mg capsule bottle pack

301984 - PHARMACOR DABIGATRAN dabigatran etexilate (as mesilate) 75 mg capsule blister pack

PHARMACOR DABIGATRAN 110 mg:

301987 - PHARMACOR DABIGATRAN  
dabigatran etexilate (as mesilate) 110 mg capsule  
blister pack

301988 - PHARMACOR DABIGATRAN  
dabigatran etexilate (as mesilate) 110 mg capsule  
bottle pack

PHARMACOR DABIGATRAN 150 mg:

301982 - PHARMACOR DABIGATRAN  
dabigatran etexilate (as mesilate) 150 mg capsule  
blister pack

301985 - PHARMACOR DABIGATRAN  
dabigatran etexilate (as mesilate) 150 mg capsule  
bottle pack

**This leaflet was prepared in 04/2024.**