1. The probabil Sunday is:	ity of a leap ye	ear selected at ra	andom contain 53
	(b) 1/7	(c) 2/7	(d) 53/365
			marble is drawn at
random. The pro			
-	_	(c) 0/5	
			0.85. What is the
probability that i	-		
			(d) none of these
			ted from the numbers
(1, 2, 3,,1	-		
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3
5. What are the			
(a) 4	(b) 5	(c) 8	(d) 7
			ed at random from the
numbers (1,2,3,	35) is:		
(a) 12/35	(b) 11/3	(c) 13/35	(d) none of these
7. The sum of the			
(a) 2	(b) 1 (c	e) 0 (d) nor	ne of these.
8. The following	g probabilities	are given; choo	se the correct answer
for that which is	not possible.		
(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.
9. If three coins	are tossed sin	nultaneously, th	an the probability of
getting at least t			
		(c) $\frac{1}{2}$	
10. A letter is c			
ASSASSINAT	ION�. The pro	bability that the	eletter chosen has:
(a) 6/1 <mark>3</mark>	(b) 7/13	(c) 1	(d) none of these.
	-	•	tting an even number.
(A) 2/3	(B) 1	(C) 5/6	(D) 1/2
10 T			
		e same time. Fil	nd the probability of
getting both hea		(D) 0	
(A) 3/4 (B) 1/	4 (C) 1/2	(D) 0	
13. Two dice are sum of 9 is:	thrown simul	taneously. The բ	probability of getting a

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	4/9					
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.									
(A) 3/4	(B) 27/50	(C) 1/ <mark>4</mark>	(D) 29/100					
of drawing	15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:								
(A) 5	(B) 10	(C) 15	(D) 20						
	t random fron ive bulb is:	ontains 12 defention this box. The							
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100									
18. What is (A) 1/7	s the probabili (B) 53/366	ity of getting 55 (C) 2/ <mark>7</mark>		in a leap year? 7/366					
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13									
20. A game	e of chance co ly to come to ,then the prob	onsists of spin rest pointing to	ning an arro o one of the	number an odd number is:					
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4									

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:									
	(B) 31/365	_	(D) 1/133225						
2. Then the pro	x is chosen at raphability that x² < 2/5 (C) 3/5	< 2 is?	ımbers -2, -1, 0 , 1,						
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7									
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$									
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?									
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$						
27. Runs scored by batsman in 5 one day matches are 50, 70, 82,									
	e standard devia b) 25.49	c) 25.29	d) 25.69						
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.									
a) 13, 15		c) 18, 15	d) 13, 16						
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is									
a) $\frac{1}{2}$	b) $\frac{1}{3}$	c) $\frac{1}{4}$	d) $\frac{1}{6}$						
		d 3. The value of I c) 27 d)	• •						
	31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?								

a) 3	b) 4	c) 5	d) 7					
32.Out of the following values, which one is not possible in probability? a) $P(x) = 1$ b) $\sum x P(x) = 3$ c) $P(x) = 0.5$ d) $P(x) = -0.5$								
, , ,								
33.lf E(x) = a) 2	2 and E(z) = 4 , b) 6	c) 0		fficient data				
34.The cov	ariance of two	independen	t random variable	e is				
a) 1	b) 0	c) - 1	d) Und	efined				
35.If Σ P(x) a) 0	b) 1	the value of		ufficient data				
36.If P(x) = a) 1	0.5 and x = 4, t b) 0.5	t hen E(x) = 1 c) 4	d) <mark>2</mark>					
37.In a disc is always?	rete probability	y distributio	n, the sum of all p	probabilities				
•	b) Infinite	c) 1	d) Unde	efined				
=	obability of hit	ting the tar	get is 0.4, find me	ean and				
variance. a) 0.4, 0.2 <mark>4</mark>	b) 0.6, 0).24	c) 0.4, 0.16	d) 0.6, 0.16				
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6								
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 								

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0										
	Variance (X)				given by 2) - (E(X))2	d) (E(X))2			
43.Mean of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2) - (E(X))2$ d) $(E(X))2$										
44.N a) 0	44.Mean of a constant 'a' is a) 0									
45.Variance of a constant 'a' is . a) 0										
46.F	46. Find the mean and variance of X?									
	Х	0	1	2	3	4				
	f(x)	1/9	2/9	3/9	2/9	1/9				
a) 2, 4/3 b) 3, 4/3		3, 4/3	c) 2, 2/3			d) 3, 2/3				
47.F	47.Find the expectation of a random variable X?									

	Х	0	1	2	3		
	f(x)	1/6	2/6	2/6	1/6		
a) ().5		b) 1.5		c) 2	2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**
- a) \sqrt{np}

- b) \sqrt{pq} c) (np)2 d) \sqrt{npq}