1. Clinical Question & Core Hypothesis

Clinical Question

TLS-like lymphoid aggregates are often associated with improved immunotherapy response, yet many TLS-rich bladder tumors fail to benefit—what intrinsic tumor programs modulate whether local immune organization translates into effective anti-tumor immunity? 2) Tier 1 paths used: I built on MSH6 (MMR/MSI axis) because loss/defect increases neoantigen burden and should enhance immune recognition, and on eIF4E (translation control) because altered cap-dependent translation can selectively upregulate immune-evasive or antigen-processing–suppressive proteins; together they provide opposing forces (neoantigen supply vs. immune-evasion at the proteome level) that plausibly determine whether TLSs are effective. 3) Tier 2 evidence that supports the claim: metabolic state (HDL-C) can modulate the tumor immune microenvironment and affect lymphoid organization/function; DPYD and epigenetic markers (SDC2/TFPI2) point to additional metabolic/pharmacogenomic and epigenetic contexts that may modify immune visibility; AI-based biomarker accessibility supports combining histology (TLS) with molecular readouts for clinical stratification. 4) Why this endpoint: I selected therapy_response (immunotherapy response proxy) because the mechanistic tension between neoantigen generation (MSH6/MSI) and proteome-level immune evasion (eIF4E) most directly predicts susceptibility to checkpoint blockade and is the highest-impact, actionable clinical question emerging from the tier-1 synthesis. 5) What I excluded and why: I deliberately excluded exploratory Tier 3 signals (SARIFA, ER/TP53 links, CSF AD markers) from the core hypothesis because they are indirect or low-confidence for bladder-cancer immunobiology; I also limited anchors to two (MSH6, eIF4E) to keep the mechanistic question sharp—other Tier 2 entities were relegated to secondary_modifiers for follow-up rather than included in the core claim?

Working Hypothesis

TLS-like lymphoid aggregates in bladder tumors predict superior response to immune checkpoint blockade predominantly when the tumor harbors mismatch repair–deficiency-driven neoantigen load (MSH6 deficiency/MSI-H) and when eIF4E-mediated translation is low, limiting expression of immune-evasive proteins and preserving antigen presentation. This defines a two-axis biomarker model (MSH6/MSI status × eIF4E activity) with a predicted hierarchy of responses: TLS-high, MSH6-deficient, low-eIF4E tumors show the strongest therapy_response; TLS-high with MSH6-deficient but high eIF4E activity or TLS-high with MSH6-proficient but low eIF4E activity show intermediate responses; TLS-high with MSH6-proficient and high eIF4E activity, or TLS-low tumors, show attenuated responses. Anchors MSH6 and eIF4E jointly determine whether TLS-associated local immunity translates into durable anti-tumor activity, enabling a clinically actionable stratification that combines histology with molecular and translational readouts.

Element	Value
Imaging feature	TLS-like lymphoid aggregates
Anchor gene(s)	MSH6, eIF4E
Primary endpoint	therapy_response

2. Why This Gene — Not Others

Primary hypothesis gene(s)

MSH6: MSH6 (MMR/MSI axis) because loss/defect increases neoantigen burden and should enhance immune recognition, and on eIF4E (translation control) because altered cap-dependent translation can selectively upregulate immune-evasive or antigen-processing–suppressive proteins; together they provide opposing forces (neoantigen supply vs.

eIF4E: eIF4E (translation control) because altered cap-dependent translation can selectively upregulate immune-evasive or antigen-processing–suppressive proteins; together they provide opposing forces (neoantigen supply vs.

Alternative / secondary genes considered

Tier	Gene	Role / Reason excluded from core
Secondary	HDL-C	Supporting modifier from knowledge graph
Secondary	DPYD	Supporting modifier from knowledge graph
Secondary	SDC2	Supporting modifier from knowledge graph
Secondary	TFPI2	Supporting modifier from knowledge graph
Secondary	MSI/dMMR	Supporting modifier from knowledge graph
Secondary	AI_based_biomarkers	Supporting modifier from knowledge graph

3. Clinically Testable Predictions

Prediction 1: retrospective analysis of bladder cancer cohorts treated with ICIs, assessing TLS density, MSH6/MSI status, and proxies of eIF4E activity (e.g., p-eIF4E levels or translation-signature scores);

Prediction 2: integrate secondary modifiers (HDL-C, DPYD, SDC2, TFPI2) as exploratory covariates;

Prediction 3: validate in prospective datasets or organoid/PDX models engineered to recapitulate TLS and MSI/MMR status;

4. Minimum Viable Validation Plan

A. Retrospective Cohort Analysis

Data source: TCGA-BLCA (Firehose Legacy), ~400 patients - Cox PH model with interaction term for the imaging feature and anchor gene - Kaplan-Meier 2x2 stratification

B. Tissue-Level Validation

Methods: Multiplex IF / IHC / spatial transcriptomics

Key readouts: - Immune structure markers (e.g., CD20, CD3, PNAd) - Effector cell localization (CD8) relative to tumor nests - Anchor gene protein / phospho-protein expression - Stromal features (collagen density by SHG, if applicable)

Goal: Confirm that the proposed mechanism operates at the spatial / tissue level.

C. Functional Validation

Models: Ex vivo organotypic cultures, 3D co-culture, syngeneic mouse models

Perturbations: Genetic (KO/rescue) or pharmacologic modulation of anchor gene

Readouts: T-cell infiltration, cytotoxicity (granzyme B / IFN-gamma), tumor growth / survival

Goal: Establish causal relationship between anchor gene activity and immune efficacy.

5. Strengths, Caveats & Alternative Directions

Strengths

- Novel mechanistic hypothesis derived from knowledge graph analysis.

Caveats & Risks

- Hypothesis is computationally derived and requires experimental validation.

Alternative Directions

- HDL-C: Could serve as an alternative mechanistic axis if the primary hypothesis is not supported.
- DPYD: Could serve as an alternative mechanistic axis if the primary hypothesis is not supported.