

Espace Étale of a Presheaf

Sahil Karawade

Remark: This is actually given as a exercises question in Hartshorne.

Let X be the topological space and \mathcal{F} be the presheaf on X .

Define topological space $Sp\acute{e}(\mathcal{F})$, called espace étalé of \mathcal{F} as follows:

As a set $Sp\acute{e}(\mathcal{F}) = \bigcup_{p \in X} \mathcal{F}_p$ and define projection map $\pi : Sp\acute{e}(\mathcal{F}) \rightarrow X$ given by $s_p \mapsto p$. For each open set $U \subseteq X$ and each section $S \in \mathcal{F}(U)$ we obtain $\bar{s} : U \rightarrow Sp\acute{e}(\mathcal{F})$ given by $p \mapsto s_p = [U, s] \in \mathcal{F}_p$. Then we have $\pi \circ \bar{s} = id_U$. We make $Sp\acute{e}(\mathcal{F})$ into topological space by giving it the strongest topology such that $\bar{s} : U \rightarrow Sp\acute{e}(\mathcal{F})$ for all U . and all $s \in \mathcal{F}(U)$ are continuous.

Denote $\mathcal{O}(X)$ to be the set of all open subsets of X then $\bar{s}(U)$ form the basic open subset of $Sp\acute{e}(\mathcal{F})$ for $U \in \mathcal{O}(X)$: fix $x \in X$ and $s_x \in Sp\acute{e}(\mathcal{F})$ then $s_x \in \mathcal{F}$ so there exists $U \subseteq X$ such that $x \in U$ and $s \in \mathcal{F}(U) \implies s_x \in \bar{s}$.

Therefore $\{\bar{s}(U)\}_{U \in \mathcal{O}(X)}$ cover $Sp\acute{e}(\mathcal{F})$. Now, take $u_x \in \bar{s}(U) \cap \bar{t}(V)$ which will imply that $u_x = s_x$ and $u_x = t_x$ and hence there will exist $W \subset U \cap V$ with $x \in W$ and $s|_W = u|_W = t|_W \in \mathcal{F}(W)$. So we have the following maps $\bar{s}|_W : W \rightarrow Sp\acute{e}(\mathcal{F})$ and similarly maps $\bar{u}|_W$ and $\bar{t}|_W$. From above we have $\overline{s|_W}(W) = \overline{u|_W}(W) = \overline{t|_W}(W)$. For $x \in W$ and $u|_W \in \mathcal{F}(W)$, $(u|_W)_x \in \overline{u|_W}(W)$. Therefore $u_x = (u|_W)_x \in \overline{u|_W}(W) = \overline{t|_W}(W) = \overline{s|_W}(W) \subset \bar{s}(U)$ and similarly $u_x = (u|_W)_x \in \overline{u|_W}(W) = \overline{s|_W}(W) = \overline{t|_W}(W) \subset \bar{t}(V)$. Hence $u_x \in \bar{s}(U) \cap \bar{t}(V)$. Also, $\bar{s} : U \rightarrow Sp\acute{e}(\mathcal{F})$ is continuous since for $\bar{t}(V)$ a basic open subset of $Sp\acute{e}(\mathcal{F})$ we have $\bar{s}^{-1}(\bar{t}(V)) = \{p \in U | \bar{s}(p) \in \bar{t}(V)\} = \{p \in U \cap V | s_p = t_p\}$ which is open: let $x \in \bar{s}^{-1}(\bar{t}(V))$ then $s_p = t_p$ and by definition there exists $W \subseteq U \cap V$ with $x \in W$ and $s|_W = t|_W$. Therefore $W \subseteq \bar{s}^{-1}(\bar{t}(V))$.

Inclusion, i , of $\bar{s}(U)$ in $Sp\acute{e}(\mathcal{F})$ is continous and π is continuous so $\pi \circ i$ is continuous.

Definition 1 (Local homeomorphism) $f : X \rightarrow Y$ is continuous then for all $x \in X$ there exists open set U containing x such that $f(U)$ is open in Y and $f|_U : U \rightarrow f(U)$ is homeomorphism.

Now we show that $\pi : Sp\acute{e}(\mathcal{F}) \rightarrow X$ is local homeomorphism. Let $s_p \in Sp\acute{e}(\mathcal{F})$ then there exists $U \in \mathcal{O}(X)$ and $s \in \mathcal{F}(U)$. By definition $s_p = [U, s]$ with $p \in U$ and $s \in \mathcal{F}(U)$ and $\pi(\bar{s}(U)) = U$ is open. To show: $\pi|_{\bar{s}(U)} : \bar{s}(U) \rightarrow \pi(\bar{s}(U))$ is homeomorphism.

$\pi|_{\bar{s}(U)} : \bar{s}(U) \rightarrow U$ is continuous and $\bar{s} : U \rightarrow Sp\acute{e}(\mathcal{F})$ is continuous. Therefore $\pi|_{\bar{s}(U)}$ has continuous inverse given by \bar{s} . Hence π is local homeomorphism and \bar{s} is a section of $Sp\acute{e}(\mathcal{F})$.

Now, $\mathcal{F}^+(U) = \{\bar{s} : U \rightarrow Sp\acute{e}(\mathcal{F}) | \bar{s}(x) = s_x \text{ for all } s \in \mathcal{F}(U)\}$.

For open subset U define $\varphi(U) : \mathcal{F}(U) \rightarrow \mathcal{F}^+(U)$ by $s \mapsto \bar{s}$.

Consider the following construction, let \mathcal{F} and \mathcal{G} be presheaves on X . $\psi : \mathcal{F} \rightarrow \mathcal{G}$ be morphism of presheaves. If $\sigma \in \mathcal{F}^+(U)$ and x is point in U then $\sigma(x) \in \mathcal{F}_x$, that is $\sigma(x) = [V, s]$ for $s \in \mathcal{F}(V)$. Define $\sigma'(x) = (\psi_V(s))_x$ then $\sigma' : U \rightarrow Sp\acute{e}(\mathcal{G})$ is well defined and continuous such that $\pi \circ \sigma' = id_U$. Define $\psi_U^+(\sigma) = \sigma'$. This gives $\psi_U^+ : \mathcal{F}^+(U) \rightarrow \mathcal{G}^+(U)$ morphism of sheaves and ψ is unique since sheafification does not alter stalks. If \mathcal{G} is a sheaf then we get that \mathcal{F}^+ is the sheafification of \mathcal{F} .